

<p style="text-align: center;"><b>Table of Contents</b></p> <p style="text-align: center;"><b>Section II – County Soil and Site Information Reports</b></p>
---

	<b>Issue Date</b>	<b>Date of Last Review</b>	<b>Responsible Staff</b>
<i>Use and Explanation of Soil Interpretations</i>			
<i>Explanation of Key Phrases Used in Soil Interpretations</i>			
<b>Soils Legends</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Acreage and Proportionate Extent of the Soils</i>			
<b>Soil Descriptions - Nontechnical</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>Use and Explanation of Nontechnical Descriptions</i>			
<i>*Nontechnical Soils Description Report</i>			
<b>Soil Descriptions - Technical</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Map Unit Description Report</i>			
<b>Cropland Interpretations - Technical</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Land Capability and Yields Per Acre of Crops</i>			
<i>*Prime Farmland Report</i>			
<i>*Soil Rating for Plant Growth Index</i>			
<i>*Soil Properties for Conservation Planning</i>			
<b>Rangeland, Grazed Forestland, Native Pastureland Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Rangeland Productivity Report</i>			
<i>*Range Site Descriptions</i>			
<b>Forestland Interpretations</b>	<b>1/93</b>	<b>1/93</b>	<b>SOI</b>
<i>Use and Explanation of Forestland Interpretations</i>			
<i>*Woodland Management and Productivity</i>			
<b>Nonagricultural Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Building Site Development Report</i>			
<i>*Construction Materials Report</i>			
<b>Recreation Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Recreational Interpretations</i>			
<b>Wildlife Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Wildlife Interpretations Report</i>			

<b>Pastureland and Hayland Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Yields Per Acre of Pasture and Hayland</i>			
	<b>Issue Date</b>	<b>Date of Last Review</b>	<b>Responsible Staff</b>
<b>Mined Land Interpretations</b>	<b>1/93</b>	<b>1/93</b>	<b>SOI</b>
<i>Use and Explanation of Mined Land Interpretations</i>			
<b>Windbreak Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Conservation Tree and Shrub Management Report</i>			
<b>Engineering Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Engineering Index Properties</i>			
<i>*Physical Properties of the Soils</i>			
<i>*Chemical Properties of the Soils</i>			
<i>*Water Features</i>			
<i>*Soil Features</i>			
<i>*Water Management Report</i>			
<b>Waste Disposal Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>*Sanitary Facilities Report</i>			
<i>*Agricultural Waste Management Report</i>			
<b>Hydric Soil Interpretations</b>	<b>4/02</b>	<b>4/02</b>	<b>SOI</b>
<i>Use and Explanation of Hydric Soil Interpretations</i>			
<i>*Hydric Soils List</i>			
<b>HEL Interpretations</b>	<b>See Report</b>		<b>SOI</b>
<i>Use and Explanation of Highly Erodible Land Interpretations</i>			
<i>*Highly Erodible Lands Report</i>			
1	<i>Highly Erodible Land</i>		
2	<i>Potentially Highly Erodible Land</i>		
3	<i>Nonhighly Erodible Land</i>		

*\*County specific computer generated reports.*

## ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS

Brown County, Nebraska: Published

Map symbol	Soil name	Acres	Percent
Aa	Almeria Loamy Fine Sand, 0 To 2 Percent Slopes-----	544	*
Ae	Almeria Fine Sandy Loam, 0 To 2 Percent Slopes-----	737	*
Af	Almeria-Histosols Complex, Channeled-----	1,799	0.2
An	Anselmo Fine Sandy Loam, 0 To 2 Percent Slopes-----	3,376	0.4
AnC	Anselmo Fine Sandy Loam, 2 To 6 Percent Slopes-----	4,355	0.6
AnD	Anselmo Fine Sandy Loam, 6 To 11 Percent Slopes-----	915	0.1
AtF	Anselmo-Brunswick Fine Sandy Loams, 11 To 30 Percent Slopes-----	2,776	0.4
Ba	Barney Fine Sandy Loam, Channeled-----	1,380	0.2
Bd	Bolent Fine Sandy Loam, 0 To 2 Percent Slopes-----	805	0.1
Bo	Brocksburg Loam, 0 To 1 Percent Slopes-----	1,116	0.1
BrD	Brunswick Fine Sandy Loam, 3 To 9 Percent Slopes-----	843	0.1
DuB	Dunday Loamy Fine Sand, 0 To 3 Percent Slopes-----	5,218	0.7
DuD	Dunday Loamy Fine Sand, 3 To 9 Percent Slopes-----	4,319	0.6
Eo	Els Loamy Sand, 0 To 2 Percent Slopes-----	4,191	0.5
EpB	Els-Ipage Fine Sands, 0 To 3 Percent Slopes-----	19,823	2.5
Es	Elsmere Loamy Fine Sand, 0 To 2 Percent Slopes-----	2,784	0.4
Fe	Fluvaquents, Sandy-----	286	*
Gn	Gannett Fine Sandy Loam, 0 To 2 Percent Slopes-----	1,825	0.2
IdB	Inavale Loamy Fine Sand, 0 To 3 Percent Slopes-----	2,573	0.3
IkB	Inavale Sand, Channeled-----	1,013	0.1
In	Inavale-Barney Complex, Channeled-----	2,890	0.4
IpB	Ipage Fine Sand, 0 To 3 Percent Slopes-----	11,372	1.5
IsB	Ipage Loamy Sand, 0 To 3 Percent Slopes-----	2,794	0.4
Jn	Jansen Fine Sandy Loam, 0 To 2 Percent Slopes-----	2,635	0.3
JnC	Jansen Fine Sandy Loam, 2 To 6 Percent Slopes-----	2,370	0.3
Jo	Jansen Loam, 0 To 2 Percent Slopes-----	4,005	0.5
JoC	Jansen Loam, 2 To 6 Percent Slopes-----	4,279	0.5
Jr	Jansen-Meadin Complex, 0 To 2 Percent Slopes-----	3,037	0.4
JtB	Jansen-Sandose Complex, 0 To 3 Percent Slopes-----	1,529	0.2
Jw	Johnstown Fine Sandy Loam, 0 To 1 Percent Slopes-----	3,943	0.5
JwB	Johnstown Fine Sandy Loam, 1 To 3 Percent Slopes-----	2,578	0.3
Jy	Johnstown Loam, 0 To 1 Percent Slopes-----	15,746	2.0
JyB	Johnstown Loam, 1 To 3 Percent Slopes-----	3,902	0.5
JyC	Johnstown Loam, 3 To 6 Percent Slopes-----	1,069	0.1
LcG	Labu-Sansarc Silty Clays, 11 To 40 Percent Slopes-----	574	*
LfB	Libory Loamy Fine Sand, 0 To 3 Percent Slopes-----	3,517	0.4
Lo	Loup Fine Sandy Loam, 0 To 2 Percent Slopes-----	7,997	1.0
Lp	Loup Fine Sandy Loam, Wet, 0 To 2 Percent Slopes-----	2,105	0.3
LtB	Loup-Els Complex, 0 To 3 Percent Slopes-----	6,080	0.8
Ma	Marlake Fine Sandy Loam, 0 To 1 Percent Slopes-----	3,048	0.4
McG	Mckelvie-Tassel-Ronson Complex, 15 To 70 Percent Slopes-----	25,949	3.3
MeB	Meadin Sandy Loam, 0 To 3 Percent Slopes-----	1,128	0.1
MeF	Meadin Sandy Loam, 3 To 30 Percent Slopes-----	3,487	0.4
Oe	O'Neill Fine Sandy Loam, 0 To 2 Percent Slopes-----	3,715	0.5
On	O'Neill Loam, 0 To 2 Percent Slopes-----	1,052	0.1
OsC	O'Neill-Meadin Sandy Loams, 2 To 6 Percent Slopes-----	3,451	0.4
OsD	O'Neill-Meadin Sandy Loams, 6 To 11 Percent Slopes-----	2,627	0.3
Pg	Pits, Sand And Gravel-----	440	*
PtB	Pivot Loamy Sand, 0 To 3 Percent Slopes-----	3,060	0.4
RtB	Ronson-Tassel Fine Sandy Loams, 0 To 3 Percent Slopes-----	1,822	0.2
RtC	Ronson-Tassel Fine Sandy Loams, 3 To 6 Percent Slopes-----	2,246	0.3
RtD	Ronson-Tassel Fine Sandy Loams, 6 To 11 Percent Slopes-----	945	0.1
ScB	Sandose Loamy Fine Sand, 0 To 3 Percent Slopes-----	4,867	0.6
SkB	Simeon Loamy Sand, 0 To 3 Percent Slopes-----	5,844	0.7
SkD	Simeon Loamy Sand, 3 To 9 Percent Slopes-----	2,802	0.4
SvD	Simeon-Valentine Fine Sands, 0 To 9 Percent Slopes-----	19,245	2.5
Tn	Tryon Loamy Fine Sand, 0 To 2 Percent Slopes-----	6,059	0.8
To	Tryon Loamy Fine Sand, Wet, 0 To 2 Percent Slopes-----	1,869	0.2
TpB	Tryon-Els Complex, 0 To 3 Percent Slopes-----	9,460	1.2
VaB	Valentine Fine Sand, 0 To 3 Percent Slopes-----	2,697	0.3
VaD	Valentine Fine Sand, 3 To 9 Percent Slopes-----	40,105	5.1
VaE	Valentine Fine Sand, Rolling-----	210,376	26.9
VaF	Valentine Fine Sand, Rolling And Hilly-----	169,412	21.7
VbB	Valentine Loamy Fine Sand, 0 To 3 Percent Slopes-----	1,890	0.2
VbD	Valentine Loamy Fine Sand, 3 To 9 Percent Slopes-----	23,963	3.1
VfD	Valentine-Els Fine Sands, 0 To 9 Percent Slopes-----	49,684	6.4
VhD	Valentine-Libory Complex, 0 To 9 Percent Slopes-----	4,426	0.6
VpD	Valentine-Pivot Complex, 0 To 9 Percent Slopes-----	4,254	0.5
VrD	Valentine-Sandose Loamy Fine Sands, 0 To 9 Percent Slopes-----	4,386	0.6
VsG2	Valentine-Simeon Complex, 9 To 40 Percent Slopes, Eroded-----	6,870	0.9
VtE	Valentine-Tassel Complex, 3 To 17 Percent Slopes-----	2,284	0.3
VwE	Valentine-Tryon Complex, 0 To 17 Percent Slopes-----	8,645	1.1
VxB	Vetal Loam, 1 To 3 Percent Slopes-----	1,625	0.2
zwa	Water > 40 Acres-----	4,281	0.5
zwb	Water < 40 Acres-----	614	*
	Total-----	781,728	100.0

\* Less than 0.1 percent.



# NONTECHNICAL SOIL DESCRIPTIONS Brown County, Nebraska

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the NonTechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

## Aa Almeria Loamy Fine Sand, 0 To 2 Percent Slopes

Almeria soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Wet Subirrigated - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 5w.

## Ae Almeria Fine Sandy Loam, 0 To 2 Percent Slopes

Almeria soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Wet Subirrigated range site. It is in the nonirrigated land capability classification 5w.

## Af Almeria-Histosols Complex, Channeled

Almeria soil makes up 55 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level fen on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is very poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Wet Land range site. It is in the nonirrigated land capability classification 6w.

Histosols soil makes up 45 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level fen on valley. The runoff class is negligible. <parent material is missing> This soil is very poorly drained. The slowest permeability is rapid. It has a very high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. It is in the nonirrigated land capability classification 8w.

## An Anselmo Fine Sandy Loam, 0 To 2 Percent Slopes

Anselmo soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

## AnC Anselmo Fine Sandy Loam, 2 To 6 Percent Slopes

Anselmo soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hill on tableland, valley side on valley. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

## AnD Anselmo Fine Sandy Loam, 6 To 11 Percent Slopes

Anselmo soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hill on tableland, valley side on valley. The runoff class is low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued  
Brown County, Nebraska

AtF Anselmo-Brunswick Fine Sandy Loams, 11 To 30 Percent Slopes

Anselmo soil makes up 60 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately steep to steep summit, shoulder valley side on valley. The runoff class is medium. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 3 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Brunswick soil makes up 40 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a strongly sloping to steep backslope valley side on valley. The runoff class is medium. <parent material is missing> The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Ba Barney Fine Sandy Loam, Channeled

Barney soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is very poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 6 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Land range site. It is in the nonirrigated land capability classification 6w.

Bd Bolent Fine Sandy Loam, 0 To 2 Percent Slopes

Bolent soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is very low. The parent material consists of sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

Bo Brocksburg Loam, 0 To 1 Percent Slopes

Brocksburg soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level flat on tableland. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

BrD Brunswick Fine Sandy Loam, 3 To 9 Percent Slopes

Brunswick soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hill on tableland, valley side on valley. The runoff class is low. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

DuB Dunday Loamy Fine Sand, 0 To 3 Percent Slopes

Dunday soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on tableland, interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

DuD Dunday Loamy Fine Sand, 3 To 9 Percent Slopes

Dunday soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on tableland, interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

**Eo Els Loamy Sand, 0 To 2 Percent Slopes**

Els soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

**EpB Els-Ipage Fine Sands, 0 To 3 Percent Slopes**

Els soil makes up 65 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 6e.

Ipage soil makes up 35 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands over sandy alluvium. This soil is moderately well drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil is in the Sandy Lowland 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

**Es Elsmere Loamy Fine Sand, 0 To 2 Percent Slopes**

Elsmere soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy alluvium and/or eolian sands. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

**Fe Fluvaquents, Sandy**

Fluvaquents soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on valley. The runoff class is negligible. The parent material consists of silty alluvium. This soil is very poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It is in the nonirrigated land capability classification 8w.

**Gn Gannett Fine Sandy Loam, 0 To 2 Percent Slopes**

Gannett soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is poorly drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Wet Subirrigated range site. It is in the nonirrigated land capability classification 5w.

**IdB Inavale Loamy Fine Sand, 0 To 3 Percent Slopes**

Inavale soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

**IkB Inavale Sand, Channeled**

Inavale soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6w.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

In Inavale-Barney Complex, Channeled

Inavale soil makes up 60 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6w.

Barney soil makes up 40 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is very poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 6 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Land - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6w.

IpB Ipage Fine Sand, 0 To 3 Percent Slopes

Ipage soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping stream terrace on valley, hummock on interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands over sandy alluvium. This soil is moderately well drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil is in the Sandy Lowland 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

IsB Ipage Loamy Sand, 0 To 3 Percent Slopes

Ipage soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on interdune on sandhills, stream terrace on valley. The runoff class is negligible. The parent material consists of eolian sands over sandy alluvium. This soil is moderately well drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Jn Jansen Fine Sandy Loam, 0 To 2 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

JnC Jansen Fine Sandy Loam, 2 To 6 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping valley side on valley, hill on tableland. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Jo Jansen Loam, 0 To 2 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

JoC Jansen Loam, 2 To 6 Percent Slopes

Jansen soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hill on tableland, valley side on valley. The runoff class is medium. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.



NONTECHNICAL SOIL DESCRIPTIONS--Continued  
Brown County, Nebraska

---

Jr Jansen-Meadin Complex, 0 To 2 Percent Slopes

Jansen soil makes up 50 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping knoll on tableland. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Meadin soil makes up 50 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is very low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

JtB Jansen-Sandose Complex, 0 To 3 Percent Slopes

Jansen soil makes up 50 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is medium. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Sandose soil makes up 50 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping knoll on tableland. The runoff class is low. The parent material consists of eolian sands over loamy sediments. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Jw Johnstown Fine Sandy Loam, 0 To 1 Percent Slopes

Johnstown soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level flat on tableland. The runoff class is low. The parent material consists of loess over gravelly sand. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

JwB Johnstown Fine Sandy Loam, 1 To 3 Percent Slopes

Johnstown soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping hill on tableland. The runoff class is medium. The parent material consists of loess over gravelly sand. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Jy Johnstown Loam, 0 To 1 Percent Slopes

Johnstown soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level flat on tableland. The runoff class is low. The parent material consists of loess over gravelly sand. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

JyB Johnstown Loam, 1 To 3 Percent Slopes

Johnstown soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping hill on tableland. The runoff class is medium. The parent material consists of loess over gravelly sand. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

---

JyC Johnstown Loam, 3 To 6 Percent Slopes

Johnstown soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping hill on tableland, valley side on valley. The runoff class is medium. The parent material consists of loess over gravelly sand. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

LcG Labu-Sansarc Silty Clays, 11 To 40 Percent Slopes

Labu soil makes up 55 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a strongly sloping to steep backslope valley side on valley. The runoff class is very high. The parent material consists of residuum weathered from clayey shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a very low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Clayey - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Sansarc soil makes up 45 percent of the map unit. This map unit is in the Southern Rolling Pierre Shale Plains Major Land Resource Area. This soil occurs on a strongly sloping to steep summit, shoulder valley side on valley. The runoff class is very high. The parent material consists of clayey residuum weathered from shale. The soil is 4 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a very low available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Shallow Clay - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

LfB Libory Loamy Fine Sand, 0 To 3 Percent Slopes

Libory soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands over loess. This soil is moderately well drained. The slowest permeability is moderately slow. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 18 inches. This soil is in the Sandy Lowland 22-25" P.z. range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Lo Loup Fine Sandy Loam, 0 To 2 Percent Slopes

Loup soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is very low. The parent material consists of sandy alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 9 inches. This soil is in the Wet Subirrigated range site. It is in the nonirrigated land capability classification 5w.

Lp Loup Fine Sandy Loam, Wet, 0 To 2 Percent Slopes

Loup soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is very poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. This soil is in the Wet Land range site. It is in the nonirrigated land capability classification 5w.

LtB Loup-Els Complex, 0 To 3 Percent Slopes

Loup soil makes up 50 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is very low. The parent material consists of sandy alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 9 inches. This soil is in the Wet Subirrigated range site. It is in the nonirrigated land capability classification 5w.

Els soil makes up 50 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

Ma Marlake Fine Sandy Loam, 0 To 1 Percent Slopes

Marlake soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level depression on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over alluvium. This soil is very poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It is in the nonirrigated land capability classification 8w.

McG Mckelvie-Tassel-Ronson Complex, 15 To 70 Percent Slopes

Mckelvie soil makes up 50 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately steep to very steep backslope valley side on valley. The runoff class is low. The parent material consists of sandy residuum weathered from sandstone. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 7e.

Longpine soil makes up 30 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately steep to very steep shoulder valley side on valley. The runoff class is very high. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 7s.

MeB Meadin Sandy Loam, 0 To 3 Percent Slopes

Meadin soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland, knoll on tableland. The runoff class is very low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

MeF Meadin Sandy Loam, 3 To 30 Percent Slopes

Meadin soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to steep hill on tableland, valley side on valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

Oe O'Neill Fine Sandy Loam, 0 To 2 Percent Slopes

O'Neill soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is very low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

On O'Neill Loam, 0 To 2 Percent Slopes

O'Neill soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flat on tableland. The runoff class is low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2c.

OsC O'Neill-Meadin Sandy Loams, 2 To 6 Percent Slopes

O'Neill soil makes up 60 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hill on tableland, valley side on valley. The runoff class is very low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued  
Brown County, Nebraska

Meadin soil makes up 40 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hill on tableland, valley side on valley. The runoff class is very low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

OsD O'Neill-Meadin Sandy Loams, 6 To 11 Percent Slopes

O'Neill soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hill on tableland, valley side on valley. The runoff class is low. The parent material consists of coarse-loamy alluvium over sandy and gravelly alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Meadin soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hill on tableland, valley side on valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

PtB Pivot Loamy Sand, 0 To 3 Percent Slopes

Pivot soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on tableland, interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over gravelly alluvium. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

RtB Ronson-Tassel Fine Sandy Loams, 0 To 3 Percent Slopes

Ronson soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on tableland. The runoff class is very low. The parent material consists of residuum weathered from calcareous sandstone. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Longpine soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping knoll on tableland. The runoff class is medium. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

RtC Ronson-Tassel Fine Sandy Loams, 3 To 6 Percent Slopes

Ronson soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping backslope hill on tableland. The runoff class is very low. The parent material consists of residuum weathered from calcareous sandstone. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Longpine soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping summit, shoulder hill on tableland. The runoff class is medium. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

RtD Ronson-Tassel Fine Sandy Loams, 6 To 11 Percent Slopes

Ronson soil makes up 55 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping backslope hill on tableland. The runoff class is low. The parent material consists of residuum weathered from calcareous sandstone. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Longpine soil makes up 45 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping summit, shoulder hill on tableland. The runoff class is high. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

ScB Sandose Loamy Fine Sand, 0 To 3 Percent Slopes

Sandose soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping interdune on sandhills, swale on tableland. The runoff class is negligible. The parent material consists of eolian sands over loamy sediments. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

SkB Simeon Loamy Sand, 0 To 3 Percent Slopes

Simeon soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping interdune on sandhills, hummock on tableland. The runoff class is negligible. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel 17-25" P.z. range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

SkD Simeon Loamy Sand, 3 To 9 Percent Slopes

Simeon soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hill on tableland, valley side on valley. The runoff class is very low. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel 17-25" P.z. range site. It is in the nonirrigated land capability classification 6s.

SvD Simeon-Valentine Fine Sands, 0 To 9 Percent Slopes

Simeon soil makes up 60 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on tableland, interdune on sandhills. The runoff class is negligible. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel 17-25" P.z. range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

Valentine soil makes up 40 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping interdune on sandhills, dune on tableland. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Tn Tryon Loamy Fine Sand, 0 To 2 Percent Slopes

Tryon soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Subirrigated range site. It is in the nonirrigated land capability classification 5w.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

To Tryon Loamy Fine Sand, Wet, 0 To 2 Percent Slopes

Tryon soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is very poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Land range site. It is in the nonirrigated land capability classification 5w.

TpB Tryon-Els Complex, 0 To 3 Percent Slopes

Tryon soil makes up 55 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Subirrigated range site. It is in the nonirrigated land capability classification 5w.

Els soil makes up 45 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 6e.

VaB Valentine Fine Sand, 0 To 3 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

VaD Valentine Fine Sand, 3 To 9 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

VaE Valentine Fine Sand, Rolling

Valentine soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a strongly sloping to steep dune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. It is in the nonirrigated land capability classification 6e.

VaF Valentine Fine Sand, Rolling And Hilly

Valentine soil makes up 70 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a strongly sloping to steep dune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. It is in the nonirrigated land capability classification 6e.

Valentine soil makes up 30 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a steep to very steep dune on sandhills. The runoff class is low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Choppy Sands 22-25" P.z. range site. It is in the nonirrigated land capability classification 7e.

VbB Valentine Loamy Fine Sand, 0 To 3 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping interdune on sandhills, hummock on tableland. The runoff class is negligible. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued  
Brown County, Nebraska

VbD Valentine Loamy Fine Sand, 3 To 9 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping interdune on sandhills, dune on tableland. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

VfD Valentine-Els Fine Sands, 0 To 9 Percent Slopes

Valentine soil makes up 55 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Els soil makes up 45 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 6e.

VhD Valentine-Libory Complex, 0 To 9 Percent Slopes

Valentine soil makes up 60 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Libory soil makes up 40 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands over loess. This soil is moderately well drained. The slowest permeability is moderately slow. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 18 inches. This soil is in the Sandy Lowland 22-25" P.z. range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

VpD Valentine-Pivot Complex, 0 To 9 Percent Slopes

Valentine soil makes up 60 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Pivot soil makes up 40 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over gravelly alluvium. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

VrD Valentine-Sandose Loamy Fine Sands, 0 To 9 Percent Slopes

Valentine soil makes up 65 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Sandose soil makes up 35 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of eolian sands over loamy sediments. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Sandy 22-25" P.z. range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued  
Brown County, Nebraska

---

VsG2 Valentine-Simeon Complex, 9 To 40 Percent Slopes, Eroded

Valentine soil makes up 65 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a strongly sloping to steep backslope valley side on valley. The runoff class is low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. It is in the nonirrigated land capability classification 7e.

Simeon soil makes up 35 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a strongly sloping to steep summit, shoulder valley side on valley. The runoff class is low. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel 17-25" P.z. range site. It is in the nonirrigated land capability classification 6s.

VtE Valentine-Tassel Complex, 3 To 17 Percent Slopes

Valentine soil makes up 60 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Longpine soil makes up 40 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep backslope hill on sandhills. The runoff class is very high. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately rapid. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Shallow Limy - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6s.

VwE Valentine-Tryon Complex, 0 To 17 Percent Slopes

Valentine soil makes up 65 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on interdune on sandhills. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands 22-25" P.z. range site. It is in the nonirrigated land capability classification 6e.

Tryon soil makes up 35 percent of the map unit. This map unit is in the Nebraska Sand Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on interdune on sandhills. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is very poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Land range site. It is in the nonirrigated land capability classification 5w.

VxB Vetat Loam, 1 To 3 Percent Slopes

Vetat soil makes up 100 percent of the map unit. This map unit is in the Dakota-Nebraska Eroded Tableland Major Land Resource Area. This soil occurs on a gently sloping footslope swale on tableland, valley side on valley. The runoff class is very low. The parent material consists of loamy alluvium over eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.





**Aa—Almeria loamy fine sand, 0 to 2 percent slopes**

## Map Unit Composition

Almeria: 100 percent

## Component Descriptions

Almeria

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Flood plain on valley

Parent material: Sandy alluvium

Slope: 0 to 2 percent

Drainage class: Poorly drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 5.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: About 0 to 18 inches

Runoff class: Negligible

Ecological site: Wet Subirrigated - Veg. Zone 3

Land capability (nonirrigated): 5w

## Typical Profile:

H1—0 to 5 inches; loamy fine sand

H2—5 to 60 inches; stratified sand to fine sandy loam

Runoff class: Negligible

Ecological site: Wet Subirrigated

Land capability (nonirrigated): 5w

## Typical Profile:

H1—0 to 5 inches; fine sandy loam

H2—5 to 60 inches; stratified sand to fine sandy loam

**Af—Almeria-Histosols complex, Channeled**

## Map Unit Composition

Almeria: 55 percent

Histosols: 45 percent

## Component Descriptions

Almeria

MLRA: 65 - Nebraska Sand Hills

Landform: Flood plain on valley

Parent material: Sandy alluvium

Slope: 0 to 2 percent

Drainage class: Very poorly drained

Slowest permeability: Moderately rapid (About 1.98 in/hr)

Available water capacity: Low (About 5.8 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: About 0 to 0 inches

Runoff class: Negligible

Ecological site: Wet Land

Land capability (nonirrigated): 6w

## Typical Profile:

H1—0 to 5 inches; fine sandy loam

H2—5 to 60 inches; stratified sand to fine sandy loam

Histosols

MLRA: 65 - Nebraska Sand Hills

Landform: Fen on valley

Slope: 0 to 1 percent

Drainage class: Very poorly drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Very high (About 12.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: About 0 to 0 inches

Runoff class: Negligible

**Ae—Almeria fine sandy loam, 0 to 2 percent slopes**

## Map Unit Composition

Almeria: 100 percent

## Component Descriptions

Almeria

MLRA: 65 - Nebraska Sand Hills

Landform: Flood plain on valley

Parent material: Sandy alluvium

Slope: 0 to 2 percent

Drainage class: Poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Low (About 5.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: About 0 to 18 inches

*Land capability (nonirrigated): 8w*

*Typical Profile:*

H1—0 to 26 inches; mucky peat  
H2—26 to 60 inches; fine sand

## **An—Anselmo fine sandy loam, 0 to 2 percent slopes**

### **Map Unit Composition**

Anselmo: 100 percent

### **Component Descriptions**

Anselmo

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flat on tableland

*Parent material:* Loamy eolian deposits

*Slope:* 0 to 2 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 7.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated): 2e*

*Land capability (nonirrigated): 2e*

*Typical Profile:*

H1—0 to 7 inches; fine sandy loam  
H2—7 to 18 inches; fine sandy loam  
H3—18 to 33 inches; fine sandy loam  
H4—33 to 60 inches; loamy fine sand

### **Minor Components**

**Perched Wt**

## **AnC—Anselmo fine sandy loam, 2 to 6 percent slopes**

### **Map Unit Composition**

Anselmo: 100 percent

### **Component Descriptions**

Anselmo

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Loamy eolian deposits

*Slope:* 2 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 7.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated): 3e*

*Land capability (nonirrigated): 3e*

*Typical Profile:*

H1—0 to 7 inches; fine sandy loam  
H2—7 to 18 inches; fine sandy loam  
H3—18 to 33 inches; fine sandy loam  
H4—33 to 60 inches; loamy fine sand

### **Minor Components**

**Perched Wt**

## **AnD—Anselmo fine sandy loam, 6 to 11 percent slopes**

### **Map Unit Composition**

Anselmo: 100 percent

### **Component Descriptions**

Anselmo

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Loamy eolian deposits

*Slope:* 6 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 7.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Sandy - Veg. Zone 3  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 7 inches; fine sandy loam  
 H2—7 to 18 inches; fine sandy loam  
 H3—18 to 33 inches; fine sandy loam  
 H4—33 to 60 inches; loamy fine sand

## **AtF—Anselmo-Brunswick fine sandy loams, 11 to 30 percent slopes**

### **Map Unit Composition**

Anselmo: 60 percent  
 Brunswick: 40 percent

### **Component Descriptions**

#### **Anselmo**

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Valley side on valley

*Hillslope position:* Summit, shoulder

*Parent material:* Loamy eolian deposits

*Slope:* 17 to 30 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 7.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam  
 H2—6 to 28 inches; fine sandy loam  
 H3—28 to 37 inches; fine sandy loam  
 H4—37 to 60 inches; fine sand

#### **Brunswick**

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Valley side on valley

*Hillslope position:* Backslope

*Slope:* 11 to 30 percent

*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 1.98 in/hr)

*Available water capacity:* Low (About 3.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 4 inches; fine sandy loam  
 H2—4 to 26 inches; fine sandy loam  
 H3—26 to 33 inches; fine sandy loam  
 Cr—33 to 60 inches; weathered bedrock

### **Minor Components**

#### **Perched Wt**

## **Ba—Barney fine sandy loam, Channeled**

### **Map Unit Composition**

Barney: 100 percent

### **Component Descriptions**

#### **Barney**

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flood plain on valley

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Very poorly drained

*Slowest permeability:* Moderately rapid (About 1.98 in/hr)

*Available water capacity:* Low (About 4.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Frequent

*Depth to seasonal water saturation:* About 0 to 12 inches

*Runoff class:* Negligible

*Ecological site:* Wet Land

*Land capability (nonirrigated):* 6w

*Typical Profile:*

H1—0 to 10 inches; fine sandy loam  
 H2—10 to 20 inches; stratified sand to loam  
 H3—20 to 60 inches; coarse sand

**Bd—Bolent fine sandy loam, 0 to 2 percent slopes**

## Map Unit Composition

Bolent: 100 percent

## Component Descriptions

Bolent

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Flood plain on valley*Parent material:* Sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Low (About 5.1 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* About 18 to 36 inches*Runoff class:* Very low*Ecological site:* Subirrigated - Veg. Zone 3*Land capability (irrigated):* 4w*Land capability (nonirrigated):* 4w*Typical Profile:*

H1—0 to 5 inches; fine sandy loam

H2—5 to 60 inches; stratified sand to fine sandy loam

**Minor Components****Almeria***Slope:* 0 to 2 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**Bo—Brocksburg loam, 0 to 1 percent slopes**

## Map Unit Composition

Brocksburg: 100 percent

## Component Descriptions

Brocksburg

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Flat on tableland*Parent material:* Loamy alluvium over sandy and gravelly alluvium*Slope:* 0 to 1 percent*Drainage class:* Well drained*Slowest permeability:* Moderately slow (About 0.20 in/hr)*Available water capacity:* Moderate (About 7.6 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Silty - Veg. Zone 3*Land capability (irrigated):* 2s*Land capability (nonirrigated):* 2s*Typical Profile:*

H1—0 to 16 inches; loam

H2—16 to 36 inches; clay loam

H3—36 to 60 inches; coarse sand

**Minor Components****Perched Wt****BrD—Brunswick fine sandy loam, 3 to 9 percent slopes**

## Map Unit Composition

Brunswick: 100 percent

## Component Descriptions

Brunswick

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Hill on tableland, valley side on valley*Parent material:* Calcareous loamy residuum weathered from sandstone*Slope:* 3 to 9 percent*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)*Drainage class:* Well drained*Slowest permeability:* Moderately rapid (About 1.98 in/hr)*Available water capacity:* Low (About 3.3 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Sandy - Veg. Zone 3*Land capability (irrigated):* 4e

*Land capability (nonirrigated): 4e*

*Typical Profile:*

H1—0 to 4 inches; fine sandy loam  
H2—4 to 17 inches; fine sandy loam  
H3—17 to 36 inches; fine sand  
Cr—36 to 60 inches; weathered bedrock

**Minor Components**

**Wt At 0-1 Foot**

**DuB—Dunday loamy fine sand, 0 to 3 percent slopes**

Map Unit Composition

Dunday: 100 percent

Component Descriptions

Dunday

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hummock on tableland, interdune on sandhills

*Parent material:* Eolian sands

*Slope:* 0 to 3 percent

*Drainage class:* Somewhat excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 5.3 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated): 3e*

*Land capability (nonirrigated): 4e*

*Typical Profile:*

H1—0 to 18 inches; loamy fine sand  
H2—18 to 60 inches; fine sand

**Minor Components**

**Wt At 0-1 Foot**

**DuD—Dunday loamy fine sand, 3 to 9 percent slopes**

Map Unit Composition

Dunday: 100 percent

Component Descriptions

Dunday

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Dune on tableland, interdune on sandhills

*Parent material:* Eolian sands

*Slope:* 3 to 8 percent

*Drainage class:* Somewhat excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 5.3 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated): 4e*

*Land capability (nonirrigated): 4e*

*Typical Profile:*

H1—0 to 18 inches; loamy fine sand  
H2—18 to 60 inches; fine sand

**Minor Components**

**Wt At 0-1 Foot**

**Eo—Els loamy sand, 0 to 2 percent slopes**

Map Unit Composition

Els: 100 percent

Component Descriptions

Els

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Sandy eolian deposits over sandy alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 4.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated

*Land capability (irrigated):* 4w  
*Land capability (nonirrigated):* 4w

**Typical Profile:**

H1—0 to 6 inches; loamy sand  
 H2—6 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**Marlake**

*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained

**EpB—Els-lpage fine sands, 0 to 3 percent slopes**

**Map Unit Composition**

Els: 65 percent  
 lpage: 35 percent

**Component Descriptions**

**Els**

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Swale on interdune on sandhills  
*Parent material:* Sandy eolian deposits over sandy alluvium

*Slope:* 0 to 3 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.3 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 18 to 36 inches  
*Runoff class:* Negligible  
*Ecological site:* Subirrigated  
*Land capability (irrigated):* 4w  
*Land capability (nonirrigated):* 6e

**Typical Profile:**

H1—0 to 6 inches; fine sand  
 H2—6 to 60 inches; fine sand

**lpage**

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Hummock on interdune on sandhills  
*Parent material:* Eolian sands over sandy alluvium

*Slope:* 0 to 3 percent  
*Drainage class:* Moderately well drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.2 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 36 to 60 inches  
*Runoff class:* Negligible  
*Ecological site:* Sandy Lowland 22-25" P.z.  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

**Typical Profile:**

H1—0 to 7 inches; fine sand  
 H2—7 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**Marlake**

*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained

**Es—Elsmere loamy fine sand, 0 to 2 percent slopes**

**Map Unit Composition**

Elsmere: 100 percent

**Component Descriptions**

**Elsmere**

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Swale on interdune on sandhills  
*Parent material:* Sandy alluvium and/or eolian sands  
*Slope:* 0 to 2 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.3 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 18 to 36 inches  
*Runoff class:* Negligible  
*Ecological site:* Subirrigated  
*Land capability (irrigated):* 4w  
*Land capability (nonirrigated):* 4w

*Typical Profile:*

H1—0 to 14 inches; loamy fine sand  
H2—14 to 60 inches; fine sand

**Minor Components****Loup**

*Slope:* 0 to 2 percent  
*Drainage class:* Poorly drained  
*Ecological site:* Wet Subirrigated

**Fe—Fluvaquents, sandy**

## Map Unit Composition

Fluvaquents: 100 percent

## Component Descriptions

## Fluvaquents

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flood plain on valley

*Parent material:* Silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Very poorly drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 5.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Frequent

*Depth to seasonal water saturation:* About 0 to 0 inches

*Runoff class:* Negligible

*Land capability (nonirrigated):* 8w

*Typical Profile:*

H1—0 to 60 inches; stratified loamy fine sand

**Gn—Gannett fine sandy loam, 0 to 2 percent slopes**

## Map Unit Composition

Gannett: 100 percent

## Component Descriptions

## Gannett

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Sandy alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Low (About 5.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* About 0 to 18 inches

*Runoff class:* Negligible

*Ecological site:* Wet Subirrigated

*Land capability (nonirrigated):* 5w

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam

H2—6 to 23 inches; loam

H3—23 to 60 inches; fine sand

**IdB—Inavale loamy fine sand, 0 to 3 percent slopes**

## Map Unit Composition

Inavale: 100 percent

## Component Descriptions

## Inavale

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flood plain on valley

*Parent material:* Sandy alluvium

*Slope:* 0 to 3 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 5.0 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy Lowland - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 8 inches; loamy fine sand

H2—8 to 60 inches; stratified fine sand to loam

**Minor Components****Almeria**

*Slope:* 0 to 2 percent

*Drainage class:* Poorly drained

*Ecological site:* Wet Subirrigated - Veg. Zone 3



**Fluvaquents***Slope:* 0 to 1 percent*Drainage class:* Very poorly drained**IkB—Inavale sand, Channeled**

## Map Unit Composition

Inavale: 100 percent

## Component Descriptions

Inavale

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Flood plain on valley*Parent material:* Sandy alluvium*Slope:* 0 to 3 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 4.7 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 3*Land capability (nonirrigated):* 6w*Typical Profile:*

H1—0 to 8 inches; sand

H2—8 to 60 inches; stratified coarse sand to loam

**Minor Components****Almeria***Slope:* 0 to 2 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**Fluvaquents***Slope:* 0 to 1 percent*Drainage class:* Very poorly drained**In—Inavale-Barney complex, Channeled**

## Map Unit Composition

Inavale: 60 percent

Barney: 40 percent

## Component Descriptions

Inavale

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Flood plain on valley*Parent material:* Sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 5.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 3*Land capability (nonirrigated):* 6w*Typical Profile:*

H1—0 to 10 inches; loamy fine sand

H2—10 to 60 inches; stratified sand to loam

**Barney***MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Flood plain on valley*Parent material:* Loamy alluvium over sandy and gravelly alluvium*Slope:* 0 to 2 percent*Drainage class:* Very poorly drained*Slowest permeability:* Moderately rapid (About 1.98 in/hr)*Available water capacity:* Low (About 4.8 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Frequent*Depth to seasonal water saturation:* About 0 to 12 inches*Runoff class:* Negligible*Ecological site:* Wet Land - Veg. Zone 3*Land capability (nonirrigated):* 6w*Typical Profile:*

H1—0 to 3 inches; fine sandy loam

H3—7 to 60 inches; sand

H2—10 to 20 inches; stratified sand to loam

**Minor Components****Fluvaquents***Slope:* 0 to 1 percent*Drainage class:* Very poorly drained**IpB—Ipage fine sand, 0 to 3 percent slopes**

**Map Unit Composition**

lpage: 100 percent

**Component Descriptions**

lpage

MLRA: 65 - Nebraska Sand Hills

Landform: Stream terrace on valley, hummock  
on interdune on sandhills

Parent material: Eolian sands over sandy  
alluvium

Slope: 0 to 3 percent

Drainage class: Moderately well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 4.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 36 to  
60 inches

Runoff class: Negligible

Ecological site: Sandy Lowland 22-25" P.z.

Land capability (irrigated): 4e

Land capability (nonirrigated): 6e

**Typical Profile:**

H1—0 to 7 inches; fine sand

H2—7 to 60 inches; fine sand

**Minor Components****Tryon**

Slope: 0 to 2 percent

Drainage class: Very poorly drained

Ecological site: Wet Land

**IsB—Ipage loamy sand, 0 to 3 percent slopes****Map Unit Composition**

lpage: 100 percent

**Component Descriptions**

lpage

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Hummock on interdune on sandhills,  
stream terrace on valley

Parent material: Eolian sands over sandy  
alluvium

Slope: 0 to 3 percent

Drainage class: Moderately well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 4.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 36 to  
60 inches

Runoff class: Negligible

Ecological site: Sandy Lowland - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

**Typical Profile:**

H1—0 to 7 inches; loamy sand

H2—7 to 60 inches; sand

**Minor Components****Tryon**

Slope: 0 to 2 percent

Drainage class: Very poorly drained

Ecological site: Wet Land

**Jn—Jansen fine sandy loam, 0 to 2 percent slopes****Map Unit Composition**

Jansen: 100 percent

**Component Descriptions**

Jansen

MLRA: 66 - Dakota-Nebraska Eroded Tableland

Landform: Flat on tableland

Parent material: Loamy alluvium over sandy and  
gravelly alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About  
0.20 in/hr)

Available water capacity: Low (About 5.8 inches)

Shrink-swell potential: Moderate (About 4.5  
LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6  
feet

Runoff class: Low

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

**Typical Profile:**

H1—0 to 13 inches; fine sandy loam

H2—13 to 30 inches; clay loam

H3—30 to 60 inches; sand

**Minor Components**  
**Perched Wt**

**JnC—Jansen fine sandy loam, 2 to 6 percent slopes**

Map Unit Composition

Jansen: 100 percent

Component Descriptions

Jansen

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Valley side on valley, hill on tableland

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 2 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Low (About 5.8 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 13 inches; fine sandy loam

H2—13 to 30 inches; clay loam

H3—30 to 60 inches; coarse sand

**Minor Components**  
**Perched Wt**

**Jo—Jansen loam, 0 to 2 percent slopes**

Map Unit Composition

Jansen: 100 percent

Component Descriptions

Jansen

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flat on tableland

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 2s

*Land capability (nonirrigated):* 2s

*Typical Profile:*

H1—0 to 13 inches; loam

H2—13 to 30 inches; loam

H3—30 to 60 inches; sand

**Minor Components**  
**Perched Wt**

**JoC—Jansen loam, 2 to 6 percent slopes**

Map Unit Composition

Jansen: 100 percent

Component Descriptions

Jansen

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 2 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 3  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 13 inches; loam  
 H2—13 to 30 inches; clay loam  
 H3—30 to 60 inches; gravelly coarse sand

**Minor Components**  
**Perched Wt**

**Jr—Jansen-Meadin complex, 0 to 2 percent slopes**

Map Unit Composition

Jansen: 50 percent  
 Meadin: 50 percent

Component Descriptions

Jansen

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Knoll on tableland  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* Low (About 4.8 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Silty - Veg. Zone 3  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 7 inches; fine sandy loam  
 H2—7 to 22 inches; clay loam  
 H3—22 to 60 inches; sand

Meadin

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Flat on tableland  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Excessively drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Low (About 3.3 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Shallow To Gravel - Veg. Zone 3  
*Land capability (irrigated):* 4s  
*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 5 inches; sandy loam  
 H2—5 to 10 inches; loamy sand  
 H3—10 to 60 inches; gravelly coarse sand

**Minor Components**  
**Perched Wt**

**JtB—Jansen-Sandose complex, 0 to 3 percent slopes**

Map Unit Composition

Jansen: 50 percent  
 Sandose: 50 percent

Component Descriptions

Jansen

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Flat on tableland  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* Moderate (About 6.1 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Silty - Veg. Zone 3  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 11 inches; fine sandy loam  
 H2—11 to 32 inches; clay loam

H3—32 to 60 inches; coarse sand

**Sandose**

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Knoll on tableland

*Parent material:* Eolian sands over loamy sediments

*Slope:* 0 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 7.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 10 inches; loamy fine sand

H2—10 to 38 inches; loamy fine sand

H3—38 to 60 inches; loam

**Jw—Johnstown fine sandy loam, 0 to 1 percent slopes**

**Map Unit Composition**

Johnstown: 100 percent

**Component Descriptions**

Johnstown

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flat on tableland

*Parent material:* Loess over gravelly sand

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 9.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 21 inches; fine sandy loam

H2—21 to 44 inches; clay loam

H3—44 to 50 inches; silty clay loam

H4—50 to 60 inches; gravelly coarse sand

**Minor Components**

**Perched Wt**

**JwB—Johnstown fine sandy loam, 1 to 3 percent slopes**

**Map Unit Composition**

Johnstown: 100 percent

**Component Descriptions**

Johnstown

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland

*Parent material:* Loess over gravelly sand

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 9.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 21 inches; fine sandy loam

H2—21 to 44 inches; clay loam

H3—44 to 50 inches; silty clay loam

H4—50 to 60 inches; sand

**Minor Components**

**Perched Wt**

**Jy—Johnstown loam, 0 to 1 percent slopes**

**Map Unit Composition**

Johnstown: 100 percent

**Component Descriptions**

Johnstown

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flat on tableland

*Parent material:* Loess over gravelly sand

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

**Typical Profile:**

H1—0 to 21 inches; loam

H2—21 to 44 inches; clay loam

H3—44 to 50 inches; silty clay loam

H4—50 to 60 inches; coarse sand

**Minor Components**

**Perched Wt**

**JyB—Johnstown loam, 1 to 3 percent slopes****Map Unit Composition**

Johnstown: 100 percent

**Component Descriptions**

Johnstown

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland

*Parent material:* Loess over gravelly sand

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

**Typical Profile:**

H1—0 to 21 inches; loam

H2—21 to 44 inches; clay loam

H3—44 to 50 inches; silty clay loam

H4—50 to 60 inches; sand

**Minor Components**

**Perched Wt**

**JyC—Johnstown loam, 3 to 6 percent slopes****Map Unit Composition**

Johnstown: 100 percent

**Component Descriptions**

Johnstown

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Loess over gravelly sand

*Slope:* 3 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Silty - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

**Typical Profile:**

H1—0 to 21 inches; loam

H2—21 to 44 inches; clay loam

H3—44 to 50 inches; silty clay loam

H4—50 to 60 inches; loamy sand

**Minor Components**  
**Perched Wt**

**LcG—Labu-Sansarc silty clays, 11 to 40 percent slopes**

Map Unit Composition

Labu: 55 percent  
Sansarc: 45 percent

Component Descriptions

Labu

MLRA: 63B - Southern Rolling Pierre Shale  
Plains

*Landform:* Valley side on valley

*Hillslope position:* Backslope

*Parent material:* Residuum weathered from  
clayey shale

*Slope:* 11 to 30 percent

*Depth to restrictive feature:* 20 to 40 inches to  
bedrock (paralithic)

*Drainage class:* Well drained

*Slowest permeability:* Very slow (About 0.00  
in/hr)

*Available water capacity:* Very low (About 2.5  
inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6  
feet

*Runoff class:* Very high

*Ecological site:* Clayey - Veg. Zone 3

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 5 inches; silty clay

H2—5 to 23 inches; silty clay

Cr—23 to 60 inches; weathered bedrock

Sansarc

MLRA: 63B - Southern Rolling Pierre Shale  
Plains

*Landform:* Valley side on valley

*Hillslope position:* Summit, shoulder

*Parent material:* Clayey residuum weathered  
from shale

*Slope:* 11 to 40 percent

*Depth to restrictive feature:* 4 to 20 inches to  
bedrock (paralithic)

*Drainage class:* Well drained

*Slowest permeability:* Very slow (About 0.00  
in/hr)

*Available water capacity:* Very low (About 1.1  
inches)

*Shrink-swell potential:* Very high (About 17.0  
LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6  
feet

*Runoff class:* Very high

*Ecological site:* Shallow Clay - Veg. Zone 3

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 3 inches; silty clay

H2—3 to 12 inches; clay

Cr—12 to 60 inches; weathered bedrock

**LfB—Libory loamy fine sand, 0 to 3 percent slopes**

Map Unit Composition

Libory: 100 percent

Component Descriptions

Libory

MLRA: 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Eolian sands over loess

*Slope:* 0 to 3 percent

*Drainage class:* Moderately well drained

*Slowest permeability:* Moderately slow (About  
0.20 in/hr)

*Available water capacity:* High (About 9.2  
inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* About 18 to  
36 inches

*Runoff class:* Very low

*Ecological site:* Sandy Lowland 22-25" P.z.

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 14 inches; loamy fine sand

H2—14 to 25 inches; loamy fine sand

H3—25 to 60 inches; loam

**Minor Components**

**Wt At 0-1 Foot**

**Lo—Loup fine sandy loam, 0 to 2 percent slopes**

## Map Unit Composition

Loup: 100 percent

## Component Descriptions

## Loup

*MLRA:* 65 - Nebraska Sand Hills*Landform:* Swale on interdune on sandhills*Parent material:* Sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Poorly drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Low (About 5.1 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 0 to 18 inches*Runoff class:* Very low*Ecological site:* Wet Subirrigated*Land capability (nonirrigated):* 5w*Typical Profile:*

H1—0 to 10 inches; fine sandy loam

H2—10 to 60 inches; fine sand

**Lp—Loup fine sandy loam, Wet, 0 to 2 percent slopes**

## Map Unit Composition

Loup: 100 percent

## Component Descriptions

## Loup

*MLRA:* 65 - Nebraska Sand Hills*Landform:* Swale on interdune on sandhills*Parent material:* Sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Very poorly drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Low (About 5.1 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 0 to 0 inches*Runoff class:* Negligible*Ecological site:* Wet Land*Land capability (nonirrigated):* 5w*Typical Profile:*

H1—0 to 10 inches; fine sandy loam

H2—10 to 60 inches; fine sand

**LtB—Loup-Els complex, 0 to 3 percent slopes**

## Map Unit Composition

Loup: 50 percent

Els: 50 percent

## Component Descriptions

## Loup

*MLRA:* 65 - Nebraska Sand Hills*Landform:* Swale on interdune on sandhills*Parent material:* Sandy alluvium*Slope:* 0 to 2 percent*Drainage class:* Poorly drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Low (About 5.2 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 0 to 18 inches*Runoff class:* Very low*Ecological site:* Wet Subirrigated*Land capability (nonirrigated):* 5w*Typical Profile:*

H1—0 to 11 inches; fine sandy loam

H2—11 to 60 inches; sand

## Els

*MLRA:* 65 - Nebraska Sand Hills*Landform:* Hummock on interdune on sandhills*Parent material:* Sandy eolian deposits over sandy alluvium*Slope:* 0 to 3 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 4.4 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 18 to 36 inches*Runoff class:* Negligible*Ecological site:* Subirrigated*Land capability (irrigated):* 4w*Land capability (nonirrigated):* 4w



*Typical Profile:*

H1—0 to 6 inches; loamy fine sand  
H2—6 to 60 inches; fine sand

**Minor Components****Marlake**

*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained

**Loup**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**Ma—Marlake fine sandy loam, 0 to 1 percent slopes**

## Map Unit Composition

Marlake: 100 percent

## Component Descriptions

## Marlake

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Depression on interdune on sandhills  
*Parent material:* Sandy eolian deposits over alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained  
*Slowest permeability:* Moderately rapid (About 1.98 in/hr)  
*Available water capacity:* Low (About 5.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 0 to 0 inches  
*Runoff class:* Negligible  
*Land capability (nonirrigated):* 8w

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam  
H2—6 to 60 inches; loamy fine sand

**McG—Mckelvie-Tassel-Ronson complex, 15 to 70 percent slopes**

## Map Unit Composition

Mckelvie: 50 percent

Longpine: 30 percent

Minor components: 20 percent

## Component Descriptions

## Mckelvie

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Valley side on valley  
*Hillslope position:* Backslope  
*Parent material:* Sandy residuum weathered from sandstone  
*Slope:* 15 to 70 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.9 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Sands - Veg. Zone 3  
*Land capability (nonirrigated):* 7e

*Typical Profile:*

H1—0 to 6 inches; loamy fine sand  
H2—6 to 60 inches; fine sand

## Longpine

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Valley side on valley  
*Hillslope position:* Shoulder  
*Parent material:* Calcareous loamy residuum weathered from sandstone  
*Slope:* 15 to 70 percent  
*Depth to restrictive feature:* 10 to 20 inches to bedrock (paralithic)  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Very low (About 1.4 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very high  
*Ecological site:* Shallow Limy - Veg. Zone 3  
*Land capability (nonirrigated):* 7s

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam  
H2—6 to 10 inches; gravelly fine sandy loam  
Cr—10 to 60 inches; unweathered bedrock

**Minor Components****Ronson**

*Composition:* About 20 percent  
*Landform:* valley side on valley  
*Slope:* 15 to 30 percent

*Depth to restrictive feature:* 20 to 40 inches  
to bedrock (paralithic)  
*Drainage class:* Well drained  
*Ecological site:* Savannah - Veg. Zone 3

**Map Unit Composition**

Meadin: 100 percent

**Fluvaquents**

*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained

**Barney**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**MeB—Meadin sandy loam, 0 to 3 percent slopes****Map Unit Composition**

Meadin: 100 percent

**Component Descriptions**

Meadin

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Flat on tableland, knoll on tableland  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 3 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Moderately rapid (About 1.98 in/hr)  
*Available water capacity:* Low (About 3.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low  
*Ecological site:* Shallow To Gravel - Veg. Zone 3  
*Land capability (irrigated):* 4s  
*Land capability (nonirrigated):* 6s

**Typical Profile:**

H1—0 to 7 inches; sandy loam  
H2—7 to 12 inches; very gravelly loamy sand, loamy sand  
H3—12 to 60 inches; gravelly coarse sand

**MeF—Meadin sandy loam, 3 to 30 percent slopes****Component Descriptions**

Meadin

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Hill on tableland, valley side on valley  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 3 to 30 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Moderately rapid (About 1.98 in/hr)

*Available water capacity:* Low (About 3.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Shallow To Gravel - Veg. Zone 3  
*Land capability (nonirrigated):* 6s

**Typical Profile:**

H1—0 to 7 inches; sandy loam  
H2—7 to 12 inches; sandy loam  
H3—12 to 60 inches; gravelly coarse sand

**Minor Components**

**Wt At 0-1 Foot**

**Oe—O'Neill fine sandy loam, 0 to 2 percent slopes****Map Unit Composition**

O'Neill: 100 percent

**Component Descriptions**

O'Neill

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Flat on tableland

*Parent material:* Coarse-loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 1.98 in/hr)

*Available water capacity:* Low (About 4.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 8 inches; fine sandy loam

H2—8 to 26 inches; fine sandy loam

H3—26 to 60 inches; gravelly coarse sand

**Minor Components**

**Perched Wt**

**On—O'Neill loam, 0 to 2 percent slopes**

Map Unit Composition

O'Neill: 100 percent

Component Descriptions

O'Neill

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Flat on tableland

*Parent material:* Coarse-loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Low (About 4.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 2s

*Land capability (nonirrigated):* 2c

*Typical Profile:*

H1—0 to 8 inches; loam

H2—8 to 26 inches; fine sandy loam

H3—26 to 60 inches; coarse sand

**OsC—O'Neill-Meadin sandy loams, 2 to 6 percent slopes**

Map Unit Composition

O'Neill: 60 percent

Meadin: 40 percent

Component Descriptions

O'Neill

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Coarse-loamy alluvium over sandy and gravelly alluvium

*Slope:* 2 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 1.98 in/hr)

*Available water capacity:* Low (About 4.8 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 8 inches; sandy loam

H2—8 to 30 inches; fine sandy loam

H3—30 to 60 inches; sand

Meadin

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 2 to 6 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Moderately rapid (About 1.98 in/hr)

*Available water capacity:* Low (About 3.5 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Shallow To Gravel - Veg. Zone 3

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 8 inches; sandy loam

H2—8 to 12 inches; loamy sand

H3—12 to 60 inches; gravelly coarse sand

**Minor Components**

**Perched Wt**

## OsD—O'Neill-Meadin sandy loams, 6 to 11 percent slopes

### Map Unit Composition

O'Neill: 55 percent  
Meadin: 45 percent

### Component Descriptions

#### O'Neill

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Coarse-loamy alluvium over  
sandy and gravelly alluvium

*Slope:* 6 to 11 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About  
1.98 in/hr)

*Available water capacity:* Low (About 4.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6  
feet

*Runoff class:* Low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

#### Typical Profile:

- H1—0 to 6 inches; sandy loam
- H2—6 to 24 inches; fine sandy loam
- H3—24 to 60 inches; gravelly coarse sand

#### Meadin

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Loamy alluvium over sandy and  
gravelly alluvium

*Slope:* 6 to 11 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Moderately rapid (About  
1.98 in/hr)

*Available water capacity:* Low (About 3.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6  
feet

*Runoff class:* Low

*Ecological site:* Shallow To Gravel - Veg. Zone 3

*Land capability (nonirrigated):* 6s

#### Typical Profile:

- H1—0 to 9 inches; sandy loam
- H2—9 to 12 inches; loamy sand

H3—12 to 60 inches; gravelly coarse sand

### Minor Components Wt At 0-1 Foot

## Pg—Pits, sand And Gravel

### Map Unit Composition

Pits: 100 percent

### Component Descriptions

#### Pits

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Tableland

*Slope:* 0 to 30 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.5 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6  
feet

*Runoff class:* Very low

*Land capability (nonirrigated):* 8s

### Minor Components

#### Marlake

*Slope:* 0 to 1 percent

*Drainage class:* Very poorly drained

## PtB—Pivot loamy sand, 0 to 3 percent slopes

### Map Unit Composition

Pivot: 100 percent

### Component Descriptions

#### Pivot

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hummock on tableland, interdune on  
sandhills

*Parent material:* Sandy eolian deposits over  
gravelly alluvium

*Slope:* 0 to 3 percent

*Drainage class:* Somewhat excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Sandy - Veg. Zone 3  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 16 inches; loamy sand  
 H2—16 to 27 inches; loamy sand  
 H3—27 to 60 inches; coarse sand

**Minor Components**

**Wt At 0-1 Foot**

**RtB—Ronson-Tassel fine sandy loams, 0 to 3 percent slopes**

Map Unit Composition

Ronson: 55 percent  
 Longpine: 45 percent

Component Descriptions

Ronson  
*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Swale on tableland  
*Parent material:* Residuum weathered from calcareous sandstone  
*Slope:* 0 to 3 percent  
*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Low (About 3.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sandy - Veg. Zone 3  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 13 inches; fine sandy loam  
 H2—13 to 27 inches; sandy loam  
 Cr—27 to 60 inches; weathered bedrock

Longpine  
*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Knoll on tableland  
*Parent material:* Calcareous loamy residuum weathered from sandstone  
*Slope:* 0 to 3 percent  
*Depth to restrictive feature:* 10 to 20 inches to bedrock (paralithic)  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Very low (About 1.9 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Shallow Limy - Veg. Zone 3  
*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam  
 H2—6 to 17 inches; gravelly fine sandy loam  
 Cr—17 to 60 inches; unweathered bedrock

**Minor Components**

**Perched Wt**

**RtC—Ronson-Tassel fine sandy loams, 3 to 6 percent slopes**

Map Unit Composition

Ronson: 55 percent  
 Longpine: 45 percent

Component Descriptions

Ronson  
*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Hill on tableland  
*Hillslope position:* Backslope  
*Parent material:* Residuum weathered from calcareous sandstone  
*Slope:* 3 to 6 percent  
*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Low (About 3.9 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 13 inches; fine sandy loam

H2—13 to 37 inches; sandy loam

Cr—37 to 60 inches; weathered bedrock

Longpine

MLRA: 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland

*Hillslope position:* Summit, shoulder

*Parent material:* Calcareous loamy residuum weathered from sandstone

*Slope:* 3 to 6 percent

*Depth to restrictive feature:* 10 to 20 inches to bedrock (paralithic)

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Very low (About 1.3 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Shallow Limy - Veg. Zone 3

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam

H2—6 to 9 inches; gravelly fine sandy loam

Cr—9 to 60 inches; unweathered bedrock

**Minor Components**

**Perched Wt**

**RtD—Ronson-Tassel fine sandy loams, 6 to 11 percent slopes**

Map Unit Composition

Ronson: 55 percent

Longpine: 45 percent

Component Descriptions

Ronson

MLRA: 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland

*Hillslope position:* Backslope

*Parent material:* Residuum weathered from calcareous sandstone

*Slope:* 6 to 11 percent

*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Low (About 3.1 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 12 inches; fine sandy loam

H2—12 to 24 inches; fine sandy loam

Cr—24 to 60 inches; weathered bedrock

Longpine

MLRA: 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on tableland

*Hillslope position:* Summit, shoulder

*Parent material:* Calcareous loamy residuum weathered from sandstone

*Slope:* 6 to 11 percent

*Depth to restrictive feature:* 10 to 20 inches to bedrock (paralithic)

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Very low (About 1.5 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Shallow Limy - Veg. Zone 3

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam

H2—6 to 11 inches; gravelly fine sandy loam

Hr—11 to 60 inches; unweathered bedrock

**Minor Components**

**Perched Wt**

**ScB—Sandose loamy fine sand, 0 to 3 percent slopes**

## Map Unit Composition

Sandose: 100 percent

## Component Descriptions

Sandose

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Interdune on sandhills, swale on tableland

*Parent material:* Eolian sands over loamy sediments

*Slope:* 0 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 8.1 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy - Veg. Zone 3

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 15 inches; loamy fine sand

H2—15 to 26 inches; loamy fine sand

H3—26 to 46 inches; loam

H4—46 to 60 inches; very fine sandy loam

**Minor Components****Perched Wt****SkB—Simeon loamy sand, 0 to 3 percent slopes**

## Map Unit Composition

Simeon: 100 percent

## Component Descriptions

Simeon

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Interdune on sandhills, hummock on tableland

*Parent material:* Sandy and gravelly alluvium

*Slope:* 0 to 3 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 4.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Shallow To Gravel 17-25" P.z.

*Land capability (irrigated):* 4s

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 5 inches; loamy sand

H2—5 to 60 inches; coarse sand

**Minor Components****Wt At 0-1 Foot****SkD—Simeon loamy sand, 3 to 9 percent slopes**

## Map Unit Composition

Simeon: 100 percent

## Component Descriptions

Simeon

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Hill on tableland, valley side on valley

*Parent material:* Sandy and gravelly alluvium

*Slope:* 3 to 9 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 4.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Shallow To Gravel 17-25" P.z.

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 5 inches; loamy sand

H2—5 to 60 inches; coarse sand

**Minor Components****Wt At 0-1 Foot**

## **SvD—Simeon-Valentine fine sands, 0 to 9 percent slopes**

### Map Unit Composition

Simeon: 60 percent  
Valentine: 40 percent

### Component Descriptions

#### Simeon

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on tableland, interdune on sandhills

*Parent material:* Sandy and gravelly alluvium

*Slope:* 0 to 3 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 4.8 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Shallow To Gravel 17-25" P.z.

*Land capability (irrigated):* 4s

*Land capability (nonirrigated):* 6s

#### Typical Profile:

H1—0 to 5 inches; fine sand

H2—5 to 60 inches; coarse sand

#### Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Interdune on sandhills, dune on tableland

*Parent material:* Eolian sands

*Slope:* 3 to 9 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands 22-25" P.z.

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

#### Typical Profile:

H1—0 to 6 inches; fine sand

H2—6 to 60 inches; fine sand

## **Minor Components**

### **Wt At 0-1 Foot**

## **Tn—Tryon loamy fine sand, 0 to 2 percent slopes**

### Map Unit Composition

Tryon: 100 percent

### Component Descriptions

#### Tryon

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Sandy eolian deposits over sandy alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.8 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* About 0 to 18 inches

*Runoff class:* Negligible

*Ecological site:* Wet Subirrigated

*Land capability (nonirrigated):* 5w

#### Typical Profile:

H1—0 to 5 inches; loamy fine sand

H2—5 to 60 inches; fine sand

## **To—Tryon loamy fine sand, Wet, 0 to 2 percent slopes**

### Map Unit Composition

Tryon: 100 percent

### Component Descriptions

#### Tryon

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Sandy eolian deposits over sandy alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Very poorly drained

*Slowest permeability:* Rapid (About 5.95 in/hr)



*Available water capacity:* Low (About 3.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 0 to 0 inches  
*Runoff class:* Negligible  
*Ecological site:* Wet Land  
*Land capability (nonirrigated):* 5w

*Typical Profile:*  
 H1—0 to 6 inches; loamy fine sand  
 H2—6 to 60 inches; fine sand

### **TpB—Tryon-Els complex, 0 to 3 percent slopes**

#### Map Unit Composition

Tryon: 55 percent  
 Els: 45 percent

#### Component Descriptions

##### Tryon

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Swale on interdune on sandhills  
*Parent material:* Sandy eolian deposits over sandy alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 3.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 0 to 18 inches  
*Runoff class:* Negligible  
*Ecological site:* Wet Subirrigated  
*Land capability (nonirrigated):* 5w

*Typical Profile:*  
 H1—0 to 5 inches; loamy fine sand  
 H2—5 to 60 inches; fine sand

##### Els

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Hummock on interdune on sandhills  
*Parent material:* Sandy eolian deposits over sandy alluvium  
*Slope:* 0 to 3 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.3 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 18 to 36 inches  
*Runoff class:* Negligible  
*Ecological site:* Subirrigated  
*Land capability (irrigated):* 4w  
*Land capability (nonirrigated):* 6e

*Typical Profile:*  
 H1—0 to 6 inches; fine sand  
 H2—6 to 60 inches; fine sand

#### Minor Components

##### Marlake

*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained

### **VaB—Valentine fine sand, 0 to 3 percent slopes**

#### Map Unit Composition

Valentine: 100 percent

#### Component Descriptions

##### Valentine

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Hummock on interdune on sandhills  
*Parent material:* Eolian sands  
*Slope:* 0 to 3 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 3.6 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Sandy 22-25" P.z.  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

*Typical Profile:*  
 H1—0 to 4 inches; fine sand  
 H2—4 to 60 inches; fine sand

#### Minor Components

##### Tryon

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**VaD—Valentine fine sand, 3 to 9 percent slopes**

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Dune on interdune on sandhills

*Parent material:* Eolian sands

*Slope:* 3 to 9 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands 22-25" P.z.

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 4 inches; fine sand

H2—4 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent

*Drainage class:* Very poorly drained

*Ecological site:* Wet Land

**Marlake**

*Slope:* 0 to 1 percent

*Drainage class:* Very poorly drained

**VaE—Valentine fine sand, Rolling**

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Dune on sandhills

*Parent material:* Eolian sands

*Slope:* 9 to 24 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands 22-25" P.z.

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 4 inches; fine sand

H2—4 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent

*Drainage class:* Very poorly drained

*Ecological site:* Wet Land

**VaF—Valentine fine sand, Rolling And Hilly**

Map Unit Composition

Valentine: 70 percent

Valentine: 30 percent

Component Descriptions

Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Dune on sandhills

*Parent material:* Eolian sands

*Slope:* 9 to 24 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands 22-25" P.z.

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 4 inches; fine sand

H2—4 to 60 inches; fine sand

Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Dune on sandhills  
*Parent material:* Eolian sands  
*Slope:* 24 to 60 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 3.6 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Choppy Sands 22-25" P.z.  
*Land capability (nonirrigated):* 7e

*Typical Profile:*

H1—0 to 4 inches; fine sand  
 H2—4 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**VbB—Valentine loamy fine sand, 0 to 3 percent slopes**

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Interdune on sandhills, hummock on tableland

*Parent material:* Eolian sands  
*Slope:* 0 to 3 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.1 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Sandy - Veg. Zone 3  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 7 inches; loamy fine sand  
 H2—7 to 13 inches; loamy fine sand

H3—13 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**VbD—Valentine loamy fine sand, 3 to 9 percent slopes**

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Interdune on sandhills, dune on tableland

*Parent material:* Eolian sands  
*Slope:* 3 to 9 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.1 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands 22-25" P.z.  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 7 inches; loamy fine sand  
 H2—7 to 13 inches; loamy fine sand  
 H3—13 to 60 inches; fine sand

**Minor Components**

**Tryon**

*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Ecological site:* Wet Land

**VfD—Valentine-Els fine sands, 0 to 9 percent slopes**

Map Unit Composition

Valentine: 55 percent  
Els: 45 percent

#### Component Descriptions

##### Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Dune on interdune on sandhills

*Parent material:* Eolian sands

*Slope:* 3 to 9 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands 22-25" P.z.

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

##### Typical Profile:

H1—0 to 4 inches; fine sand

H2—4 to 60 inches; fine sand

##### Els

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Sandy eolian deposits over sandy alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 4.3 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Subirrigated

*Land capability (irrigated):* 4w

*Land capability (nonirrigated):* 6e

##### Typical Profile:

H1—0 to 6 inches; fine sand

H2—6 to 60 inches; fine sand

#### Minor Components

##### Tryon

*Slope:* 0 to 2 percent

*Drainage class:* Very poorly drained

*Ecological site:* Wet Land

##### Marlake

*Slope:* 0 to 1 percent

*Drainage class:* Very poorly drained

## VhD—Valentine-Libory complex, 0 to 9 percent slopes

#### Map Unit Composition

Valentine: 60 percent

Libory: 40 percent

#### Component Descriptions

##### Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Dune on interdune on sandhills

*Parent material:* Eolian sands

*Slope:* 3 to 9 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sands 22-25" P.z.

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

##### Typical Profile:

H1—0 to 6 inches; fine sand

H2—6 to 60 inches; fine sand

##### Libory

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Swale on interdune on sandhills

*Parent material:* Eolian sands over loess

*Slope:* 0 to 3 percent

*Drainage class:* Moderately well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 9.3 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Sandy Lowland 22-25" P.z.

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

##### Typical Profile:

H1—0 to 10 inches; loamy fine sand

H2—10 to 24 inches; loamy fine sand

H3—24 to 60 inches; silty clay loam

#### Minor Components

**Wt At 0-1 Foot**

H3—28 to 60 inches; coarse sand

**VpD—Valentine-Pivot complex, 0 to 9 percent slopes**

## Map Unit Composition

Valentine: 60 percent  
Pivot: 40 percent

## Component Descriptions

## Valentine

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Dune on interdune on sandhills*Parent material:* Eolian sands*Slope:* 3 to 9 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.6 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Sands - Veg. Zone 3*Land capability (irrigated):* 4e*Land capability (nonirrigated):* 6e*Typical Profile:*

H1—0 to 5 inches; fine sand

H2—5 to 60 inches; fine sand

## Pivot

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland*Landform:* Swale on interdune on sandhills*Parent material:* Sandy eolian deposits over gravelly alluvium*Slope:* 0 to 3 percent*Drainage class:* Somewhat excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.8 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy - Veg. Zone 3*Land capability (irrigated):* 3e*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 10 inches; loamy fine sand

H2—10 to 28 inches; loamy fine sand

**Minor Components****Wt At 0-1 Foot****VrD—Valentine-Sandose loamy fine sands, 0 to 9 percent slopes**

## Map Unit Composition

Valentine: 65 percent  
Sandose: 35 percent

## Component Descriptions

## Valentine

*MLRA:* 65 - Nebraska Sand Hills*Landform:* Dune on interdune on sandhills*Parent material:* Eolian sands*Slope:* 3 to 9 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 4.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Sands 22-25" P.z.*Land capability (irrigated):* 4e*Land capability (nonirrigated):* 6e*Typical Profile:*

H1—0 to 7 inches; loamy fine sand

H2—7 to 10 inches; loamy fine sand

H3—10 to 60 inches; fine sand

## Sandose

*MLRA:* 65 - Nebraska Sand Hills*Landform:* Swale on interdune on sandhills*Parent material:* Eolian sands over loamy sediments*Slope:* 0 to 3 percent*Drainage class:* Well drained*Slowest permeability:* Moderately slow (About 0.20 in/hr)*Available water capacity:* Moderate (About 8.4 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible

*Ecological site:* Sandy 22-25" P.z.  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 10 inches; loamy fine sand  
 H2—10 to 20 inches; loamy fine sand  
 H3—20 to 40 inches; sandy clay loam  
 H4—40 to 60 inches; very fine sandy loam

**Minor Components**  
**Perched Wt**

**VsG2—Valentine-Simeon complex,  
 9 to 40 percent slopes, Eroded**

Map Unit Composition

Valentine: 65 percent  
 Simeon: 35 percent

Component Descriptions

Valentine

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Valley side on valley

*Hillslope position:* Backslope

*Parent material:* Eolian sands

*Slope:* 9 to 40 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Sands 22-25" P.z.

*Land capability (nonirrigated):* 7e

*Typical Profile:*

H1—0 to 6 inches; fine sand  
 H2—6 to 60 inches; fine sand

Simeon

*MLRA:* 65 - Nebraska Sand Hills

*Landform:* Valley side on valley

*Hillslope position:* Summit, shoulder

*Parent material:* Sandy and gravelly alluvium

*Slope:* 9 to 30 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 4.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Shallow To Gravel 17-25" P.z.

*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 7 inches; loamy sand  
 H2—7 to 60 inches; sand

**Minor Components**  
**Marlake**

*Slope:* 0 to 1 percent

*Drainage class:* Very poorly drained

**VtE—Valentine-Tassel complex, 3  
 to 17 percent slopes**

Map Unit Composition

Valentine: 60 percent  
 Longpine: 40 percent

Component Descriptions

Valentine

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Dune on sandhills

*Parent material:* Eolian sands

*Slope:* 3 to 17 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 5.95 in/hr)

*Available water capacity:* Low (About 3.7 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands - Veg. Zone 3

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 7 inches; fine sand  
 H2—7 to 60 inches; fine sand

Longpine

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland

*Landform:* Hill on sandhills

*Hillslope position:* Backslope

*Parent material:* Calcareous loamy residuum weathered from sandstone

*Slope:* 3 to 17 percent

*Depth to restrictive feature:* 10 to 20 inches to bedrock (paralithic)  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Very low (About 1.7 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very high  
*Ecological site:* Shallow Limy - Veg. Zone 3  
*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 6 inches; fine sandy loam  
 H2—6 to 13 inches; gravelly fine sandy loam  
 Cr—13 to 60 inches; unweathered bedrock

## **VwE—Valentine-Tryon complex, 0 to 17 percent slopes**

### Map Unit Composition

Valentine: 65 percent  
 Tryon: 35 percent

### Component Descriptions

#### Valentine

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Dune on interdune on sandhills  
*Parent material:* Eolian sands  
*Slope:* 3 to 17 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 3.6 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands 22-25" P.z.  
*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 4 inches; fine sand  
 H2—4 to 60 inches; fine sand

#### Tryon

*MLRA:* 65 - Nebraska Sand Hills  
*Landform:* Swale on interdune on sandhills

*Parent material:* Sandy eolian deposits over sandy alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Very poorly drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 3.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* About 0 to 0 inches  
*Runoff class:* Negligible  
*Ecological site:* Wet Land  
*Land capability (nonirrigated):* 5w

*Typical Profile:*

H1—0 to 6 inches; loamy fine sand  
 H2—6 to 60 inches; fine sand

### Minor Components

#### Marlake

*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained

## **VxB—Vetal loam, 1 to 3 percent slopes**

### Map Unit Composition

Vetal: 100 percent

### Component Descriptions

#### Vetal

*MLRA:* 66 - Dakota-Nebraska Eroded Tableland  
*Landform:* Swale on tableland, valley side on valley  
*Hillslope position:* Footslope  
*Parent material:* Loamy alluvium over eolian deposits  
*Slope:* 1 to 3 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Moderate (About 8.9 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sandy - Veg. Zone 3  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 16 inches; loam  
H2—16 to 31 inches; loam  
H3—31 to 56 inches; loam  
H4—56 to 60 inches; loamy fine sand

**Minor Components****Wt At 0-1 Foot***Depth to seasonal water saturation:* More than 6 feet**zwb—Water < 40 Acres**

Map Unit Composition

Water: 100 percent

**zwa—Water > 40 Acres**

Map Unit Composition

Water: 100 percent

Component Descriptions

Water

*MLRA:* 65 - Nebraska Sand Hills

Component Descriptions

Water

*MLRA:* 65 - Nebraska Sand Hills*Depth to seasonal water saturation:* More than 6 feet



Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive land-forming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes. In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

(Class 1) soils have slight limitations that restrict their use.

(Class 2) soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

(Class 3) soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

(Class 4) soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

(Class 5) soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 6) soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 7) soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

(Class 8) soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, e, w, s, or c, to the class numeral, for example, 2e. The letter e shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; w shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); s shows that the soil is limited mainly because it is shallow, droughty, or stony; and c, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by w, s, or c because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2e-4 and 3e-6. These units are not given in all soil surveys.

The capability classification of map units in this survey area is given in the section "Detailed Soil Map Units" and in the Land Capability and Component Yields table.

#### Crop Yield Estimates

The average yields per acre that can be expected of the principal crops under a high level of management are shown in "Land Capability and Component Yields" table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors. The land capability classification of map units in the survey area also is shown in the table.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, animal waste manure, and green manure crops; and harvesting that ensures the smallest possible loss.

For yields of irrigated crops, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in this table, are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service (NRCS) or the Cooperative Extension Service (CES) can provide information about the management and productivity of the soils for those crops.

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued  
Brown County, Nebraska

PAGE 2 of 5

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Alfalfa hay		Corn		Winter wheat	
	N	I	N	I	N	I	N	I
			Tons		Bu		Bu	
Aa: ALMERIA-----	5w	---	---	---	---	---	---	---
Ae: ALMERIA-----	5w	---	---	---	---	---	---	---
Af: ALMERIA-----	6w	---	---	---	---	---	---	---
HISTOSOLS-----	8w	---	---	---	---	---	---	---
An: ANSELMO-----	2e	2e	2.40	5.00	44.00	130.00	27.00	---
AnC: ANSELMO-----	3e	3e	2.30	4.50	41.00	125.00	25.00	---
AnD: ANSELMO-----	4e	4e	1.90	3.50	37.00	110.00	21.00	---
AtF: ANSELMO-----	6e	---	---	---	---	---	---	---
BRUNSWICK-----	6e	---	---	---	---	---	---	---
Ba: BARNEY-----	6w	---	---	---	---	---	---	---
Bd: BOLENT-----	4w	4w	2.30	4.10	24.00	110.00	20.00	---
Bo: BROCKSBURG-----	2s	2s	2.00	5.50	43.00	150.00	28.00	---
BrD: BRUNSWICK-----	4e	4e	1.90	3.50	36.00	105.00	21.00	---
DuB: DUNDAY-----	4e	3e	1.50	4.00	34.00	110.00	20.00	---
DuD: DUNDAY-----	4e	4e	1.10	3.30	31.00	90.00	16.00	---
EO: ELS-----	4w	4w	1.50	4.00	34.00	110.00	20.00	---
EpB: ELS-----	6e	4w	---	3.40	---	100.00	---	---
IPAGE-----	6e	4e	---	3.40	---	100.00	---	---
Es: ELSMERE-----	4w	4w	1.60	4.00	---	115.00	18.00	---
Fe: FLUVAQUENTS-----	8w	---	---	---	---	---	---	---
Gn: GANNETT-----	5w	---	---	---	---	---	---	---
IdB: INAVALE-----	4e	3e	1.50	4.00	30.00	100.00	20.00	---
IkB: INAVALE-----	6w	---	---	---	---	---	---	---
In: INAVALE-----	6w	---	---	---	---	---	---	---
BARNEY-----	6w	---	---	---	---	---	---	---
IpB: IPAGE-----	6e	4e	---	3.90	---	100.00	---	---
IsB: IPAGE-----	4e	4e	1.20	4.50	31.00	125.00	20.00	---
Jn: JANSEN-----	2e	2e	1.80	5.00	37.00	136.00	25.00	---

## LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued

Brown County, Nebraska

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Alfalfa hay		Corn		Winter wheat	
	N	I	N	I	N	I	N	I
			Tons		Bu		Bu	
JnC: JANSEN-----	3e	3e	1.60	4.80	36.00	130.00	23.00	---
Jo: JANSEN-----	2s	2s	1.80	5.00	40.00	140.00	26.00	---
JoC: JANSEN-----	3e	3e	1.60	4.80	36.00	135.00	23.00	---
Jr: JANSEN-----	2e	2e	1.30	3.50	32.00	106.00	22.00	---
MEADIN-----	6s	4s	1.30	3.50	32.00	106.00	22.00	---
JtB: JANSEN-----	2e	2e	1.90	5.10	36.00	132.00	24.00	---
SANDOSE-----	3e	3e	1.90	5.10	36.00	132.00	24.00	---
Jw: JOHNSTOWN-----	2e	2e	2.10	6.10	56.00	145.00	37.00	---
JwB: JOHNSTOWN-----	2e	2e	2.00	6.00	53.00	143.00	34.00	---
Jy: JOHNSTOWN-----	2c	1	2.50	6.30	60.00	150.00	40.00	---
JyB: JOHNSTOWN-----	2e	2e	2.30	6.00	56.00	145.00	37.00	---
JyC: JOHNSTOWN-----	3e	3e	2.10	5.60	50.00	135.00	33.00	---
LcG: LABU-----	6e	---	---	---	---	---	---	---
SANSARC-----	6s	---	---	---	---	---	---	---
LfB: LIBORY-----	3e	3e	3.40	5.20	48.00	125.00	28.00	---
Lo: LOUP-----	5w	---	---	---	---	---	---	---
Lp: LOUP-----	5w	---	---	---	---	---	---	---
LtB: ELS-----	4w	4w	---	---	---	---	---	---
LOUP-----	5w	---	---	---	---	---	---	---
Ma: MARLAKE-----	8w	---	---	---	---	---	---	---
McG: MCKELVIE-----	7e	---	---	---	---	---	---	---
LONGPINE-----	7s	---	---	---	---	---	---	---
MeB: MEADIN-----	6s	4s	---	2.00	---	75.00	---	---
MeF: MEADIN-----	6s	---	---	---	---	---	---	---
Oe: O'NEILL-----	3e	3e	1.50	4.80	34.00	133.00	22.00	---
On: O'NEILL-----	2c	2s	1.70	5.00	36.00	135.00	24.00	---
OsC: O'NEILL-----	4e	4e	1.20	3.60	29.00	110.00	16.00	---
MEADIN-----	6s	---	1.20	3.60	29.00	110.00	16.00	---
OsD: O'NEILL-----	6e	4e	---	3.30	---	95.00	---	---

## LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued

Brown County, Nebraska

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Alfalfa hay		Corn		Winter wheat	
	N	I	N	I	N	I	N	I
			Tons		Bu		Bu	
MEADIN-----	6s	---	---	3.30	---	95.00	---	---
Pg: PITS-----	8s	---	---	---	---	---	---	---
PtB: PIVOT-----	4e	3e	1.40	4.00	31.00	115.00	18.00	---
RtB: RONSON-----	3e	3e	1.50	4.00	22.00	90.00	16.00	---
LONGPINE-----	6s	---	1.50	4.00	22.00	90.00	16.00	---
RtC: RONSON-----	3e	3e	1.40	3.90	18.00	85.00	15.00	---
LONGPINE-----	6s	---	1.40	3.90	18.00	85.00	15.00	---
RtD: RONSON-----	4e	4e	---	---	---	---	---	---
LONGPINE-----	6s	---	---	---	---	---	---	---
ScB: SANDOSE-----	3e	3e	2.00	5.20	42.00	130.00	24.00	---
SkB: SIMEON-----	6s	4s	---	2.90	---	90.00	---	---
SkD: SIMEON-----	6s	---	---	---	---	---	---	---
SvD: SIMEON-----	6s	4s	---	3.00	---	86.00	---	---
VALENTINE-----	6e	4e	---	3.00	---	86.00	---	---
Tn: TRYON-----	5w	---	---	---	---	---	---	---
To: TRYON-----	5w	---	---	---	---	---	---	---
TpB: TRYON-----	5w	---	---	---	---	---	---	---
ELS-----	6e	4w	---	---	---	---	---	---
VaB: VALENTINE-----	6e	4e	---	3.20	---	80.00	---	---
VaD: VALENTINE-----	6e	4e	---	3.20	---	80.00	---	---
VaE: VALENTINE-----	6e	---	---	---	---	---	---	---
VaF: VALENTINE-----	6e	---	---	---	---	---	---	---
VALENTINE-----	7e	---	---	---	---	---	---	---
VbB: VALENTINE-----	4e	4e	0.80	3.60	30.00	100.00	17.00	---
VbD: VALENTINE-----	6e	4e	---	3.20	---	90.00	---	---
VfD: VALENTINE-----	6e	4e	---	3.20	---	82.00	---	---
ELS-----	6e	4w	---	3.20	---	82.00	---	---
VhD: VALENTINE-----	6e	4e	---	3.80	---	96.00	---	---
LIBORY-----	3e	3e	---	3.80	---	96.00	---	---
VpD: VALENTINE-----	6e	4e	---	3.40	---	93.00	---	---

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued  
Brown County, Nebraska

PAGE 5 of 5

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

Map symbol and soil name	Land Capability		Alfalfa hay		Corn		Winter wheat	
	N	I	N	I	N	I	N	I
			Tons		Bu		Bu	
PIVOT-----	4e	3e	---	3.40	---	93.00	---	---
VrD: VALENTINE-----	6e	4e	---	3.80	---	106.00	---	---
SANDOSE-----	3e	3e	---	3.80	---	106.00	---	---
VsG2: VALENTINE-----	7e	---	---	---	---	---	---	---
SIMEON-----	6s	---	---	---	---	---	---	---
VtE: VALENTINE-----	6e	---	---	---	---	---	---	---
LONGPINE-----	6s	---	---	---	---	---	---	---
VwE: VALENTINE-----	6e	---	---	---	---	---	---	---
TRYON-----	5w	---	---	---	---	---	---	---
VxB: VETAL-----	2e	2e	2.50	5.50	50.00	140.00	32.00	---
zwa: WATER-----	---	---	---	---	---	---	---	---
zwb: WATER-----	---	---	---	---	---	---	---	---



Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
An	Anselmo fine sandy loam, 0 to 2 percent slopes	Prime farmland if irrigated
AnC	Anselmo fine sandy loam, 2 to 6 percent slopes	Prime farmland if irrigated
Bo	Brocksburg loam, 0 to 1 percent slopes	Prime farmland if irrigated
Jn	Jansen fine sandy loam, 0 to 2 percent slopes	Prime farmland if irrigated
JnC	Jansen fine sandy loam, 2 to 6 percent slopes	Prime farmland if irrigated
Jo	Jansen loam, 0 to 2 percent slopes	Prime farmland if irrigated
JoC	Jansen loam, 2 to 6 percent slopes	Prime farmland if irrigated
Jw	Johnstown fine sandy loam, 0 to 1 percent slopes	Prime farmland if irrigated
JwB	Johnstown fine sandy loam, 1 to 3 percent slopes	Prime farmland if irrigated
Jy	Johnstown loam, 0 to 1 percent slopes	Prime farmland if irrigated
JyB	Johnstown loam, 1 to 3 percent slopes	Prime farmland if irrigated
JyC	Johnstown loam, 3 to 6 percent slopes	Prime farmland if irrigated
VxB	Vetal loam, 1 to 3 percent slopes	Prime farmland if irrigated





The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
Aa	Almeria Loamy Fine Sand, 0 To 2 Percent Slopes-----	23
Ae	Almeria Fine Sandy Loam, 0 To 2 Percent Slopes-----	24
Af	Almeria-Histosols Complex, Channeled-----	13
An	Anselmo Fine Sandy Loam, 0 To 2 Percent Slopes-----	50
AnC	Anselmo Fine Sandy Loam, 2 To 6 Percent Slopes-----	48
AnD	Anselmo Fine Sandy Loam, 6 To 11 Percent Slopes-----	44
AtF	Anselmo-Brunswick Fine Sandy Loams, 11 To 30 Percent Slopes-----	19
Ba	Barney Fine Sandy Loam, Channeled-----	21
Bd	Bolent Fine Sandy Loam, 0 To 2 Percent Slopes-----	26
Bo	Brocksburg Loam, 0 To 1 Percent Slopes-----	60
BrD	Brunswick Fine Sandy Loam, 3 To 9 Percent Slopes-----	29
DuB	Dunday Loamy Fine Sand, 0 To 3 Percent Slopes-----	31
DuD	Dunday Loamy Fine Sand, 3 To 9 Percent Slopes-----	29
Eo	Els Loamy Sand, 0 To 2 Percent Slopes-----	27
EpB	Els-Ipage Fine Sands, 0 To 3 Percent Slopes-----	24
Es	Elsmere Loamy Fine Sand, 0 To 2 Percent Slopes-----	27
Fe	Fluvaquents, Sandy-----	4
Gn	Gannett Fine Sandy Loam, 0 To 2 Percent Slopes-----	41
IdB	Inavale Loamy Fine Sand, 0 To 3 Percent Slopes-----	31
IkB	Inavale Sand, Channeled-----	28
In	Inavale-Barney Complex, Channeled-----	26
IpB	Ipage Fine Sand, 0 To 3 Percent Slopes-----	25
IsB	Ipage Loamy Sand, 0 To 3 Percent Slopes-----	27
Jn	Jansen Fine Sandy Loam, 0 To 2 Percent Slopes-----	50
JnC	Jansen Fine Sandy Loam, 2 To 6 Percent Slopes-----	49
Jo	Jansen Loam, 0 To 2 Percent Slopes-----	53
JoC	Jansen Loam, 2 To 6 Percent Slopes-----	51
Jr	Jansen-Meadin Complex, 0 To 2 Percent Slopes-----	35
JtB	Jansen-Sandose Complex, 0 To 3 Percent Slopes-----	48
Jw	Johnstown Fine Sandy Loam, 0 To 1 Percent Slopes-----	68
JwB	Johnstown Fine Sandy Loam, 1 To 3 Percent Slopes-----	67
Jy	Johnstown Loam, 0 To 1 Percent Slopes-----	69
JyB	Johnstown Loam, 1 To 3 Percent Slopes-----	68
JyC	Johnstown Loam, 3 To 6 Percent Slopes-----	65
LcG	Labu-Sansarc Silty Clays, 11 To 40 Percent Slopes-----	7
LfB	Libory Loamy Fine Sand, 0 To 3 Percent Slopes-----	51
Lo	Loup Fine Sandy Loam, 0 To 2 Percent Slopes-----	30
Lp	Loup Fine Sandy Loam, Wet, 0 To 2 Percent Slopes-----	20
LtB	Loup-Els Complex, 0 To 3 Percent Slopes-----	28
Ma	Marlake Fine Sandy Loam, 0 To 1 Percent Slopes-----	3
McG	Mckelvie-Tassel-Ronson Complex, 15 To 70 Percent Slopes-----	3
MeB	Meadin Sandy Loam, 0 To 3 Percent Slopes-----	29
MeF	Meadin Sandy Loam, 3 To 30 Percent Slopes-----	21
Oe	O'Neill Fine Sandy Loam, 0 To 2 Percent Slopes-----	37
On	O'Neill Loam, 0 To 2 Percent Slopes-----	40
OsC	O'Neill-Meadin Sandy Loams, 2 To 6 Percent Slopes-----	34
OsD	O'Neill-Meadin Sandy Loams, 6 To 11 Percent Slopes-----	29
Pg	Pits, Sand And Gravel-----	16
PtB	Pivot Loamy Sand, 0 To 3 Percent Slopes-----	28
RtB	Ronson-Tassel Fine Sandy Loams, 0 To 3 Percent Slopes-----	17
RtC	Ronson-Tassel Fine Sandy Loams, 3 To 6 Percent Slopes-----	17
RtD	Ronson-Tassel Fine Sandy Loams, 6 To 11 Percent Slopes-----	14
ScB	Sandose Loamy Fine Sand, 0 To 3 Percent Slopes-----	47
SkB	Simeon Loamy Sand, 0 To 3 Percent Slopes-----	26
SkD	Simeon Loamy Sand, 3 To 9 Percent Slopes-----	25
SvD	Simeon-Valentine Fine Sands, 0 To 9 Percent Slopes-----	24
Tn	Tryon Loamy Fine Sand, 0 To 2 Percent Slopes-----	25
To	Tryon Loamy Fine Sand, Wet, 0 To 2 Percent Slopes-----	17
TpB	Tryon-Els Complex, 0 To 3 Percent Slopes-----	24
VaB	Valentine Fine Sand, 0 To 3 Percent Slopes-----	23
VaD	Valentine Fine Sand, 3 To 9 Percent Slopes-----	21
VaE	Valentine Fine Sand, Rolling-----	16
VaF	Valentine Fine Sand, Rolling And Hilly-----	12
VbB	Valentine Loamy Fine Sand, 0 To 3 Percent Slopes-----	27
VbD	Valentine Loamy Fine Sand, 3 To 9 Percent Slopes-----	25
VfD	Valentine-Els Fine Sands, 0 To 9 Percent Slopes-----	23
VhD	Valentine-Libory Complex, 0 To 9 Percent Slopes-----	34
VpD	Valentine-Pivot Complex, 0 To 9 Percent Slopes-----	24
VrD	Valentine-Sandose Loamy Fine Sands, 0 To 9 Percent Slopes-----	33
VsG2	Valentine-Simeon Complex, 9 To 40 Percent Slopes, Eroded-----	10
VtE	Valentine-Tassel Complex, 3 To 17 Percent Slopes-----	15
VwE	Valentine-Tryon Complex, 0 To 17 Percent Slopes-----	19
VxB	Vetal Loam, 1 To 3 Percent Slopes-----	57
zwa	Water > 40 Acres-----	0
zwb	Water < 40 Acres-----	0



Brown County, Nebraska: Published  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Aa:ALMERIA-----	100	N/A	5w	Not prime farmland	D	Wet Subirrigated - Veg. Zone 3		.17	.17	5	8	0
Ae:ALMERIA-----	100	N/A	5w	Not prime farmland	D	Wet Subirrigated		.24	.24	5	8	0
Af:ALMERIA-----	55	N/A	6w	Not prime farmland	D	Wet Land		.24	.24	5	8	0
Af:HISTOSOLS----	45	N/A	8w	Not prime farmland	D	Unspecified		---	---	3	8	0
An:ANSELMO-----	100	2e-	2e	Prime farmland if irrigated	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
AnC:ANSELMO-----	100	3e-	3e	Prime farmland if irrigated	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
AnD:ANSELMO-----	100	4e-	4e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
AtF:ANSELMO-----	60	N/A	6e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	5	3	86
AtF:BRUNSWICK---	40	N/A	6e	Not prime farmland	B	Sandy - Veg. Zone 3		.24	.24	3	3	86
Ba:BARNEY-----	100	N/A	6w	Not prime farmland	D	Wet Land		.20	.20	5	8	0
Bd:BOLENT-----	100	4w-	4w	Not prime farmland	A	Subirrigated - Veg. Zone 3		.24	.24	5	3	86
Bo:BROCKSBURG---	100	2s-	2s	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.28	.28	4	5	56
BrD:BRUNSWICK---	100	4e-	4e	Not prime farmland	B	Sandy - Veg. Zone 3		.24	.24	3	3	86
DuB:DUNDAY-----	100	3e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
DuD:DUNDAY-----	100	4e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
Eo:ELS-----	100	4w-	4w	Not prime farmland	A	Subirrigated		.17	.17	5	2	134
EpB:ELS-----	65	4w-	6e	Not prime farmland	A	Subirrigated		.15	.15	5	1	220
EpB:IPAGE-----	35	4e-	6e	Not prime farmland	A	Sandy Lowland 22-25" P.z.		.15	.15	5	1	220
Es:ELSMERE-----	100	4w-	4w	Not prime farmland	A	Subirrigated		.17	.17	5	2	134

Brown County, Nebraska: Published  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Fe:FLUVAQUENTS--	100	N/A	8w	Not prime farmland	D	Unspecified		.17	.17	5	8	0
Gn:GANNETT-----	100	N/A	5w	Not prime farmland	D	Wet Subirrigated		.20	.20	4	8	0
IdB:INAVALE-----	100	3e-	4e	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.17	.17	5	2	134
IkB:INAVALE-----	100	N/A	6w	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.15	.15	5	1	220
In:INAVALE-----	60	N/A	6w	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.17	.17	5	2	134
In:BARNEY-----	40	N/A	6w	Not prime farmland	D	Wet Land - Veg. Zone 3		.20	.20	5	8	0
IpB:IPAGE-----	100	4e-	6e	Not prime farmland	A	Sandy Lowland 22-25" P.z.		.15	.15	5	1	220
IsB:IPAGE-----	100	4e-	4e	Not prime farmland	A	Sandy Lowland - Veg. Zone 3		.17	.17	5	2	134
Jn:JANSEN-----	100	2e-	2e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.20	.20	4	3	86
JnC:JANSEN-----	100	3e-	3e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.20	.20	4	3	86
Jo:JANSEN-----	100	2s-	2s	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.28	.28	4	5	56
JoC:JANSEN-----	100	3e-	3e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.28	.28	4	5	56
Jr:JANSEN-----	50	2e-	2e	Not prime farmland	B	Silty - Veg. Zone 3		.20	.20	4	3	86
Jr:MEADIN-----	50	4s-	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86
JtB:JANSEN-----	50	2e-	2e	Not prime farmland	B	Silty - Veg. Zone 3		.20	.20	4	3	86
JtB:SANDOSE-----	50	3e-	3e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
Jw:JOHNSTOWN----	100	2e-	2e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.20	.20	4	3	86
JwB:JOHNSTOWN---	100	2e-	2e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.20	.20	4	3	86

Brown County, Nebraska: Published  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Jy:JOHNSTOWN----	100	1-	2c	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.28	.28	4	5	56
JyB:JOHNSTOWN---	100	2e-	2e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.28	.28	4	5	56
JyC:JOHNSTOWN---	100	3e-	3e	Prime farmland if irrigated	B	Silty - Veg. Zone 3		.28	.28	4	5	56
LcG:LABU-----	55	N/A	6e	Not prime farmland	D	Clayey - Veg. Zone 3		.32	.32	3	4	86
LcG:SANSARC----	45	N/A	6s	Not prime farmland	D	Shallow Clay - Veg. Zone 3		.37	.37	2	4	86
LfB:LIBORY-----	100	3e-	3e	Not prime farmland	A	Sandy Lowland 22-25" P.z.		.17	.17	5	2	134
Lo:LOUP-----	100	N/A	5w	Not prime farmland	D	Wet Subirrigated		.20	.20	3	8	0
Lp:LOUP-----	100	N/A	5w	Not prime farmland	D	Wet Land		.20	.20	3	8	0
LtB:LOUP-----	50	N/A	5w	Not prime farmland	D	Wet Subirrigated		.20	.20	3	8	0
LtB:ELS-----	50	4w-	4w	Not prime farmland	A	Subirrigated		.17	.17	5	2	134
Ma:MARLAKE-----	100	N/A	8w	Not prime farmland	D	Unspecified		.20	.20	5	8	0
McG:MCKELVIE----	50	N/A	7e	Not prime farmland	A	Sands - Veg. Zone 3		.17	.17	5	2	134
McG:LONGPINE----	30	N/A	7s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.24	.24	2	3	86
MeB:MEADIN-----	100	4s-	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86
MeF:MEADIN-----	100	N/A	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86
Oe:O'NEILL-----	100	3e-	3e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
On:O'NEILL-----	100	2s-	2c	Not prime farmland	B	Sandy - Veg. Zone 3		.28	.28	4	5	56
OsC:O'NEILL-----	60	4e-	4e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
OsC:MEADIN-----	40	N/A	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86

Brown County, Nebraska: Published  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
OsD:O'NEILL-----	55	4e-	6e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	4	3	86
OsD:MEADIN-----	45	N/A	6s	Not prime farmland	A	Shallow To Gravel - Veg. Zone 3		.20	.20	5	3	86
Pg:PITS-----	100	N/A	8s	Not prime farmland	A	Unspecified		.15	.15	2	8	0
PtB:PIVOT-----	100	3e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
RtB:RONSON-----	55	3e-	3e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	3	3	86
RtB:LONGPINE----	45	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.24	.24	2	3	86
RtC:RONSON-----	55	3e-	3e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	3	3	86
RtC:LONGPINE----	45	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.24	.24	2	3	86
RtD:RONSON-----	55	4e-	4e	Not prime farmland	B	Sandy - Veg. Zone 3		.20	.20	3	3	86
RtD:LONGPINE----	45	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.24	.24	2	3	86
ScB:SANDOSE-----	100	3e-	3e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
SkB:SIMEON-----	100	4s-	6s	Not prime farmland	A	Shallow To Gravel 17-25" P.Z.		.17	.17	5	2	134
SkD:SIMEON-----	100	N/A	6s	Not prime farmland	A	Shallow To Gravel 17-25" P.Z.		.17	.17	5	2	134
SvD:SIMEON-----	60	4s-	6s	Not prime farmland	A	Shallow To Gravel 17-25" P.Z.		.15	.15	5	1	250
SvD:VALENTINE---	40	4e-	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
Tn:TRYON-----	100	N/A	5w	Not prime farmland	D	Wet Subirrigated		.17	.17	5	8	0
To:TRYON-----	100	N/A	5w	Not prime farmland	D	Wet Land		.17	.17	5	8	0
TpB:TRYON-----	55	N/A	5w	Not prime farmland	D	Wet Subirrigated		.17	.17	5	8	0
TpB:ELS-----	45	4w-	6e	Not prime farmland	A	Subirrigated		.15	.15	5	1	220

Brown County, Nebraska: Published  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
VaB:VALENTINE---	100	4e-	6e	Not prime farmland	A	Sandy 22-25" P.Z.		.15	.15	5	1	250
VaD:VALENTINE---	100	4e-	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
VaE:VALENTINE---	100	N/A	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
VaF:VALENTINE---	70	N/A	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
	30	N/A	7e	Not prime farmland	A	Choppy Sands 22- 25" P.Z.		.15	.15	5	1	250
VbB:VALENTINE---	100	4e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
VbD:VALENTINE---	100	4e-	6e	Not prime farmland	A	Sands 22-25" P.Z.		.17	.17	5	2	134
VfD:VALENTINE---	55	4e-	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
VfD:ELS-----	45	4w-	6e	Not prime farmland	A	Subirrigated		.15	.15	5	1	220
VhD:VALENTINE---	60	4e-	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
VhD:LIBORY-----	40	3e-	3e	Not prime farmland	A	Sandy Lowland 22-25" P.Z.		.17	.17	5	2	134
VpD:VALENTINE---	60	4e-	6e	Not prime farmland	A	Sands - Veg. Zone 3		.15	.15	5	1	250
VpD:PIVOT-----	40	3e-	4e	Not prime farmland	A	Sandy - Veg. Zone 3		.17	.17	5	2	134
VrD:VALENTINE---	65	4e-	6e	Not prime farmland	A	Sands 22-25" P.Z.		.17	.17	5	2	134
VrD:SANDOSE-----	35	3e-	3e	Not prime farmland	A	Sandy 22-25" P.Z.		.17	.17	5	2	134
VsG2:VALENTINE--	65	N/A	7e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250
VsG2:SIMEON-----	35	N/A	6s	Not prime farmland	A	Shallow To Gravel 17-25" P.Z.		.17	.17	5	2	134
VtE:VALENTINE---	60	N/A	6e	Not prime farmland	A	Sands - Veg. Zone 3		.15	.15	5	1	250
VtE:LONGPINE----	40	N/A	6s	Not prime farmland	D	Shallow Limy - Veg. Zone 3		.24	.24	2	3	86
VwE:VALENTINE---	65	N/A	6e	Not prime farmland	A	Sands 22-25" P.Z.		.15	.15	5	1	250

Brown County, Nebraska: Published  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
VwE:TRYON-----	35	N/A	5w	Not prime farmland	D	Wet Land		.17	.17	5	8	0
VxB:VETAL-----	100	2e-	2e	Prime farmland if irrigated	B	Sandy - Veg. Zone 3		.28	.28	5	5	56
zwa:WATER-----	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---
zwb:WATER-----	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---





RANGELAND PRODUCTIVITY  
Brown County, Nebraska

# Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

**Rangeland.** Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

**Grazed Forest Land.** Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

**Native Pasture.** Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

## Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued  
Brown County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Aa:				
Almeria-----	Wet Subirrigated - Veg. Zone 3	5,800	5,200	4,700
Ae:				
Almeria-----	Wet Subirrigated	5,800	5,200	4,700
Af:				
Almeria-----	Wet Land	6,000	5,400	5,000
Histosols-----	---	---	---	---
An:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
AnC:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
AnD:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
AtF:				
Anselmo-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Brunswick-----	Sandy - Veg. Zone 3	3,000	2,600	2,200
Ba:				
Barney-----	Wet Land	6,000	5,400	5,000
Bd:				
Bolent-----	Subirrigated - Veg. Zone 3	5,500	5,000	4,200
Bo:				
Brocksburg-----	Silty - Veg. Zone 3	3,700	3,200	2,700
BrD:				
Brunswick-----	Sandy - Veg. Zone 3	3,000	2,600	2,200
DuB:				
Dunday-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
DuD:				
Dunday-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
Eo:				
Els-----	Subirrigated	5,500	5,300	5,000
EpB:				
Els-----	Subirrigated	5,500	5,300	5,000
Ipage-----	Sandy Lowland 22-25" P.z.	3,500	3,200	3,000
Es:				
Elsmere-----	Subirrigated	5,500	5,300	5,000
Fe:				
Fluvaquents-----	---	---	---	---
Gn:				
Gannett-----	Wet Subirrigated	5,800	5,500	5,300
IdB:				
Inavale-----	Sandy Lowland - Veg. Zone 3	3,500	3,000	2,200
IkB:				
Inavale-----	Sandy Lowland - Veg. Zone 3	3,300	2,800	2,200
In:				
Inavale-----	Sandy Lowland - Veg. Zone 3	3,300	2,800	2,200
Barney-----	Wet Land - Veg. Zone 3	6,000	5,400	5,000
IpB:				
Ipape-----	Sandy Lowland 22-25" P.z.	3,500	3,200	3,000
IsB:				
Ipape-----	Sandy Lowland - Veg. Zone 3	3,500	3,200	3,000
Jn:				
Jansen-----	Silty - Veg. Zone 3	3,500	3,300	3,000
JnC:				
Jansen-----	Silty - Veg. Zone 3	3,500	3,300	3,000
Jo:				
Jansen-----	Silty - Veg. Zone 3	3,700	3,200	2,700
JoC:				
Jansen-----	Silty - Veg. Zone 3	3,700	3,200	2,700
Jr:				
Jansen-----	Silty - Veg. Zone 3	3,500	3,300	3,000
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
JtB:				
Jansen-----	Silty - Veg. Zone 3	3,500	3,300	3,000
Sandose-----	Sandy - Veg. Zone 3	3,500	3,000	2,200
Jw:				
Johnstown-----	Silty - Veg. Zone 3	3,800	3,500	3,000
JwB:				
Johnstown-----	Silty - Veg. Zone 3	3,800	3,500	3,000
Jy:				
Johnstown-----	Silty - Veg. Zone 3	3,800	3,500	3,000
JyB:				
Johnstown-----	Silty - Veg. Zone 3	3,800	3,500	3,000
JyC:				
Johnstown-----	Silty - Veg. Zone 3	3,800	3,500	3,000
LcG:				
Labu-----	Clayey - Veg. Zone 3	3,400	3,000	2,600
Sansarc-----	Shallow Clay - Veg. Zone 3	2,500	2,100	1,500
LfB:				
Libory-----	Sandy Lowland 22-25" P.z.	4,300	3,500	2,700
Lo:				
Loup-----	Wet Subirrigated	5,800	5,500	5,300
Lp:				
Loup-----	Wet Land	6,000	5,600	5,300
LtB:				

RANGELAND PRODUCTIVITY--Continued  
Brown County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Els-----	Subirrigated	5,500	5,300	5,000
Loup-----	Wet Subirrigated	5,800	5,500	5,300
Ma:				
Marlake-----	---	---	---	---
McG:				
Mckelvie-----	Sands - Veg. Zone 3	2,500	2,000	1,500
Longpine-----	Shallow Limy - Veg. Zone 3	2,100	1,800	1,600
MeB:				
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
MeF:				
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
Oe:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
On:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
OsC:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
OsD:				
O'Neill-----	Sandy - Veg. Zone 3	3,500	3,300	3,000
Meadin-----	Shallow To Gravel - Veg. Zone 3	1,500	1,300	1,100
Pg:				
Pits-----	---	---	---	---
PtB:				
Pivot-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
RtB:				
Ronson-----	Sandy - Veg. Zone 3	3,000	2,800	1,700
Longpine-----	Shallow Limy - Veg. Zone 3	2,100	1,800	1,600
RtC:				
Ronson-----	Sandy - Veg. Zone 3	3,000	2,800	1,700
Longpine-----	Shallow Limy - Veg. Zone 3	2,100	1,800	1,600
RtD:				
Ronson-----	Sandy - Veg. Zone 3	3,000	2,800	1,700
Longpine-----	Shallow Limy - Veg. Zone 3	2,100	1,800	1,600
ScB:				
Sandose-----	Sandy - Veg. Zone 3	3,500	3,000	2,200
SkB:				
Simeon-----	Shallow To Gravel 17-25" P.z.	1,800	1,600	1,100
SkD:				
Simeon-----	Shallow To Gravel 17-25" P.z.	1,400	1,200	700
SvD:				
Simeon-----	Shallow To Gravel 17-25" P.z.	1,500	1,300	800
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
Tn:				
Tryon-----	Wet Subirrigated	5,800	5,500	5,300
To:				
Tryon-----	Wet Land	6,000	5,800	5,500
TpB:				
Tryon-----	Wet Subirrigated	5,800	5,500	5,300
Els-----	Subirrigated	5,500	5,300	5,000
VaB:				
Valentine-----	Sandy 22-25" P.z.	3,300	3,000	2,600
VaD:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
VaE:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
VaF:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
Valentine-----	Choppy Sands 22-25" P.z.	2,800	2,400	2,000
VbB:				
Valentine-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
VbD:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
VfD:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
Els-----	Subirrigated	5,500	5,300	5,000
VhD:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
Libory-----	Sandy Lowland 22-25" P.z.	4,300	3,500	2,700
VpD:				
Valentine-----	Sands - Veg. Zone 3	3,000	2,600	2,200
Pivot-----	Sandy - Veg. Zone 3	3,300	3,000	2,600
VrD:				
Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
Sandose-----	Sandy 22-25" P.z.	3,500	3,000	2,200
VsG2:				
Valentine-----	Sands 22-25" P.z.	2,800	2,400	2,000
Simeon-----	Shallow To Gravel 17-25" P.z.	1,400	1,200	700
VtE:				
Valentine-----	Sands - Veg. Zone 3	3,000	2,600	2,200
Longpine-----	Shallow Limy - Veg. Zone 3	2,100	1,800	1,600

RANGELAND PRODUCTIVITY--Continued  
Brown County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
VwE: Valentine-----	Sands 22-25" P.z.	3,000	2,600	2,200
Tryon-----	Wet Land	6,000	5,800	5,500
VxB: Vetal-----	Sandy - Veg. Zone 3	3,000	2,300	1,700
zwa: Water-----	---	---	---	---
zwb: Water-----	---	---	---	---



BUILDING SITE DEVELOPMENT  
Brown County, Nebraska

### Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. These tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Ae: Almeria-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Af: Almeria-----	55	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Histosols-----	45	Very limited Subsidence Flooding Depth to saturated zone Content of organic matter	1.00 1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone Content of organic matter	1.00 1.00 1.00
An: Anselmo-----	100	Not limited		Not limited		Not limited	
AnC: Anselmo-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
AnD: Anselmo-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
AtF: Anselmo-----	60	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Brunswick-----	40	Very limited Slope	1.00	Very limited Slope Depth to soft bedrock	1.00 0.42	Very limited Slope	1.00
Ba: Barney-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Bd: Bolent-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Bo: Brocksburg-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
BrD: Brunswick-----	100	Not limited		Somewhat limited Depth to soft bedrock	0.42	Somewhat limited Slope	0.48
DuB: Dunday-----	100	Not limited		Not limited		Not limited	
DuD: Dunday-----	100	Not limited		Not limited		Somewhat limited Slope	0.48
Eo: Els-----	100	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
EpB: Els-----	65	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
Ipage-----	35	Not limited		Somewhat limited Depth to saturated zone	0.61	Not limited	
Es: Elsmere-----	100	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07



BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Fe: Fluvaquents-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Gn: Gannett-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
IdB: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
IkB: Inavale-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
In: Inavale-----	60	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Barney-----	40	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
IpB: Ipage-----	100	Not limited		Somewhat limited Depth to saturated zone	0.61	Not limited	
IsB: Ipage-----	100	Not limited		Somewhat limited Depth to saturated zone	0.61	Not limited	
Jn: Jansen-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
JnC: Jansen-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell Slope	0.50 0.00
Jo: Jansen-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
JoC: Jansen-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell Slope	0.50 0.00
Jr: Jansen-----	50	Not limited		Not limited		Not limited	
Meadin-----	50	Not limited		Not limited		Not limited	
JtB: Jansen-----	50	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
Sandose-----	50	Not limited		Not limited		Not limited	
Jw: Johnstown-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
JwB: Johnstown-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Jy: Johnstown-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
JyB: Johnstown-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
JyC: Johnstown-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LcG: Labu-----	55	Very limited Shrink-swell Slope	1.00 1.00	Very limited Shrink-swell Slope Depth to soft bedrock	1.00 1.00 0.42	Very limited Slope Shrink-swell	1.00 1.00
Sansarc-----	45	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 1.00 1.00
LfB: Libory-----	100	Somewhat limited Depth to saturated zone	0.98	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.98
Lo: Loup-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Lp: Loup-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
LtB: Els-----	50	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
Loup-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Ma: Marlake-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
McG: Mckelvie-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Longpine-----	30	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Slope Depth to soft bedrock	1.00 1.00
MeB: Meadin-----	100	Not limited		Not limited		Not limited	
MeF: Meadin-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Oe: O'Neill-----	100	Not limited		Not limited		Not limited	
On: O'Neill-----	100	Not limited		Not limited		Not limited	
OsC: O'Neill-----	60	Not limited		Not limited		Somewhat limited Slope	0.00
Meadin-----	40	Not limited		Not limited		Somewhat limited Slope	0.00
OsD: O'Neill-----	55	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Meadin-----	45	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Pg: Pits-----	100	Not rated		Not rated		Not rated	
PtB: Pivot-----	100	Not limited		Not limited		Not limited	
RtB: Ronson-----	55	Not limited		Somewhat limited Depth to soft bedrock	0.42	Not limited	
Longpine-----	45	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock	1.00

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
RtC: Ronson-----	55	Not limited		Somewhat limited Depth to soft bedrock	0.42	Somewhat limited Slope	0.12
Longpine-----	45	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock Slope	0.12
RtD: Ronson-----	55	Somewhat limited Slope	0.04	Somewhat limited Depth to soft bedrock Slope	0.42	Very limited Slope	1.00
Longpine-----	45	Somewhat limited Depth to soft bedrock Slope	1.00	Very limited Depth to soft bedrock Slope	0.04	Very limited Depth to soft bedrock Slope	1.00
ScB: Sandose-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
SkB: Simeon-----	100	Not limited		Not limited		Not limited	
SkD: Simeon-----	100	Not limited		Not limited		Somewhat limited Slope	0.48
SvD: Simeon-----	60	Not limited		Not limited		Not limited	
Valentine-----	40	Not limited		Not limited		Somewhat limited Slope	0.48
Tn: Tryon-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
To: Tryon-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
TpB: Tryon-----	55	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Els-----	45	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
VaB: Valentine-----	100	Not limited		Not limited		Not limited	
VaD: Valentine-----	100	Not limited		Not limited		Somewhat limited Slope	0.48
VaE: Valentine-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
VaF: Valentine-----	70	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Valentine-----	30	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
VbB: Valentine-----	100	Not limited		Not limited		Not limited	
VbD: Valentine-----	100	Not limited		Not limited		Somewhat limited Slope	0.48
VfD: Valentine-----	55	Not limited		Not limited		Somewhat limited Slope	0.48
Els-----	45	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
VhD: Valentine-----	60	Not limited		Not limited		Somewhat limited Slope	0.48
Libory-----	40	Somewhat limited Depth to saturated zone	0.98	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.98
VpD: Valentine-----	60	Not limited		Not limited		Somewhat limited Slope	0.48
Pivot-----	40	Not limited		Not limited		Not limited	

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VrD: Valentine-----	65	Not limited		Not limited		Somewhat limited Slope	0.48
Sandose-----	35	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
VsG2: Valentine-----	65	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Simeon-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
VtE: Valentine-----	60	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Longpine-----	40	Somewhat limited Depth to soft bedrock Slope	1.00 0.16	Very limited Depth to soft bedrock Slope	1.00 0.16	Very limited Depth to soft bedrock Slope	1.00 1.00
VwE: Valentine-----	65	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Tryon-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
VxB: Vetal-----	100	Not limited		Not limited		Not limited	
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Frost action	0.50	Flooding	0.80	Droughty	0.05
Ae: Almeria-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Frost action	0.50	Flooding	0.80	Droughty	0.00
Af: Almeria-----	55	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Frost action	0.50	Flooding	0.80		
Histosols-----	45	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Subsidence	1.00	Cutbanks cave	1.00	Content of organic matter	1.00
		Flooding	1.00	Content of organic matter	1.00	Depth to saturated zone	1.00
An: Anselmo-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
AnC: Anselmo-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
AnD: Anselmo-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.04
		Slope	0.04	Slope	0.04		
AtF: Anselmo-----	60	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Cutbanks cave	1.00		
Brunswick-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to soft bedrock	0.42	Depth to bedrock	0.42
				Cutbanks cave	0.10	Droughty	0.05
Ba: Barney-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Frost action	0.50	Flooding	0.80	Droughty	0.02
				Depth to dense layer	0.50		
Bd: Bolent-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Somewhat limited Flooding	0.60
		Frost action	0.50	Depth to saturated zone	1.00	Droughty	0.09
		Depth to saturated zone	0.03	Flooding	0.60	Depth to saturated zone	0.03
Bo: Brocksburg-----	100	Somewhat limited Shrink-swell	0.50	Very limited Cutbanks cave	1.00	Not limited	
		Frost action	0.50				
BrD: Brunswick-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Depth to bedrock	0.42
				Depth to soft bedrock	0.42	Droughty	0.23
DuB: Dunday-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.04

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
DuD: Dunday-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.04
Eo: Els-----	100	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Droughty Depth to saturated zone	0.48 0.03
EpB: Els-----	65	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Droughty Depth to saturated zone	0.54 0.03
Ipage-----	35	Somewhat limited Frost action	0.50	Very limited Cutbanks cave Depth to saturated zone	1.00 0.61	Somewhat limited Droughty	0.63
Es: Elsmere-----	100	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Droughty Depth to saturated zone	0.41 0.03
Fe: Fluvaquents-----	100	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Gn: Gannett-----	100	Frost action	0.50	Flooding	0.80	Droughty	0.00
		Very limited Frost action Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
IdB: Inavale-----	100	Somewhat limited Flooding	0.40	Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.17
IkB: Inavale-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding Too sandy Droughty	0.60 0.50 0.34
In: Inavale-----	60	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding Droughty	0.60 0.14
Barney-----	40	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
		Frost action	0.50	Flooding Depth to dense layer	0.80 0.50	Droughty	0.07
IpB: Ipage-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave Depth to saturated zone	1.00 0.61	Somewhat limited Droughty	0.63
IsB: Ipage-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave Depth to saturated zone	1.00 0.61	Somewhat limited Droughty	0.43
Jn: Jansen-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
JnC: Jansen-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
Jo: Jansen-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
JoC: Jansen-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
Jr: Jansen-----	50	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Meadin-----	50	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.87
JtB: Jansen-----	50	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
Sandose-----	50	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.01
Jw: Johnstown-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
JwB: Johnstown-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
Jy: Johnstown-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
JyB: Johnstown-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
JyC: Johnstown-----	100	Somewhat limited Shrink-swell Frost action	0.50 0.50	Very limited Cutbanks cave	1.00	Not limited	
LcG: Labu-----	55	Very limited Shrink-swell Slope	1.00 1.00	Very limited Slope Too clayey Depth to soft bedrock Cutbanks cave	1.00 0.72 0.42 0.10	Very limited Too clayey Slope Droughty Depth to bedrock	1.00 1.00 0.85 0.42
Sansarc-----	45	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00 1.00	Very limited Depth to soft bedrock Too clayey Slope Cutbanks cave	1.00 1.00 1.00 1.00 0.10	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 1.00 1.00 1.00
LfB: Libory-----	100	Somewhat limited Depth to saturated zone	0.75	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Depth to saturated zone	0.75
Lo: Loup-----	100	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.03
Lp: Loup-----	100	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.03
LtB: Els-----	50	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Droughty Depth to saturated zone	0.48 0.03
Loup-----	50	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.01

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ma: Marlake-----	100	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
McG: Mckelvie-----	50	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.21
Longpine-----	30	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Droughty Slope	1.00 1.00 1.00
MeB: Meadin-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.73
MeF: Meadin-----	100	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.73
Oe: O'Neill-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.02
On: O'Neill-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
OsC: O'Neill-----	60	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Meadin-----	40	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.69
OsD: O'Neill-----	55	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Droughty Slope	0.08 0.04
Meadin-----	45	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Droughty Slope	0.63 0.04
Pg: Pits-----	100	Not rated		Not rated		Not rated	
PtB: Pivot-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.26
RtB: Ronson-----	55	Somewhat limited Frost action	0.50	Somewhat limited Depth to soft bedrock Cutbanks cave	0.42 0.10	Somewhat limited Depth to bedrock Droughty	0.42 0.10
Longpine-----	45	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock Droughty	1.00 1.00
RtC: Ronson-----	55	Somewhat limited Frost action	0.50	Somewhat limited Depth to soft bedrock Cutbanks cave	0.42 0.10	Somewhat limited Depth to bedrock Droughty	0.42 0.00
Longpine-----	45	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock Droughty	1.00 1.00
RtD: Ronson-----	55	Somewhat limited Frost action Slope	0.50 0.04	Somewhat limited Depth to soft bedrock Cutbanks cave Slope	0.42 0.10 0.04	Somewhat limited Depth to bedrock Droughty Slope	0.42 0.36 0.04
Longpine-----	45	Somewhat limited Depth to soft bedrock Slope	1.00 0.04	Very limited Depth to soft bedrock Cutbanks cave Slope	1.00 0.10 0.04	Very limited Depth to bedrock Droughty Slope	1.00 1.00 0.04



BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
ScB: Sandose-----	100	Somewhat limited Shrink-swell	0.50	Very limited Cutbanks cave	1.00	Not limited	
SkB: Simeon-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.22
SkD: Simeon-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.22
SvD: Simeon-----	60	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.30
Valentine-----	40	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.87
Tn: Tryon-----	100	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.79
To: Tryon-----	100	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.76
TpB: Tryon-----	55	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.79
Els-----	45	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Droughty Depth to saturated zone	0.54 0.03
VaB: Valentine-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.89
VaD: Valentine-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.89
VaE: Valentine-----	100	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.89
VaF: Valentine-----	70	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.89
Valentine-----	30	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Droughty	1.00 0.89
VbB: Valentine-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.52
VbD: Valentine-----	100	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.52
VfD: Valentine-----	55	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.89
Els-----	45	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Droughty Depth to saturated zone	0.54 0.03
VhD: Valentine-----	60	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.87
Libory-----	40	Somewhat limited Depth to saturated zone	0.75	Very limited Cutbanks cave Depth to saturated zone	1.00 1.00	Somewhat limited Depth to saturated zone	0.75
VpD: Valentine-----	60	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.88
Pivot-----	40	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.27

BUILDING SITE DEVELOPMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VrD: Valentine-----	65	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.63
Sandose-----	35	Somewhat limited Shrink-swell	0.50	Very limited Cutbanks cave	1.00	Not limited	
VsG2: Valentine-----	65	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00	Very limited Slope	1.00
Simeon-----	35	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00	Droughty	0.87
VtE: Valentine-----	60	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00	Somewhat limited Droughty	0.86
Longpine-----	40	Somewhat limited Depth to soft bedrock Slope	1.00 0.16	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 0.16 0.10	Very limited Depth to bedrock	1.00
VwE: Valentine-----	65	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Droughty Slope	1.00 0.16
Tryon-----	35	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Somewhat limited Droughty Slope	0.89 0.16
VxB: Vetal-----	100	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Very limited Depth to saturated zone Droughty	1.00 0.76
zwa: Water-----	100	Not rated		Not rated		Not limited	
zwb: Water-----	100	Not rated		Not rated		Not rated	



CONSTRUCTION MATERIALS  
Brown County, Nebraska

### Construction Materials

These tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In the first table, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Aa: Almeria-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.65 0.97
Ae: Almeria-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.97
Af: Almeria-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.97
Histosols-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
An: Anselmo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
AnC: Anselmo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
AnD: Anselmo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.57
AtF: Anselmo-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.96
Brunswick-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.09
Ba: Barney-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.95 0.97
Bd: Bolent-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
Bo: Brocksburg-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.95
BrD: Brunswick-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.06
DuB: Dunday-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
DuD: Dunday-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
Eo: Els-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.45

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
EpB: Els-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.99
Ipge-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Es: Elsmere-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
Fe: Fluvaquents-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.70 0.70
Gn: Gannett-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
IdB: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.70 0.98
IkB: Inavale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Bottom layer	0.98
In: Inavale-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.70 0.98
Barney-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.95 0.97
IpB: Ipge-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good	
IsB: Ipge-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.45
Jn: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
JnC: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
Jo: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
JoC: Jansen-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
Jr: Jansen-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Meadin-----	50	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
JtB: Jansen-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.00
Sandose-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.03 0.20
Jw: Johnstown-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.95
JwB: Johnstown-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.95
Jy: Johnstown-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.95
JyB: Johnstown-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.95
JyC: Johnstown-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.22
LcG: Labu-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sansarc-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LfB: Libory-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.62
Lo: Loup-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.08
Lp: Loup-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.08
LtB: Els-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
Loup-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.08
Ma: Marlake-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.17

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
McG: Mckelvie-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
Longpine-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
MeB: Meadin-----	100	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
MeF: Meadin-----	100	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
Oe: O'Neill-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.97
On: O'Neill-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
OsC: O'Neill-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
Meadin-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
OsD: O'Neill-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.97
Meadin-----	45	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
Pg: Pits-----	100	Not rated		Not rated	
PtB: Pivot-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.39 0.96
RtB: Ronson-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.09
Longpine-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
RtC: Ronson-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.09
Longpine-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
RtD: Ronson-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.01



CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Longpine-----	45	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00 0.00
ScB: Sandose-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Fair Thickest layer Bottom layer	0.03 0.09 0.09
SkB: Simeon-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Fair Thickest layer Bottom layer	0.33 0.91 0.91
SkD: Simeon-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Fair Thickest layer Bottom layer	0.33 0.91 0.91
SvD: Simeon-----	60	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Fair Bottom layer Thickest layer	0.91 0.98 0.98
Valentine-----	40	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good	
Tn: Tryon-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good Thickest layer	0.65
To: Tryon-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good Thickest layer	0.65
TpB: Tryon-----	55	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good Thickest layer	0.65
Els-----	45	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good Thickest layer	0.99
VaB: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good	
VaD: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good	
VaE: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good	
VaF: Valentine-----	70	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good	
Valentine-----	30	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good	
VbB: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00 0.00	Good Thickest layer	0.70

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
VbD: Valentine-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.70
VfD: Valentine-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Els-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.99
VhD: Valentine-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Libory-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.62
VpD: Valentine-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Pivot-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.70 0.96
VrD: Valentine-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.70
Sandose-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.07
VsG2: Valentine-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Simeon-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.33 0.91
VtE: Valentine-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Longpine-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
VwE: Valentine-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Tryon-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.65
VxB: Vetal-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.57
zwa: Water-----	100	Not rated		Not rated	

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
zwb: Water-----	100	Not rated		Not rated	

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.96	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
Ae: Almeria-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.99	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
Af: Almeria-----	55	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.99	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
Histosols-----	45	Fair Low content of organic matter	0.12	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone Content of organic matter	0.00 0.00
An: Anselmo-----	100	Fair Low content of organic matter	0.12	Good		Good	
AnC: Anselmo-----	100	Fair Low content of organic matter	0.12	Good		Good	
AnD: Anselmo-----	100	Fair Low content of organic matter	0.12	Good		Fair Slope	0.96
AtF: Anselmo-----	60	Fair Low content of organic matter	0.12	Fair Slope	0.02	Poor Slope	0.00
Brunswick-----	40	Fair Droughty Low content of organic matter Depth to bedrock	0.11 0.12 0.58	Poor Depth to bedrock Slope	0.00 0.32	Poor Slope Depth to bedrock	0.00 0.58
Ba: Barney-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.81	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone Hard to reclaim Rock fragments	0.00  0.00 0.97
Bd: Bolent-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.89	Fair Depth to saturated zone	0.76	Poor Too sandy  Depth to saturated zone	0.00  0.76
Bo: Brocksburg-----	100	Fair Low content of organic matter	0.12	Good		Fair Hard to reclaim	0.92

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
BrD: Brunswick-----	100	Poor Too sandy Droughty Low content of organic matter Depth to bedrock	0.00 0.03 0.12 0.58	Poor Depth to bedrock	0.00	Poor Too sandy Depth to bedrock	0.00 0.58
DuB: Dunday-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.92	Good		Poor Too sandy	0.00
DuD: Dunday-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.92	Good		Poor Too sandy	0.00
EO: Els-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.50	Fair Depth to saturated zone	0.76	Poor Too sandy Depth to saturated zone	0.00 0.76
EpB: Els-----	65	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.46	Fair Depth to saturated zone	0.76	Poor Too sandy Depth to saturated zone	0.00 0.76
Ipage-----	35	Poor Too sandy Wind erosion Low content of organic matter Droughty Too acid	0.00 0.00 0.12 0.39 0.97	Good		Poor Too sandy	0.00
Es: Elsmere-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.42	Fair Depth to saturated zone	0.76	Poor Too sandy Depth to saturated zone	0.00 0.76
Fe: Fluvaquents-----	100	Poor Too sandy	0.00	Poor Depth to saturated zone	0.00	Poor Too sandy Depth to saturated zone	0.00 0.00
Gn: Gannett-----	100	Fair Low content of organic matter Droughty	0.12 0.99	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
IdB: Inavale-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.82	Good		Poor Too sandy	0.00
IkB: Inavale-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.71	Good		Poor Too sandy	0.00
In: Inavale-----	60	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.85	Good		Poor Too sandy	0.00
Barney-----	40	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.74	Poor Depth to saturated zone	0.00	Poor Hard to reclaim  Too sandy  Depth to saturated zone Rock fragments	0.00  0.00 0.00 0.97
IpB: Ipage-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty Too acid	0.00 0.00 0.12 0.39 0.97	Good		Poor Too sandy	0.00
IsB: Ipage-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty Too acid	0.00 0.00 0.12 0.53 0.97	Good		Poor Too sandy	0.00
Jn: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion Droughty	0.12 0.97 0.99 0.99	Good		Fair Hard to reclaim	0.98
JnC: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion Droughty	0.12 0.97 0.99 0.99	Good		Fair Hard to reclaim	0.98
Jo: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion	0.12 0.97 0.99	Good		Fair Hard to reclaim	0.98
JoC: Jansen-----	100	Fair Low content of organic matter Too acid Water erosion	0.12 0.97 0.99	Good		Fair Hard to reclaim	0.98

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Jr: Jansen-----	50	Poor Too sandy Low content of organic matter Droughty Too acid Water erosion	0.00 0.12 0.73 0.97 0.99	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.03 0.98
Meadin-----	50	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.03 0.12 0.97	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.00 0.00
JtB: Jansen-----	50	Fair Low content of organic matter Too acid Water erosion	0.12 0.97 0.99	Good		Fair Hard to reclaim	0.98
Sandose-----	50	Poor Wind erosion Too sandy Low content of organic matter Water erosion	0.00 0.28 0.88 0.99	Good		Fair Too sandy	0.28
Jw: Johnstown-----	100	Fair Water erosion	0.99	Fair Shrink-swell	0.99	Fair Hard to reclaim	0.98
JwB: Johnstown-----	100	Fair Water erosion	0.99	Fair Shrink-swell	0.99	Fair Hard to reclaim	0.98
Jy: Johnstown-----	100	Fair Water erosion	0.99	Fair Shrink-swell	0.95	Fair Hard to reclaim	0.98
JyB: Johnstown-----	100	Fair Water erosion	0.99	Fair Shrink-swell	0.95	Fair Hard to reclaim	0.98
JyC: Johnstown-----	100	Fair Water erosion	0.99	Fair Shrink-swell	0.95	Fair Hard to reclaim	0.98
LcG: Labu-----	55	Poor Too clayey Droughty Depth to bedrock Low content of organic matter	0.00 0.00 0.58 0.88	Poor Depth to bedrock Slope Shrink-swell	0.00 0.32 0.83	Poor Too Clayey Slope Depth to bedrock	0.00 0.00 0.58
Sansarc-----	45	Poor Too clayey Droughty Depth to bedrock Low content of organic matter Water erosion	0.00 0.00 0.00 0.60 0.99	Poor Depth to bedrock Shrink-swell Slope	0.00 0.00 0.00	Poor Too Clayey Depth to bedrock Slope	0.00 0.00 0.00
LfB: Libory-----	100	Poor Wind erosion Low content of organic matter Water erosion	0.00 0.12 0.90	Fair Depth to saturated zone	0.14	Fair Depth to saturated zone	0.14

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lo: Loup-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.88 0.88	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
Lp: Loup-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.88 0.88	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
LtB: Els-----	50	Poor Too sandy  Wind erosion  Low content of organic matter Droughty	0.00  0.00  0.12 0.50	Fair Depth to saturated zone	0.76	Poor Too sandy  Depth to saturated zone	0.00  0.76
Loup-----	50	Poor Too sandy  Low content of organic matter Droughty	0.00  0.88 0.91	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
Ma: Marlake-----	100	Fair Too sandy  Low content of organic matter	0.41  0.88	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone Too sandy	0.00  0.41
McG: Mckelvie-----	50	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.80	Poor Slope	0.00	Poor Too sandy Slope	0.00 0.00
Longpine-----	30	Poor Droughty Depth to bedrock Low content of organic matter	0.00 0.00 0.88	Poor Depth to bedrock Slope	0.00 0.00	Poor Depth to bedrock Slope Rock fragments	0.00 0.00 0.97
MeB: Meadin-----	100	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.08 0.12 0.97	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.00 0.00
MeF: Meadin-----	100	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.08 0.12 0.97	Fair Slope	0.92	Poor Too sandy Rock fragments Hard to reclaim  Slope	0.00 0.00 0.00  0.00
Oe: O'Neill-----	100	Fair Low content of organic matter Droughty Too acid	0.12  0.48 0.84	Good		Fair Hard to reclaim	0.92



CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
On: O'Neill-----	100	Fair Low content of organic matter Droughty Too acid	0.12 0.63 0.84	Good		Fair Hard to reclaim	0.92
OsC: O'Neill-----	60	Fair Low content of organic matter Droughty Too acid	0.12 0.71 0.84	Good		Fair Hard to reclaim	0.92
Meadin-----	40	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.09 0.12 0.97	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.00 0.00
OsD: O'Neill-----	55	Fair Low content of organic matter Droughty Too acid	0.12 0.35 0.84	Good		Fair Hard to reclaim Slope	0.92 0.96
Meadin-----	45	Poor Too sandy Droughty Low content of organic matter Too acid	0.00 0.11 0.12 0.97	Good		Poor Too sandy Rock fragments Hard to reclaim Slope	0.00 0.00 0.00 0.96
Pg: Pits-----	100	Not rated		Not rated		Not rated	
PtB: Pivot-----	100	Poor Wind erosion Too sandy Low content of organic matter Droughty	0.00 0.02 0.12 0.19	Good		Fair Too sandy Hard to reclaim	0.02 0.92
RtB: Ronson-----	55	Fair Droughty Depth to bedrock Low content of organic matter	0.07 0.58 0.88	Poor Depth to bedrock	0.00	Fair Depth to bedrock	0.58
Longpine-----	45	Poor Droughty Depth to bedrock Low content of organic matter	0.00 0.00 0.88	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments	0.00 0.00
RtC: Ronson-----	55	Fair Droughty Depth to bedrock Low content of organic matter	0.20 0.58 0.88	Poor Depth to bedrock	0.00	Fair Depth to bedrock	0.58
Longpine-----	45	Poor Droughty Depth to bedrock Low content of organic matter	0.00 0.00 0.88	Poor Depth to bedrock	0.00	Poor Depth to bedrock	0.00
RtD: Ronson-----	55	Fair Droughty Depth to bedrock Low content of organic matter	0.01 0.58 0.88	Poor Depth to bedrock	0.00	Fair Depth to bedrock Slope	0.58 0.96

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Longpine-----	45	Poor Droughty Depth to bedrock Low content of organic matter	0.00 0.00 0.88	Poor Depth to bedrock	0.00	Poor Depth to bedrock Slope Rock fragments	0.00 0.96 0.97
ScB: Sandose-----	100	Poor Wind erosion Too sandy Low content of organic matter Water erosion	0.00 0.36 0.88 0.99	Good		Fair Too sandy	0.36
SkB: Simeon-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.79	Good		Poor Too sandy	0.00
SkD: Simeon-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.79	Good		Poor Too sandy	0.00
SvD: Simeon-----	60	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.74	Good		Poor Too sandy	0.00
Valentine-----	40	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.13	Good		Poor Too sandy	0.00
Tn: Tryon-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.18	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
To: Tryon-----	100	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.20	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
TpB: Tryon-----	55	Poor Too sandy  Low content of organic matter Droughty	0.00  0.12 0.18	Poor Depth to saturated zone	0.00	Poor Too sandy  Depth to saturated zone	0.00  0.00
Els-----	45	Poor Too sandy  Wind erosion  Low content of organic matter Droughty	0.00  0.00  0.12 0.46	Fair Depth to saturated zone	0.76	Poor Too sandy  Depth to saturated zone	0.00  0.76

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VaB: Valentine-----	100	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Good		Poor Too sandy	0.00
VaD: Valentine-----	100	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Good		Poor Too sandy	0.00
VaE: Valentine-----	100	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Fair Slope	0.92	Poor Too sandy Slope	0.00 0.00
VaF: Valentine-----	70	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Fair Slope	0.92	Poor Too sandy Slope	0.00 0.00
Valentine-----	30	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Poor Slope	0.00	Poor Slope Too sandy	0.00 0.00
VbB: Valentine-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.35	Good		Poor Too sandy	0.00
VbD: Valentine-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.35	Good		Poor Too sandy	0.00
VfD: Valentine-----	55	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Good		Poor Too sandy	0.00
Els-----	45	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.46	Fair Depth to saturated zone	0.76	Poor Too sandy Depth to saturated zone	0.00 0.76
VhD: Valentine-----	60	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.13	Good		Poor Too sandy	0.00

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Libory-----	40	Poor Wind erosion Low content of organic matter Water erosion	0.00 0.12 0.90	Fair Depth to saturated zone	0.14	Fair Depth to saturated zone	0.14
VpD: Valentine-----	60	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Good		Poor Too sandy	0.00
Pivot-----	40	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.18	Good		Poor Too sandy Hard to reclaim	0.00 0.92
VrD: Valentine-----	65	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.28	Good		Poor Too sandy	0.00
Sandose-----	35	Poor Wind erosion Low content of organic matter Water erosion	0.00 0.88 0.99	Good		Good	
VsG2: Valentine-----	65	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.13	Poor Slope	0.00	Poor Too sandy Slope	0.00 0.00
Simeon-----	35	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.81	Fair Slope	0.50	Poor Too sandy Slope	0.00 0.00
VtE: Valentine-----	60	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.12 0.14	Good		Poor Too sandy Slope	0.00 0.84
Longpine-----	40	Poor Droughty Depth to bedrock Low content of organic matter	0.00 0.00 0.88	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.00 0.84
VwE: Valentine-----	65	Poor Too sandy Wind erosion Droughty Low content of organic matter	0.00 0.00 0.12 0.12	Good		Poor Too sandy Slope	0.00 0.84
Tryon-----	35	Poor Too sandy Low content of organic matter Droughty	0.00 0.12 0.20	Poor Depth to saturated zone	0.00	Poor Too sandy Depth to saturated zone	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VxB: Vetal-----	100	Good		Good		Good	
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	



RECREATIONAL INTERPRETATIONS  
Brown County, Nebraska

### Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Depth to saturated zone Too sandy	1.00	Too sandy	0.95	Depth to saturated zone Too sandy	1.00
			0.95	Flooding	0.40		0.95
Ae: Almeria-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Flooding	0.40	Depth to saturated zone	1.00
Af: Almeria-----	55	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.40	Flooding	1.00
Histosols-----	45	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Content of organic matter	1.00	Content of organic matter	1.00
		Content of organic matter	1.00	Flooding	0.40	Flooding	1.00
An: Anselmo-----	100	Not limited		Not limited		Not limited	
AnC: Anselmo-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
AnD: Anselmo-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
AtF: Anselmo-----	60	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Brunswick-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Depth to bedrock	1.00 0.42
Ba: Barney-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.40	Flooding	1.00
Bd: Bolent-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Bo: Brocksburg-----	100	Not limited		Not limited		Not limited	
BrD: Brunswick-----	100	Not limited		Not limited		Very limited Slope Depth to bedrock	1.00 0.42
DuB: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy Slope	0.95 0.00
DuD: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95	Very limited Slope Too sandy	1.00 0.95
Eo: Els-----	100	Somewhat limited Too sandy Depth to saturated zone	0.84 0.07	Somewhat limited Too sandy Depth to saturated zone	0.84 0.03	Somewhat limited Too sandy Depth to saturated zone	0.84 0.07
EpB: Els-----	65	Very limited Too sandy Depth to saturated zone	1.00 0.07	Very limited Too sandy Depth to saturated zone	1.00 0.03	Very limited Too sandy Depth to saturated zone Slope	1.00 0.07 0.00
Ipage-----	35	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 0.00
Es: Elsmere-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95



RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Fe: Fluvaquents-----	100	Depth to saturated zone	0.07	Depth to saturated zone	0.03	Depth to saturated zone	0.07
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Flooding Too sandy	1.00 0.96	Too sandy Flooding	0.96 0.40	Flooding Too sandy	1.00 0.96
Gn: Gannett-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
IdB: Inavale-----	100	Very limited Flooding Too sandy	1.00 0.96	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Slope	0.96 0.00
IkB: Inavale-----	100	Very limited Flooding Too sandy	1.00 1.00	Very limited Too sandy	1.00	Very limited Too sandy Flooding Slope	1.00 0.60 0.00
In: Inavale-----	60	Very limited Flooding Too sandy	1.00 0.96	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Flooding	0.96 0.60
Barney-----	40	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
IpB: Ipage-----	100	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 0.00
IsB: Ipage-----	100	Somewhat limited Too sandy	0.84	Somewhat limited Too sandy	0.84	Somewhat limited Too sandy Slope	0.84 0.00
Jn: Jansen-----	100	Not limited		Not limited		Not limited	
JnC: Jansen-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
Jo: Jansen-----	100	Not limited		Not limited		Not limited	
JoC: Jansen-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
Jr: Jansen-----	50	Not limited		Not limited		Not limited	
Meadin-----	50	Not limited		Not limited		Somewhat limited Gravel content	0.22
JtB: Jansen-----	50	Not limited		Not limited		Not limited	
Sandose-----	50	Somewhat limited Too sandy	0.44	Somewhat limited Too sandy	0.44	Somewhat limited Too sandy Slope	0.44 0.00
Jw: Johnstown-----	100	Not limited		Not limited		Not limited	
JwB: Johnstown-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Jy: Johnstown-----	100	Not limited		Not limited		Not limited	
JyB: Johnstown-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
JyC: Johnstown-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
LcG: Labu-----	55	Very limited Slope Too clayey Restricted permeability	1.00 0.50 0.45	Very limited Slope Too clayey Restricted permeability	1.00 0.50 0.45	Very limited Slope Too clayey Restricted permeability Depth to bedrock	1.00 0.50 0.45 0.42
Sansarc-----	45	Very limited Depth to bedrock Slope Too clayey Restricted permeability	1.00 1.00 0.50 0.45	Very limited Depth to bedrock Slope Too clayey Restricted permeability	1.00 1.00 0.50 0.45	Very limited Slope Depth to bedrock Too clayey Restricted permeability	1.00 1.00 0.50 0.45

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LfB: Libory-----	100	Somewhat limited Depth to saturated zone Too sandy	0.98 0.94	Somewhat limited Too sandy Depth to saturated zone	0.94 0.75	Somewhat limited Depth to saturated zone Too sandy Slope	0.98 0.94 0.00
Lo: Loup-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Lp: Loup-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
LtB: Els-----	50	Somewhat limited Too sandy Depth to saturated zone	0.95 0.07	Somewhat limited Too sandy Depth to saturated zone	0.95 0.03	Somewhat limited Too sandy Depth to saturated zone Slope	0.95 0.07 0.00
Loup-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Ma: Marlake-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
McG: Mckelvie-----	50	Very limited Slope Too sandy	1.00 0.95	Very limited Slope Too sandy	1.00 0.95	Very limited Slope Too sandy	1.00 0.95
Longpine-----	30	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.06
MeB: Meadin-----	100	Not limited		Not limited		Somewhat limited Gravel content Slope	0.22 0.00
MeF: Meadin-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Gravel content	1.00 0.22
Oe: O'Neill-----	100	Not limited		Not limited		Not limited	
On: O'Neill-----	100	Not limited		Not limited		Not limited	
OsC: O'Neill-----	60	Not limited		Not limited		Somewhat limited Slope	0.50
Meadin-----	40	Not limited		Not limited		Somewhat limited Slope Gravel content	0.50 0.22
OsD: O'Neill-----	55	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Meadin-----	45	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope Gravel content	1.00 0.22
Pg: Pits-----	100	Not rated		Not rated		Not rated	
PtB: Pivot-----	100	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy Slope	0.79 0.00
RtB: Ronson-----	55	Not limited		Not limited		Somewhat limited Slope	0.00
Longpine-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Gravel content Slope	1.00 0.06 0.00
RtC: Ronson-----	55	Not limited		Not limited		Somewhat limited Slope Depth to bedrock	0.87 0.42
Longpine-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
RtD: Ronson-----	55	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Slope Gravel content	0.87 0.06
Longpine-----	45	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Slope Depth to bedrock Very limited Depth to bedrock Slope Gravel content	1.00 0.42 1.00 1.00 0.06
ScB: Sandose-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy Slope	0.37 0.00
SkB: Simeon-----	100	Somewhat limited Too sandy	0.72	Somewhat limited Too sandy	0.72	Somewhat limited Too sandy Slope	0.72 0.00
SkD: Simeon-----	100	Somewhat limited Too sandy	0.72	Somewhat limited Too sandy	0.72	Very limited Slope Too sandy	1.00 0.72
SvD: Simeon-----	60	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 0.00
Valentine-----	40	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 1.00
Tn: Tryon-----	100	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Too sandy	1.00 0.95
To: Tryon-----	100	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Too sandy	1.00 0.95
TpB: Tryon-----	55	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Too sandy	1.00 0.95
Els-----	45	Very limited Too sandy Depth to saturated zone	1.00 0.07	Very limited Too sandy Depth to saturated zone	1.00 0.03	Very limited Too sandy Depth to saturated zone Slope	1.00 0.07 0.00
VaB: Valentine-----	100	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 0.00
VaD: Valentine-----	100	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 1.00
VaE: Valentine-----	100	Very limited Too sandy Slope	1.00 1.00	Very limited Too sandy Slope	1.00 1.00	Very limited Slope Too sandy	1.00 1.00
VaF: Valentine-----	70	Very limited Too sandy Slope	1.00 1.00	Very limited Too sandy Slope	1.00 1.00	Very limited Slope Too sandy	1.00 1.00
Valentine-----	30	Very limited Slope Too sandy	1.00 1.00	Very limited Too sandy Slope	1.00 1.00	Very limited Slope Too sandy	1.00 1.00
VbB: Valentine-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Slope	0.96 0.00
VbD: Valentine-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Very limited Slope Too sandy	1.00 0.96
VfD: Valentine-----	55	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy Slope	1.00 1.00
Els-----	45	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VhD: Valentine-----	60	Depth to saturated zone	0.07	Depth to saturated zone	0.03	Depth to saturated zone	0.07
		Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00
		Slope		Slope		Slope	
Libory-----	40	Somewhat limited Depth to saturated zone	0.98	Somewhat limited Too sandy	0.94	Somewhat limited Depth to saturated zone	0.98
		Too sandy	0.94	Depth to saturated zone	0.75	Too sandy	0.94
						Slope	0.00
VpD: Valentine-----	60	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00
						Slope	1.00
		Somewhat limited Too sandy	0.92	Somewhat limited Too sandy	0.92	Somewhat limited Too sandy	0.92
VrD: Valentine-----	65	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Very limited Slope	1.00
						Too sandy	0.96
		Somewhat limited Too sandy	0.44	Somewhat limited Too sandy	0.44	Somewhat limited Too sandy	0.44
VsG2: Valentine-----	65	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Slope	1.00
		Slope	1.00	Slope	1.00	Too sandy	1.00
		Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Simeon-----	35	Too sandy	0.72	Too sandy	0.72	Too sandy	0.72
VtE: Valentine-----	60	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00
		Slope	0.16	Slope	0.16	Slope	1.00
		Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
Longpine-----	40	Slope	0.16	Slope	0.16	Slope	1.00
						Gravel content	0.06
VwE: Valentine-----	65	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00
		Slope	0.16	Slope	0.16	Slope	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Tryon-----	35	Too sandy	0.95	Too sandy	0.95	Too sandy	0.95
VxB: Vetal-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Depth to saturated zone Too sandy Flooding	1.00 0.95 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 1.00 0.05
Ae: Almeria-----	100	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 1.00 0.00
Af: Almeria-----	55	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
Histosols-----	45	Very limited Depth to saturated zone Content of organic matter Flooding	1.00 1.00 0.40	Very limited Flooding Content of organic matter Depth to saturated zone	1.00 1.00 1.00
An: Anselmo-----	100	Not limited		Not limited	
AnC: Anselmo-----	100	Not limited		Not limited	
AnD: Anselmo-----	100	Not limited		Somewhat limited Slope	0.04
AtF: Anselmo-----	60	Somewhat limited Slope	0.98	Very limited Slope	1.00
Brunswick-----	40	Somewhat limited Slope	0.68	Very limited Slope Depth to bedrock Droughty	1.00 0.42 0.05
Ba: Barney-----	100	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 1.00 0.02
Bd: Bolent-----	100	Not limited		Somewhat limited Flooding Droughty Depth to saturated zone	0.60 0.09 0.03
Bo: Brocksburg-----	100	Not limited		Not limited	
BrD: Brunswick-----	100	Not limited		Somewhat limited Depth to bedrock Droughty	0.42 0.23
DuB: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Droughty	0.04
DuD: Dunday-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Droughty	0.04
Eo: Els-----	100	Somewhat limited Too sandy	0.84	Somewhat limited Droughty Depth to saturated zone	0.48 0.03
EpB: Els-----	65	Very limited Too sandy	1.00	Somewhat limited Droughty Depth to saturated zone	0.54 0.03
Ipage-----	35	Very limited Too sandy	1.00	Somewhat limited Droughty	0.63
Es: Elsmere-----	100	Somewhat limited Too sandy	0.95	Somewhat limited Droughty	0.41

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
				Depth to saturated zone	0.03
Fe: Fluvaquents-----	100	Very limited Depth to saturated zone Too sandy	1.00	Very limited Flooding	1.00
			0.96	Depth to saturated zone	1.00
		Flooding	0.40	Droughty	0.00
Gn: Gannett-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
IdB: Inavale-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Droughty	0.17
IkB: Inavale-----	100	Very limited Too sandy	1.00	Somewhat limited Flooding Too sandy Droughty	0.60 0.50 0.34
In: Inavale-----	60	Somewhat limited Too sandy	0.96	Somewhat limited Flooding Droughty	0.60 0.14
Barney-----	40	Very limited Depth to saturated zone Flooding	1.00	Very limited Flooding	1.00
			0.40	Depth to saturated zone Droughty	1.00 0.07
IpB: Ipage-----	100	Very limited Too sandy	1.00	Somewhat limited Droughty	0.63
IsB: Ipage-----	100	Somewhat limited Too sandy	0.84	Somewhat limited Droughty	0.43
Jn: Jansen-----	100	Not limited		Not limited	
JnC: Jansen-----	100	Not limited		Not limited	
Jo: Jansen-----	100	Not limited		Not limited	
JoC: Jansen-----	100	Not limited		Not limited	
Jr: Jansen-----	50	Not limited		Not limited	
Meadin-----	50	Not limited		Somewhat limited Droughty	0.87
JtB: Jansen-----	50	Not limited		Not limited	
Sandose-----	50	Somewhat limited Too sandy	0.44	Somewhat limited Droughty	0.01
Jw: Johnstown-----	100	Not limited		Not limited	
JwB: Johnstown-----	100	Not limited		Not limited	
Jy: Johnstown-----	100	Not limited		Not limited	
JyB: Johnstown-----	100	Not limited		Not limited	
JyC: Johnstown-----	100	Not limited		Not limited	
LcG: Labu-----	55	Somewhat limited Slope Too clayey	0.68 0.50	Very limited Too clayey Slope Droughty Depth to bedrock	1.00 1.00 0.85 0.42
Sansarc-----	45	Very limited Slope Too clayey	1.00 0.50	Very limited Too clayey Depth to bedrock Droughty Slope	1.00 1.00 1.00 1.00
LfB: Libory-----	100	Somewhat limited Too sandy  Depth to saturated zone	0.94  0.44	Somewhat limited Depth to saturated zone	0.75

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Lo: Loup-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Droughty	1.00 0.03
Lp: Loup-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Droughty	1.00 0.03
LtB: Els-----	50	Somewhat limited Too sandy	0.95	Somewhat limited Droughty Depth to saturated zone	0.48 0.03
Loup-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Droughty	1.00 0.01
Ma: Marlake-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
McG: Mckelvie-----	50	Very limited Slope Too sandy	1.00 0.95	Very limited Slope Droughty	1.00 0.21
Longpine-----	30	Very limited Slope	1.00	Very limited Depth to bedrock Droughty Slope	1.00 1.00 1.00
MeB: Meadin-----	100	Not limited		Somewhat limited Droughty	0.73
MeF: Meadin-----	100	Somewhat limited Slope	0.08	Very limited Slope Droughty	1.00 0.73
Oe: O'Neill-----	100	Not limited		Somewhat limited Droughty	0.02
On: O'Neill-----	100	Not limited		Not limited	
OsC: O'Neill-----	60	Not limited		Not limited	
Meadin-----	40	Not limited		Somewhat limited Droughty	0.69
OsD: O'Neill-----	55	Not limited		Somewhat limited Droughty Slope	0.08 0.04
Meadin-----	45	Not limited		Somewhat limited Droughty Slope	0.63 0.04
Pg: Pits-----	100	Not rated		Not rated	
PtB: Pivot-----	100	Somewhat limited Too sandy	0.79	Somewhat limited Droughty	0.26
RtB: Ronson-----	55	Not limited		Somewhat limited Depth to bedrock Droughty	0.42 0.10
Longpine-----	45	Not limited		Very limited Depth to bedrock Droughty	1.00 1.00
RtC: Ronson-----	55	Not limited		Somewhat limited Depth to bedrock Droughty	0.42 0.00
Longpine-----	45	Not limited		Very limited Depth to bedrock Droughty	1.00 1.00
RtD: Ronson-----	55	Not limited		Somewhat limited Depth to bedrock Droughty	0.42 0.36

RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Longpine-----	45	Not limited		Slope Very limited	0.04
				Depth to bedrock	1.00
				Droughty	1.00
				Slope	0.04
ScB: Sandose-----	100	Somewhat limited Too sandy	0.37	Not limited	
SkB: Simeon-----	100	Somewhat limited Too sandy	0.72	Somewhat limited Droughty	0.22
SkD: Simeon-----	100	Somewhat limited Too sandy	0.72	Somewhat limited Droughty	0.22
SvD: Simeon-----	60	Very limited Too sandy	1.00	Somewhat limited Droughty	0.30
Valentine-----	40	Very limited Too sandy	1.00	Somewhat limited Droughty	0.87
Tn: Tryon-----	100	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Droughty	1.00 0.79
To: Tryon-----	100	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Droughty	1.00 0.76
TpB: Tryon-----	55	Very limited Depth to saturated zone Too sandy	1.00 0.95	Very limited Depth to saturated zone Droughty	1.00 0.79
Els-----	45	Very limited Too sandy	1.00	Somewhat limited Droughty Depth to saturated zone	0.54 0.03
VaB: Valentine-----	100	Very limited Too sandy	1.00	Somewhat limited Droughty	0.89
VaD: Valentine-----	100	Very limited Too sandy	1.00	Somewhat limited Droughty	0.89
VaE: Valentine-----	100	Very limited Too sandy Slope	1.00 0.08	Very limited Slope Droughty	1.00 0.89
VaF: Valentine-----	70	Very limited Too sandy Slope	1.00 0.08	Very limited Slope Droughty	1.00 0.89
Valentine-----	30	Very limited Too sandy Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.89
VbB: Valentine-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Droughty	0.52
VbD: Valentine-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Droughty	0.52
VfD: Valentine-----	55	Very limited Too sandy	1.00	Somewhat limited Droughty	0.89
Els-----	45	Very limited Too sandy	1.00	Somewhat limited Droughty Depth to saturated zone	0.54 0.03
VhD: Valentine-----	60	Very limited Too sandy	1.00	Somewhat limited Droughty	0.87
Libory-----	40	Somewhat limited Too sandy Depth to saturated zone	0.94 0.44	Somewhat limited Depth to saturated zone	0.75
VpD: Valentine-----	60	Very limited Too sandy	1.00	Somewhat limited Droughty	0.88
Pivot-----	40	Somewhat limited Too sandy	0.92	Somewhat limited Droughty	0.27



RECREATIONAL INTERPRETATIONS--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
VrD: Valentine-----	65	Somewhat limited Too sandy	0.96	Somewhat limited Droughty	0.63
Sandose-----	35	Somewhat limited Too sandy	0.44	Not limited	
VsG2: Valentine-----	65	Very limited Too sandy	1.00	Very limited Slope	1.00
		Slope	1.00	Droughty	0.87
Simeon-----	35	Somewhat limited Too sandy	0.72	Very limited Slope	1.00
		Slope	0.50	Droughty	0.19
VtE: Valentine-----	60	Very limited Too sandy	1.00	Somewhat limited Droughty	0.86
				Slope	0.16
Longpine-----	40	Not limited		Very limited Depth to bedrock	1.00
				Droughty	1.00
				Slope	0.16
VwE: Valentine-----	65	Very limited Too sandy	1.00	Somewhat limited Droughty	0.89
				Slope	0.16
Tryon-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Too sandy	0.95	Droughty	0.76
VxB: Vetal-----	100	Not limited		Not limited	
zwa: Water-----	100	Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated	



# WILDLIFE INTERPRETATIONS Brown County, Nebraska

## Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

## Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

## Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and gorden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS  
Brown County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Aa: ALMERIA-----	Poor	Fair	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
Ae: ALMERIA-----	Poor	Fair	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
Af: ALMERIA-----	Very poor	Poor	Fair	Poor	Poor	Poor	Good	Good	Poor	Poor	Good	Poor
HISTOSOLS-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Good	Poor	Good	Fair
An: ANSELMO-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
AnC: ANSELMO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
AnD: ANSELMO-----	Fair	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
AtF: ANSELMO-----	Poor	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor	Good
BRUNSWICK-----	Poor	Poor	Good	---	Good	Good	Very poor	Very poor	Fair	---	Very poor	Good
Ba: BARNEY-----	Very poor	Poor	Fair	Poor	Poor	Poor	Good	Good	Poor	Fair	Good	Fair
Bd: BOLENT-----	Poor	Fair	Good	Good	Good	Good	Fair	Very poor	Fair	Good	Poor	Good
Bo: BROCKSBURG-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
BrD: BRUNSWICK-----	Poor	Poor	Good	---	Good	Good	Very poor	Very poor	Fair	---	Very poor	Good
DuB: DUNDAY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
DuD: DUNDAY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
Eo: ELS-----	Poor	Fair	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair	Fair
EpB: ELS-----	Poor	Poor	Fair	Fair	Fair	Fair	Fair	Fair	Poor	Fair	Fair	Fair
IPAGE-----	Poor	Good	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair
Es: ELSMERE-----	Poor	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Poor	Fair	Fair	Fair
Fe: FLUVAQUENTS-----	Very poor	Very poor	Poor	Very poor	Very poor	Very poor	Good	Good	Very poor	Very poor	Good	Very poor
Gn: GANNETT-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
IdB: INAVALE-----	Fair	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Good
IkB: INAVALE-----	Very poor	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued  
Brown County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
In: INVALE-----	Very poor	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Fair
BARNEY-----	Very poor	Poor	Fair	Poor	Poor	Poor	Good	Good	Poor	Fair	Good	Fair
IpB: IPAGE-----	Poor	Good	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair
IsB: IPAGE-----	Poor	Good	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair	Fair
Jn: JANSEN-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
JnC: JANSEN-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
Jo: JANSEN-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
JoC: JANSEN-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
Jr: JANSEN-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
MEADIN-----	Poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
JtB: JANSEN-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
SANDOSE-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
Jw: JOHNSTOWN-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Fair	Very poor	Good
JwB: JOHNSTOWN-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Fair	Very poor	Good
Jy: JOHNSTOWN-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Fair	Very poor	Good
JyB: JOHNSTOWN-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Fair	Very poor	Good
JyC: JOHNSTOWN-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Fair	Very poor	Good
LcG: LABU-----	Poor	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor	Fair
SANSARC-----	Very poor	Very poor	Fair	Poor	Very poor	Fair	Very poor	Very poor	Very poor	Very poor	Very poor	Fair
LfB: LIBORY-----	Fair	Fair	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Very poor	Good
Lo: LOUP-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
Lp: LOUP-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
LtB: ELS-----	Poor	Fair	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair	Fair

WILDLIFE INTERPRETATIONS--Continued  
Brown County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
LOUP-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
Ma: MARLAKE-----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Good	Good	Very poor	Very poor	Good	Very poor
McG: MCKELVIE-----	Very poor	Very poor	Fair	Fair	Fair	Good	Very poor	Very poor	Very poor	Fair	Very poor	Poor
LONGPINE-----	Very poor	Very poor	Poor	Poor	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor	Poor
MeB: MEADIN-----	Poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
MeF: MEADIN-----	Very poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
Oe: O'NEILL-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
On: O'NEILL-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
OsC: O'NEILL-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Fair
MEADIN-----	Very poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
OsD: O'NEILL-----	Poor	Fair	Good	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Fair
MEADIN-----	Very poor	Poor	Fair	Poor	Poor	Fair	Very poor	Very poor	Poor	Poor	Very poor	Fair
Pg: PITS-----	Very poor	Very poor	Poor	Poor	Poor	Poor	Very poor	Fair	Very poor	Very poor	Poor	Poor
PtB: PIVOT-----	Fair	Fair	Fair	Good	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
RtB: RONSON-----	Fair	Fair	Good	Poor	Very poor	---	Very poor	Very poor	Fair	Very poor	Very poor	Good
LONGPINE-----	Poor	Poor	Poor	Poor	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor	Poor
RtC: RONSON-----	Fair	Fair	Good	Poor	Very poor	---	Very poor	Very poor	Fair	Very poor	Very poor	Good
LONGPINE-----	Poor	Poor	Poor	Poor	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor	Poor
RtD: RONSON-----	Poor	Fair	Good	Poor	Very poor	---	Very poor	Very poor	Poor	Very poor	Very poor	Good
LONGPINE-----	Poor	Poor	Poor	Poor	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor	Poor
ScB: SANDOSE-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
SkB: SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
SkD: SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued  
Brown County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
SvD: SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
Tn: TRYON-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
To: TRYON-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
TpB: TRYON-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
ELS-----	Poor	Poor	Fair	Fair	Fair	Fair	Fair	Fair	Poor	Fair	Fair	Fair
VaB: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VaD: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VaE: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VaF: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VALENTINE-----	Very poor	Very poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
VbB: VALENTINE-----	Fair	Good	Fair	Poor	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
VbD: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
VfD: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
ELS-----	Poor	Poor	Fair	Fair	Fair	Fair	Fair	Fair	Poor	Fair	Fair	Fair
VhD: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
LIBORY-----	Fair	Fair	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Very poor	Good
VpD: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
PIVOT-----	Fair	Fair	Fair	Good	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
VrD: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
SANDOSE-----	Fair	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
VsG2: VALENTINE-----	Very poor	Very poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair
SIMEON-----	Poor	Poor	Fair	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor	Fair



WILDLIFE INTERPRETATIONS--Continued  
Brown County, Nebraska

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
VtE: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
LONGPINE-----	Poor	Poor	Poor	Poor	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor	Poor
VvE: VALENTINE-----	Poor	Fair	Fair	Poor	Poor	Poor	Very poor	Very poor	Fair	Poor	Very poor	Fair
TRYON-----	Very poor	Poor	Fair	Poor	Poor	Fair	Good	Good	Poor	Poor	Good	Fair
VxB: VETAL-----	Good	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
zwa: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---
zwb: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---



YIELDS PER ACRE OF PASTURE AND HAYLAND  
Brown County, Nebraska

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. On animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Aa: Almeria-----	5w	---	---	---
Ae: Almeria-----	5w	---	---	---
Af: Almeria-----	6w	---	---	---
Histosols-----	8w	---	---	---
An: Anselmo-----	2e	2e	2.40	5.00
AnC: Anselmo-----	3e	3e	2.30	4.50
AnD: Anselmo-----	4e	4e	1.90	3.50
AtF: Anselmo-----	6e	---	---	---
Brunswick-----	6e	---	---	---
Ba: Barney-----	6w	---	---	---
Bd: Bolent-----	4w	4w	2.30	4.10
Bo: Brocksburg-----	2s	2s	2.00	5.50
BrD: Brunswick-----	4e	4e	1.90	3.50
DuB: Dunday-----	4e	3e	1.50	4.00
DuD: Dunday-----	4e	4e	1.10	3.30
Eo: Els-----	4w	4w	1.50	4.00
EpB: Els-----	6e	4w	---	3.40
Ipage-----	6e	4e	---	3.40
Es: Elsmere-----	4w	4w	1.60	4.00
Fe: Fluvaquents-----	8w	---	---	---
Gn: Gannett-----	5w	---	---	---
IdB: Inavale-----	4e	3e	1.50	4.00
IkB: Inavale-----	6w	---	---	---
In: Inavale-----	6w	---	---	---
Barney-----	6w	---	---	---
IpB: Ipage-----	6e	4e	---	3.90
IsB: Ipage-----	4e	4e	1.20	4.50
Jn: Jansen-----	2e	2e	1.80	5.00

YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued  
Brown County, Nebraska

PAGE 3 OF 5

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
JnC: Jansen-----	3e	3e	1.60	4.80
Jo: Jansen-----	2s	2s	1.80	5.00
JoC: Jansen-----	3e	3e	1.60	4.80
Jr: Jansen-----	2e	2e	1.30	3.50
Meadin-----	6s	4s	1.30	3.50
JtB: Jansen-----	2e	2e	1.90	5.10
Sandose-----	3e	3e	1.90	5.10
Jw: Johnstown-----	2e	2e	2.10	6.10
JwB: Johnstown-----	2e	2e	2.00	6.00
Jy: Johnstown-----	2c	1	2.50	6.30
JyB: Johnstown-----	2e	2e	2.30	6.00
JyC: Johnstown-----	3e	3e	2.10	5.60
LcG: Labu-----	6e	---	---	---
Sansarc-----	6s	---	---	---
LfB: Libory-----	3e	3e	3.40	5.20
Lo: Loup-----	5w	---	---	---
Lp: Loup-----	5w	---	---	---
LtB: Els-----	4w	4w	---	---
Loup-----	5w	---	---	---
Ma: Marlake-----	8w	---	---	---
McG: Mckelvie-----	7e	---	---	---
Longpine-----	7s	---	---	---
MeB: Meadin-----	6s	4s	---	2.00
MeF: Meadin-----	6s	---	---	---
Oe: O'Neill-----	3e	3e	1.50	4.80
On: O'Neill-----	2c	2s	1.70	5.00
OsC: O'Neill-----	4e	4e	1.20	3.60
Meadin-----	6s	---	1.20	3.60
OsD: O'Neill-----	6e	4e	---	3.30

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Meadin-----	6s	---	---	3.30
Pg: Pits-----	8s	---	---	---
PtB: Pivot-----	4e	3e	1.40	4.00
RtB: Ronson-----	3e	3e	1.50	4.00
Longpine-----	6s	---	1.50	4.00
RtC: Ronson-----	3e	3e	1.40	3.90
Longpine-----	6s	---	1.40	3.90
RtD: Ronson-----	4e	4e	---	---
Longpine-----	6s	---	---	---
ScB: Sandose-----	3e	3e	2.00	5.20
SkB: Simeon-----	6s	4s	---	2.90
SkD: Simeon-----	6s	---	---	---
SvD: Simeon-----	6s	4s	---	3.00
Valentine-----	6e	4e	---	3.00
Tn: Tryon-----	5w	---	---	---
To: Tryon-----	5w	---	---	---
TpB: Tryon-----	5w	---	---	---
Els-----	6e	4w	---	---
VaB: Valentine-----	6e	4e	---	3.20
VaD: Valentine-----	6e	4e	---	3.20
VaE: Valentine-----	6e	---	---	---
VaF: Valentine-----	6e	---	---	---
Valentine-----	7e	---	---	---
VbB: Valentine-----	4e	4e	0.80	3.60
VbD: Valentine-----	6e	4e	---	3.20
VfD: Valentine-----	6e	4e	---	3.20
Els-----	6e	4w	---	3.20
VhD: Valentine-----	6e	4e	---	3.80
Libory-----	3e	3e	---	3.80
VpD: Valentine-----	6e	4e	---	3.40

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Pivot-----	4e	3e	---	3.40
VrD: Valentine-----	6e	4e	---	3.80
Sandose-----	3e	3e	---	3.80
VsG2: Valentine-----	7e	---	---	---
Simeon-----	6s	---	---	---
VtE: Valentine-----	6e	---	---	---
Longpine-----	6s	---	---	---
VwE: Valentine-----	6e	---	---	---
Tryon-----	5w	---	---	---
VxB: Vetal-----	2e	2e	2.50	5.50
zwa: Water-----	---	---	---	---
zwb: Water-----	---	---	---	---





CONSERVATION TREE AND SHRUB MANAGEMENT  
Brown County, Nebraska

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsited indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT  
Brown County,  
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Aa: Almeria-----		Well suited	Well suited	Well suited	Well suited	High Wetness
Ae: Almeria-----		Well suited	Well suited	Well suited	Well suited	High Wetness
Af: Almeria-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	High Wetness
Histosols-----		Moderately suited Rock fragments	Poorly suited Rock fragments	Poorly suited Rock fragments	Well suited	High Wetness Soil reaction
An: Anselmo-----		Well suited	Well suited	Well suited	Well suited	Low
AnC: Anselmo-----		Well suited	Well suited	Well suited	Well suited	Low
AnD: Anselmo-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
AtF: Anselmo-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Low
Brunswick-----		Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Low
Ba: Barney-----		Well suited	Well suited	Well suited	Well suited	High Wetness
Bd: Bolent-----		Well suited	Well suited	Well suited	Well suited	Low
Bo: Brocksburg-----		Well suited	Well suited	Well suited	Well suited	Low
BrD: Brunswick-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
DuB: Dunday-----		Well suited	Well suited	Well suited	Well suited	Low
DuD: Dunday-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Eo: Els-----		Well suited	Well suited	Well suited	Well suited	Low
EpB: Els-----		Well suited	Well suited	Well suited	Well suited	Low
Ipage-----		Well suited	Well suited	Well suited	Well suited	Low
Es: Elsmere-----		Well suited	Well suited	Well suited	Well suited	Low
Fe: Fluvaquents-----		Unsuited Wetness	Poorly suited Wetness	Unsuited Wetness	Unsuited Wetness	High Wetness
Gn: Gannett-----		Well suited	Well suited	Well suited	Well suited	High Wetness
IdB: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
IkB: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
In: Inavale-----		Well suited	Well suited	Well suited	Well suited	Low
Barney-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	High Wetness
IpB: Ipage-----		Well suited	Well suited	Well suited	Well suited	Low
IsB: Ipage-----		Well suited	Well suited	Well suited	Well suited	Low
Jn: Jansen-----		Well suited	Well suited	Well suited	Well suited	Low
JnC: Jansen-----		Well suited	Well suited	Well suited	Well suited	Low
Jo: Jansen-----		Well suited	Well suited	Well suited	Well suited	Low
JoC: Jansen-----		Well suited	Well suited	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT  
Brown County,  
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Jr: Jansen----- Meadin-----		Well suited Well suited	Well suited Moderately suited Rock fragments	Well suited Well suited	Well suited Well suited	Low Low
JtB: Jansen----- Sandose-----		Well suited Well suited	Well suited Well suited	Well suited Well suited	Well suited Well suited	Low Low
Jw: Johnstown-----		Well suited	Well suited	Well suited	Well suited	Low
JwB: Johnstown-----		Well suited	Well suited	Well suited	Well suited	Low
Jy: Johnstown-----		Well suited	Well suited	Well suited	Well suited	Low
JyB: Johnstown-----		Well suited	Well suited	Well suited	Well suited	Low
JyC: Johnstown-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
LcG: Labu-----		Poorly suited Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Low
Sansarc-----		Poorly suited Stickiness	Unsuited Slope Stickiness	Poorly suited Slope Stickiness	Poorly suited Slope	Low
LfB: Libory-----		Well suited	Well suited	Well suited	Well suited	Low
Lo: Loup-----		Well suited	Well suited	Well suited	Well suited	High Wetness
Lp: Loup-----		Well suited	Well suited	Well suited	Well suited	High Wetness
LtB: Els----- Loup-----		Well suited Well suited	Well suited Well suited	Well suited Well suited	Well suited Well suited	Low High Wetness
Ma: Marlake-----		Well suited	Well suited	Well suited	Well suited	High Wetness
McG: Mckelvie-----		Moderately suited Slope	Unsuited Slope	Unsuited Slope	Unsuited Slope	Low
Longpine-----		Moderately suited Rock fragments Slope	Unsuited Slope Rock fragments	Unsuited Slope Rock fragments	Unsuited Slope	Low
MeB: Meadin-----		Well suited	Moderately suited Rock fragments	Well suited	Well suited	Low
MeF: Meadin-----		Well suited	Poorly suited Slope Rock fragments	Poorly suited Slope	Poorly suited Slope	Low
Oe: O'Neill-----		Well suited	Well suited	Well suited	Well suited	Low
On: O'Neill-----		Well suited	Well suited	Well suited	Well suited	Low
OsC: O'Neill----- Meadin-----		Well suited Well suited	Well suited Moderately suited Rock fragments	Well suited Well suited	Well suited Well suited	Low Low
OsD: O'Neill-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Meadin-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT  
Brown County,  
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Pg: Pits-----		Not rated	Rock fragments Not rated	Not rated	Not rated	Not rated
PtB: Pivot-----		Well suited	Well suited	Well suited	Well suited	Low
RtB: Ronson-----		Well suited	Well suited	Well suited	Well suited	Low
Longpine-----		Well suited	Well suited	Well suited	Well suited	Low
RtC: Ronson-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Longpine-----		Moderately suited Rock fragments	Poorly suited Rock fragments Slope	Poorly suited Rock fragments	Well suited	Low
RtD: Ronson-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Longpine-----		Moderately suited Rock fragments	Poorly suited Rock fragments Slope	Poorly suited Rock fragments	Well suited	Low
ScB: Sandose-----		Well suited	Well suited	Well suited	Well suited	Low
SKB: Simeon-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Low
SkD: Simeon-----		Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	Low
SvD: Simeon-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Low
Valentine-----		Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	Low
Tn: Tryon-----		Well suited	Well suited	Well suited	Well suited	High Wetness
To: Tryon-----		Well suited	Well suited	Well suited	Well suited	High Wetness
TpB: Tryon-----		Well suited	Well suited	Well suited	Well suited	High Wetness Low
Els-----		Well suited	Well suited	Well suited	Well suited	Low
VaB: Valentine-----		Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Low
VaD: Valentine-----		Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	Low
VaE: Valentine-----		Moderately suited Sandiness	Poorly suited Slope Sandiness	Poorly suited Slope	Poorly suited Slope	Low
VaF: Valentine-----		Moderately suited Sandiness	Poorly suited Slope Sandiness	Poorly suited Slope	Poorly suited Slope	Low
Valentine-----		Moderately suited Sandiness	Unsuited Slope	Unsuited Slope	Unsuited Slope	Low

CONSERVATION TREE AND SHRUB MANAGEMENT  
Brown County,  
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
VbB: Valentine-----		Slope Well suited	Sandiness Well suited	Well suited	Well suited	Low
VbD: Valentine-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
VfD: Valentine-----		Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	Low
Els-----		Well suited	Well suited	Well suited	Well suited	Low
VhD: Valentine-----		Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	Low
Libory-----		Well suited	Well suited	Well suited	Well suited	Low
VpD: Valentine-----		Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	Low
Pivot-----		Well suited	Well suited	Well suited	Well suited	Low
VrD: Valentine-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Sandose-----		Well suited	Well suited	Well suited	Well suited	Low
VsG2: Valentine-----		Moderately suited Sandiness	Poorly suited Slope Sandiness	Poorly suited Slope	Poorly suited Slope	Low
Simeon-----		Moderately suited Sandiness	Poorly suited Slope Sandiness	Poorly suited Slope	Poorly suited Slope	Low
VtE: Valentine-----		Moderately suited Sandiness	Moderately suited Slope Sandiness	Well suited	Well suited	Low
Longpine-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
VwE: Valentine-----		Moderately suited Sandiness	Moderately suited Slope Sandiness	Well suited	Well suited	Low
Tryon-----		Well suited	Well suited	Well suited	Well suited	High Wetness
VxB: Vetal-----		Well suited	Well suited	Well suited	Well suited	Low
zwa: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated
zwb: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated



ENGINEERING INDEX PROPERTIES  
Brown County, Nebraska

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

ENGINEERING INDEX PROPERTIES--Continued  
Brown County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
	In				Pct	Pct					Pct	
Aa: Almeria-----	0-5	Loamy fine sand	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	50-80	15-55	15-20	NP-5
	5-60	Stratified sand to fine sandy loam	SM, SP, SP- SM, SC-SM	A-2, A-3, A-4	0	0	90-100	80-100	50-80	0-50	15-20	NP-5
Ae: Almeria-----	0-5	Fine sandy loam	ML, SC, SC- SM, SM	A-4	0	0	100	100	70-100	40-70	15-25	NP-10
	5-60	Stratified sand to fine sandy loam	SC-SM, SM, SP, SP-SM	A-2, A-3, A-4	0	0	90-100	80-100	50-80	0-50	15-20	NP-5
Af: Almeria-----	0-5	Fine sandy loam	ML, SC, SC- SM, SM	A-4	0	0	100	100	70-100	40-55	15-25	NP-10
	5-60	Stratified sand to fine sandy loam	SC-SM, SM, SP, SP-SM	A-2, A-3	0	0	95-100	90-100	50-75	0-30	15-20	NP-5
Histosols-----	0-26	Mucky peat	PT	A-8	0	0	---	---	---	---	---	NP
	26-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	50-90	5-30	---	NP
An: Anselmo-----	0-7	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	7-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	18-33	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	33-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
AnC: Anselmo-----	0-7	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	7-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	18-33	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	33-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
AnD: Anselmo-----	0-7	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	7-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	18-33	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	33-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
	0-7	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	7-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	18-33	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	33-60	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
AtF: Anselmo-----	0-6	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	100	60-100	30-65	15-25	NP-7
	6-28	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	100	70-95	35-65	15-25	NP-7
	28-37	Fine sandy loam	SC-SM, SM	A-4	0	0	100	100	70-85	40-50	15-25	NP-7
	37-60	Fine sand	SC-SM, SM	A-2	0	0	100	100	50-80	15-35	15-20	NP-5
Brunswick-----	0-4	Fine sandy loam	CL, ML, SC, SM	A-4	0	0	95-100	95-100	70-95	40-55	0-25	NP-10
	4-26	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	95-100	65-95	30-65	18-35	NP-10
	26-33	Fine sandy loam	SM	A-2, A-4	0	0	95-100	95-100	65-90	20-50	0-14	NP
	33-60	Weathered bedrock			---	---	---	---	---	---	---	---
Ba: Barney-----	0-10	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	95-100	90-100	60-95	40-55	18-30	NP-7
	10-20	Stratified sand to loam	ML, SM	A-2, A-4	0	0	95-100	90-100	55-80	20-60	10-20	NP-5
	20-60	Coarse sand	SC-SM, SM, SP, SP-SM	A-1, A-2, A-3	0	0	95-100	75-100	30-80	3-15	10-15	NP-5
Bd: Bolent-----	0-5	Fine sandy loam	CL-ML, SC-SM, SM, ML	A-2, A-4	0	0	100	100	75-100	30-75	15-25	NP-10
	5-60	Stratified sand to fine sandy loam	SP-SM, SP, SC-SM, SM	A-1, A-2, A-3	0	0	95-100	90-100	40-70	3-35	10-20	NP-5
Bo: Brocksburg-----	0-16	Loam	ML, CL	A-4, A-6	0	0	100	100	90-100	70-90	25-40	3-15
	16-36	Clay loam	CL	A-6, A-7	0	0	100	100	90-100	70-80	35-45	11-20
	36-60	Coarse sand	SP-SM, SP, SM	A-1, A-2, A-3	0	0	85-95	50-90	20-60	3-15	---	NP
BrD: Brunswick-----	0-4	Fine sandy loam	CL, SC, ML, SM	A-4	0	0	95-100	95-100	70-95	40-55	0-25	NP-10
	4-17	Fine sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0	0	95-100	95-100	65-95	30-65	18-35	NP-10
	17-36	Fine sand	SM	A-2, A-4	0	0	95-100	95-100	65-90	20-50	0-14	NP
	36-60	Weathered bedrock			---	---	---	---	---	---	---	---
DuB: Dunday-----	0-18	Loamy fine sand	SM, SC-SM	A-2	0	0	100	100	90-100	13-35	15-25	NP-5
	18-60	Fine sand	SP-SM, SC-SM, SM	A-2, A-3	0	0	100	100	50-95	5-35	15-25	NP-5
DuD: Dunday-----	0-18	Loamy fine sand	SM, SC-SM	A-2	0	0	100	100	90-100	13-35	15-25	NP-5
	18-60	Fine sand	SP-SM, SM, SC-SM	A-2, A-3	0	0	100	100	50-95	5-35	15-25	NP-5



ENGINEERING INDEX PROPERTIES--Continued  
Brown County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
	In											
EO: Els-----	0-6	Loamy sand	SM	A-2	0	0	95-100	85-100	60-100	25-35	---	NP
	6-60	Fine sand	SP-SM, SP, SM	A-2, A-3	0	0	90-100	90-100	70-100	4-30	---	NP
EpB: Els-----	0-6	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	70-100	5-35	---	NP
	6-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	90-100	90-100	70-100	4-30	---	NP
Ipage-----	0-7	Fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	50-100	5-30	---	NP
	7-60	Fine sand	SP-SM, SM, SP	A-2, A-3	0	0	100	95-100	50-100	2-30	---	NP
Es: Elsmere-----	0-14	Loamy fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	70-100	5-35	10-20	NP-5
	14-60	Fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	60-100	5-30	5-15	NP-5
Fe: Fluvaquents----	0-60	Stratified loamy fine sand	SM, SP-SM	A-2, A-3, A-4	0	0	100	100	50-70	5-40	15-25	NP-5
Gn: Gannett-----	0-6	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	100	95-100	30-50	15-25	NP-5
	6-23	Loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0	100	100	95-100	30-65	15-35	NP-15
	23-60	Fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	90-100	5-15	---	NP
IdB: Inavale-----	0-8	Loamy fine sand	SP-SM, SC-SM, SM	A-2, A-3	0	0	100	100	85-95	5-35	15-25	NP-5
	8-60	Stratified fine sand to loam	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
IkB: Inavale-----	0-8	Sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	65-85	5-30	15-25	NP-5
	8-60	Stratified coarse sand to loam	SM, SP-SM, SC-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
In: Inavale-----	0-10	Loamy fine sand	SM, SP-SM, SC-SM	A-3, A-2	0	0	100	100	85-95	5-35	15-25	NP-5
	10-60	Stratified sand to loam	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	100	70-90	5-30	15-25	NP-5
Barney-----	0-3	Fine sandy loam	ML, CL-ML, SC-SM, SM	A-4	0	0	95-100	90-100	60-95	40-55	18-30	NP-7
	7-60	Sand	SP-SM, SC-SM, SM, SP	A-1, A-2, A-3	0	0	95-100	75-100	30-80	3-15	10-15	NP-5
	10-20	Stratified sand to loam	ML, SM	A-2, A-4	0	0	95-100	90-100	55-80	20-60	10-20	NP-5
IpB: Ipage-----	0-7	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	50-100	5-30	---	NP
	7-60	Fine sand	SP, SM, SP-SM	A-2, A-3	0	0	100	95-100	50-100	2-30	---	NP
IsB: Ipage-----	0-7	Loamy sand	SM, SP-SM	A-2	0	0	100	100	50-90	10-35	---	NP
	7-60	Sand	SM, SP, SP-SM	A-2, A-3	0	0	100	95-100	50-100	2-30	---	NP
Jn: Jansen-----	0-13	Fine sandy loam	SM, ML	A-4	0	0	95-100	90-100	80-100	40-55	20-30	NP-6
	13-30	Clay loam	CL	A-6, A-7	0	0	95-100	90-100	80-100	50-80	30-45	10-25
	30-60	Sand	SW, SW-SM, SP-SM, SP	A-1, A-2, A-3	0	0	85-100	45-100	35-65	3-10	0-10	NP-5
JnC: Jansen-----	0-13	Fine sandy loam	SM, ML	A-4	0	0	95-100	90-100	80-100	40-55	20-30	NP-6
	13-30	Clay loam	CL	A-6, A-7	0	0	95-100	90-100	80-100	50-80	30-45	10-25
	30-60	Coarse sand	SP, SW, SW- SM, SP-SM	A-1, A-2, A-3	0	0	85-100	45-100	35-65	3-10	0-10	NP-5
Jo: Jansen-----	0-13	Loam	ML, CL	A-4, A-6	0	0	100	100	90-100	50-95	25-40	3-15
	13-30	Loam	CL	A-6, A-7	0	0	95-100	90-100	80-100	50-80	30-45	10-25
	30-60	Sand	SW-SM, SP, SW, SP-SM	A-1, A-2, A-3	0	0	85-100	45-100	35-65	3-10	0-10	NP-5
JoC: Jansen-----	0-13	Loam	ML, CL	A-4, A-6	0	0	100	100	90-100	50-95	25-40	3-15
	13-30	Clay loam	CL	A-6, A-7	0	0	95-100	90-100	80-100	50-80	30-45	10-25
	30-60	Gravelly coarse sand	SW-SM, SW, SP-SM, SP	A-1, A-2, A-3	0	0	85-100	45-100	35-65	3-10	0-10	NP-5
Jr: Jansen-----	0-7	Fine sandy loam	SM, ML	A-4	0	0	95-100	90-100	80-100	40-55	20-30	NP-6
	7-22	Clay loam	CL	A-6, A-7	0	0	95-100	90-100	80-100	50-80	30-45	10-25
	22-60	Sand	SW, SW-SM, SP-SM, SP	A-1, A-2, A-3	0	0	85-100	45-100	35-65	3-10	0-10	NP-5
Meadin-----	0-5	Sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	85-100	75-95	45-80	25-55	15-20	NP-5
	5-10	Loamy sand	GM, GP-GM, SM, SP-SM	A-2, A-3	0	0	50-90	50-90	50-65	5-35	---	NP
	10-60	Gravelly coarse sand	GP, GP-GM, SP, SP-SM	A-1	0	0	40-80	30-70	15-50	1-10	---	NP

ENGINEERING INDEX PROPERTIES--Continued  
Brown County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
JtB: Jansen-----	0-11 11-32 32-60	Fine sandy loam Clay loam Coarse sand	SM, ML CL SW-SM, SP, SP-SM, SW	A-4 A-6, A-7 A-1, A-2, A-3	0 0 0	0 0 0	95-100 95-100 85-100	90-100 90-100 45-100	80-100 80-100 35-65	40-55 50-80 3-10	20-30 30-45 0-10	NP-6 10-25 NP-5
Sandose-----	0-10 10-38 38-60	Loamy fine sand Loamy fine sand Loam	SM, SC-SM SP-SM, SM, SC-SM SC, SC-SM, CL-ML, CL	A-2 A-2, A-3 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	70-100 65-85 80-100	15-35 5-35 45-80	15-20 15-20 20-45	NP-5 NP-5 5-20
Jw: Johnstown-----	0-21 21-44 44-50 50-60	Fine sandy loam Clay loam Silty clay loam Gravelly coarse sand	CL-ML, SC-SM, ML, SM CL CL, CL-ML SM, SP-SM	A-4 A-6, A-7 A-4, A-6 A-1, A-2, A-3	0 0 0 0	0 0 0 0	100 100 100 60-100	100 100 100 50-95	70-85 90-100 85-100 25-70	40-55 80-95 50-95 5-15	15-20 30-50 20-40 5-15	NP-5 15-30 5-20 NP-5
JwB: Johnstown-----	0-21 21-44 44-50 50-60	Fine sandy loam Clay loam Silty clay loam Sand	SM, CL-ML, SC-SM, ML CL CL-ML, CL SM, SP-SM	A-4 A-6, A-7 A-4, A-6 A-1, A-2, A-3	0 0 0 0	0 0 0 0	100 100 100 60-100	100 100 100 50-95	70-85 90-100 85-100 25-70	40-55 80-95 50-95 5-15	15-20 30-50 20-40 5-15	NP-5 15-30 5-20 NP-5
Jy: Johnstown-----	0-21 21-44 44-50 50-60	Loam Clay loam Silty clay loam Coarse sand	ML, CL-ML, CL CL CL, CL-ML SM, SP-SM	A-4, A-6 A-6, A-7 A-4, A-6 A-1, A-2, A-3	0 0 0 0	0 0 0 0	100 100 100 60-100	100 100 100 50-95	85-100 90-100 85-100 25-70	70-100 80-95 50-95 5-15	20-40 30-50 20-40 5-15	3-18 15-30 5-20 NP-5
JyB: Johnstown-----	0-21 21-44 44-50 50-60	Loam Clay loam Silty clay loam Sand	CL, CL-ML, ML CL CL, CL-ML SM, SP-SM	A-6, A-4 A-6, A-7 A-4, A-6 A-1, A-2, A-3	0 0 0 0	0 0 0 0	100 100 100 60-100	100 100 100 50-95	85-100 90-100 85-100 25-70	70-100 80-95 50-95 5-15	20-40 30-50 20-40 5-15	3-18 15-30 5-20 NP-5
JyC: Johnstown-----	0-21 21-44 44-50 50-60	Loam Clay loam Silty clay loam Loamy sand	CL, CL-ML, ML CL CL, CL-ML SM, SP-SM	A-4, A-6 A-6, A-7 A-4, A-6 A-1, A-2, A-3	0 0 0 0	0 0 0 0	100 100 100 60-100	100 100 100 50-95	85-100 90-100 85-100 25-70	70-100 80-95 50-95 5-15	20-40 30-50 20-40 5-15	3-18 15-30 5-20 NP-5
LcG: Labu-----	0-5 5-23 23-60	Silty clay Silty clay Weathered bedrock	CH, MH CH, MH CH, MH	A-7 A-7 A-7	0 0 ---	0 0 ---	100 95-100 95-100	95-100 95-100 95-100	90-100 90-100 90-100	85-100 85-100 85-100	50-90 50-90 50-90	20-55 20-55 ---
Sansarc-----	0-3 3-12 12-60	Silty clay Clay Weathered bedrock	CH, MH CH, MH CH, MH	A-7 A-7 A-7	0 0 0	0 0 0	100 95-100 100	95-100 95-100 95-100	90-100 95-100 90-100	75-100 85-100 85-100	60-90 60-90 50-90	25-55 25-55 20-55
LfB: Libory-----	0-14 14-25 25-60	Loamy fine sand Loamy fine sand Loam	SM SP-SM, SM CL-ML, CL	A-2, A-4 A-2 A-4, A-6	0 0 0	0 0 0	100 100 100	100 100 100	65-85 55-80 85-100	15-45 12-35 60-95	10-20 10-20 20-40	NP-5 NP-5 4-24
Lo: Loup-----	0-10 10-60	Fine sandy loam Fine sand	SM, SC-SM SP-SM, SM	A-2, A-4 A-2, A-3	0 0	0 0	100 95-100	100 95-100	70-95 65-100	20-50 5-20	15-25 10-20	NP-6 NP-5
Lp: Loup-----	0-10 10-60	Fine sandy loam Fine sand	SM, SC-SM SP-SM, SM	A-2, A-4 A-2, A-3	0 0	0 0	100 100	100 100	70-95 65-100	20-50 5-20	15-25 10-20	NP-6 NP-5
LtB: Loup-----	0-11 11-60	Fine sandy loam Sand	SM, SC-SM SM, SP-SM	A-2, A-4 A-2, A-3	0 0	0 0	100 95-100	100 95-100	70-95 65-100	20-50 5-20	15-25 10-20	NP-6 NP-5
Els-----	0-6 6-60	Loamy fine sand Fine sand	SM SP-SM, SM, SP	A-2 A-2, A-3	0 0	0 0	95-100 90-100	85-100 90-100	60-100 70-100	25-35 4-30	---	NP NP
Ma: Marlake-----	0-6 6-60	Fine sandy loam Loamy fine sand	SM, ML SP-SM, SM	A-4 A-2, A-3, A-4	0 0	0 0	100 100	100 100	70-95 50-85	40-55 5-50	15-35 ---	NP-10 NP
McG: Mckelvie-----	0-6 6-60	Loamy fine sand Fine sand	SM, SC-SM SM, SP, SP- SM, SC-SM	A-2, A-4 A-2, A-3	0 0	0 0	100 85-100	95-100 80-100	65-100 50-95	15-40 3-35	15-20 15-20	NP-5 NP-5
Longpine-----	0-6 6-10 10-60	Fine sandy loam Gravelly fine sandy loam Unweathered bedrock	ML, SM SM SM	A-4 A-1, A-2	0 0 ---	0 0-5 ---	95-100 55-100	75-100 50-75	75-100 40-60	40-65 10-35	15-25 15-25	NP-7 NP-5 ---
MeB: Meadin-----	0-7 7-12 12-60	Sandy loam Very gravelly loamy sand, loamy sand Gravelly coarse sand	CL-ML, SC-SM, SM, ML GM, GP-GM, SM, SP-SM GP, GP-GM, SP, SP-SM	A-2, A-4 A-2, A-3 A-1	0 0 0	0 0 0	85-100 50-90 40-80	75-95 50-90 30-70	45-80 50-65 15-50	25-55 5-35 1-10	15-20 ---	NP-5 NP NP

ENGINEERING INDEX PROPERTIES--Continued  
Brown County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
MeF: Meadin-----	0-7	Sandy loam	SM, CL-ML, ML, SC-SM	A-2, A-4	0	0	85-100	75-95	45-80	25-55	15-20	NP-5
	7-12	Sandy loam	SP-SM, SM, GP-GM, GM	A-2, A-3	0	0	50-90	50-90	50-65	5-35	---	NP
	12-60	Gravelly coarse sand	SP, SP-SM, GP, GP-GM	A-1	0	0	40-80	30-70	15-50	1-10	---	NP
Oe: O'Neill-----	0-8	Fine sandy loam	SC-SM, SM, CL, ML	A-4	0	0	95-100	95-100	70-85	35-55	15-25	NP-10
	8-26	Fine sandy loam	SC-SM, SC	A-2, A-4	0	0	95-100	95-100	60-75	30-50	15-30	NP-10
	26-60	Gravelly coarse sand	SP, SP-SM	A-1, A-2, A-3	0	0	70-100	50-90	25-60	0-5	10-20	NP-5
On: O'Neill-----	0-8	Loam	CL, CL-ML, ML	A-4, A-6	0	0	95-100	95-100	85-95	60-70	20-35	3-12
	8-26	Fine sandy loam	SC-SM, SC	A-2, A-4	0	0	95-100	95-100	60-75	30-50	15-30	NP-10
	26-60	Coarse sand	SP, SP-SM	A-1, A-2, A-3	0	0	70-100	50-90	25-60	0-5	10-20	NP-5
OsC: O'Neill-----	0-8	Sandy loam	CL, ML, SM, SC-SM	A-4	0	0	95-100	95-100	70-85	35-55	15-25	NP-10
	8-30	Fine sandy loam	SC, SC-SM	A-2, A-4	0	0	95-100	95-100	60-75	30-50	15-30	NP-10
	30-60	Sand	SP-SM, SP	A-1, A-2, A-3	0	0	70-100	50-90	25-60	0-5	10-20	NP-5
	0-8	Sandy loam	ML, SC-SM, SM, CL-ML	A-2, A-4	0	0	85-100	75-95	45-80	25-55	15-20	NP-5
	8-12	Loamy sand	GM, GP-GM, SM, SP-SM	A-2, A-3	0	0	50-90	50-90	50-65	5-35	---	NP
	12-60	Gravelly coarse sand	GP, GP-GM, SP, SP-SM	A-1	0	0	40-80	30-70	15-50	1-10	---	NP
OsD: O'Neill-----	0-6	Sandy loam	SM, SC-SM, CL, ML	A-4	0	0	95-100	95-100	70-85	35-55	15-25	NP-10
	6-24	Fine sandy loam	SC, SC-SM	A-2, A-4	0	0	95-100	95-100	60-75	30-50	15-30	NP-10
	24-60	Gravelly coarse sand	SP, SP-SM	A-1, A-2, A-3	0	0	70-100	50-90	25-60	0-5	10-20	NP-5
	0-9	Sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	85-100	75-95	45-80	25-55	15-20	NP-5
	9-12	Loamy sand	GP-GM, SM, SP-SM, GM	A-2, A-3	0	0	50-90	50-90	50-65	5-35	---	NP
	12-60	Gravelly coarse sand	GP-GM, SP, SP-SM, GP	A-1	0	0	40-80	30-70	15-50	1-10	---	NP
Pg: Pits-----	0-60	Gravelly sand	SM, SP, SP- SM, GP-GM	A-1, A-2, A-3	---	0-5	45-100	40-100	0-80	0-40	0-14	NP
PtB: Pivot-----	0-16	Loamy sand	SP-SM, SM	A-2	0	0	100	100	50-90	10-35	15-20	NP-5
	16-27	Loamy sand	SM, SP-SM, SW-SM	A-2, A-3	0	0	100	95-100	50-85	5-30	10-20	NP-5
	27-60	Coarse sand	GP, SP, SP- SM, GP-GM	A-1, A-2, A-3	0	0	50-100	50-90	20-54	2-12	10-20	NP-5
RtB: Ronson-----	0-13	Fine sandy loam	SM, SC-SM	A-2, A-4	0	0	100	95-100	60-95	25-45	15-30	NP-7
	13-27	Sandy loam	SC-SM, SM, SC	A-2, A-4	0	0	100	100	60-90	25-45	15-30	NP-10
	27-60	Weathered bedrock			---	---	---	---	---	---	---	---
	0-6	Fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	40-65	15-25	NP-7
	6-17	Gravelly fine sandy loam	SM	A-1, A-2	0	0-5	55-100	50-75	40-60	10-35	15-25	NP-5
	17-60	Unweathered bedrock			---	---	---	---	---	---	---	---
RtC: Ronson-----	0-13	Fine sandy loam	SM, SC-SM	A-2, A-4	0	0	100	95-100	60-95	25-45	15-30	NP-7
	13-37	Sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	100	60-90	25-45	15-30	NP-10
	37-60	Weathered bedrock			---	---	---	---	---	---	---	---
	0-6	Fine sandy loam	SM, ML	A-4	0	0	95-100	75-100	75-100	40-65	15-25	NP-7
	6-9	Gravelly fine sandy loam	SM	A-1, A-2	0	0-5	55-100	50-75	40-60	10-35	15-25	NP-5
	9-60	Unweathered bedrock			---	---	---	---	---	---	---	---
RtD: Ronson-----	0-12	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	60-95	25-45	15-30	NP-7
	12-24	Fine sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	100	100	60-90	25-45	15-30	NP-10
	24-60	Weathered bedrock			---	---	---	---	---	---	---	---
	0-6	Fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	40-65	15-25	NP-7
	6-11	Gravelly fine sandy loam	SM	A-1, A-2	0	0-5	55-100	50-75	40-60	10-35	15-25	NP-5
	11-60	Unweathered bedrock			---	---	---	---	---	---	---	---
ScB: Sandose-----	0-15	Loamy fine sand	SC-SM, SM	A-2	0	0	100	100	70-100	15-35	15-20	NP-5
	15-26	Loamy fine sand	SM, SP-SM, SC-SM	A-2, A-3	0	0	100	100	65-85	5-35	15-20	NP-5
	26-46	Loam	CL-ML, SC, SC-SM, CL	A-4, A-6, A-7	0	0	100	100	80-100	45-80	20-45	5-20
	46-60	Very fine sandy loam	CL, ML, SC, SM	A-4	0	0	100	95-100	70-95	40-75	15-25	NP-10

ENGINEERING INDEX PROPERTIES--Continued  
Brown County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
	In											
SkB: Simeon-----	0-5	Loamy sand	SM, SP-SM	A-2, A-3	0	0	95-100	90-100	51-95	5-35	10-20	NP-5
	5-60	Coarse sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
SkD: Simeon-----	0-5	Loamy sand	SM, SP-SM	A-2, A-3	0	0	95-100	90-100	51-95	5-35	10-20	NP-5
	5-60	Coarse sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
SvD: Simeon-----	0-5	Fine sand	SM, SP-SM	A-2, A-3	0	0	95-100	90-100	51-95	5-20	10-20	NP-5
	5-60	Coarse sand	SP, SM, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
Valentine-----	0-6	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	6-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Tn: Tryon-----	0-5	Loamy fine sand	SM, SP-SM	A-2	0	0	100	100	85-100	10-30	10-20	NP-5
	5-60	Fine sand	SM, SP-SM, SP	A-2, A-3	0	0	100	100	51-95	3-30	5-15	NP-5
To: Tryon-----	0-6	Loamy fine sand	SM, SP-SM	A-2	0	0	100	100	85-100	10-30	10-20	NP-5
	6-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	51-90	5-30	5-15	NP-5
TpB: Tryon-----	0-5	Loamy fine sand	SM, SP-SM	A-2	0	0	100	100	85-100	10-30	10-20	NP-5
	5-60	Fine sand	SP-SM, SM, SP	A-2, A-3	0	0	100	100	51-95	3-30	5-15	NP-5
Els-----	0-6	Fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	70-100	5-35	---	NP
	6-60	Fine sand	SM, SP-SM, SP	A-2, A-3	0	0	90-100	90-100	70-100	4-30	---	NP
VaB: Valentine-----	0-4	Fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VaD: Valentine-----	0-4	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VaE: Valentine-----	0-4	Fine sand	SM, SP-SM, SP	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SP-SM, SM, SP	A-3, A-2	0	0	100	100	70-100	2-25	15-20	NP-5
VaF: Valentine-----	0-4	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SP-SM, SM, SP	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Valentine-----	0-4	Fine sand	SP-SM, SM, SP	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SP, SP-SM, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VbB: Valentine-----	0-7	Loamy fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	95-100	2-35	15-20	NP-5
	7-13	Loamy fine sand	SP, SM, SP-SM	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	13-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VbD: Valentine-----	0-7	Loamy fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	95-100	2-35	15-20	NP-5
	7-13	Loamy fine sand	SP-SM, SM, SP	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	13-60	Fine sand	SP-SM, SM, SP	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
VfD: Valentine-----	0-4	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Els-----	0-6	Fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	70-100	5-35	---	NP
	6-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	90-100	90-100	70-100	4-30	---	NP
VhD: Valentine-----	0-6	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	6-60	Fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Libory-----	0-10	Loamy fine sand	SM	A-2, A-4	0	0	100	100	65-85	15-45	10-20	NP-5
	10-24	Loamy fine sand	SP-SM, SM	A-2	0	0	100	100	55-80	12-35	10-20	NP-5
	24-60	Silty clay loam	CL-ML, CL	A-4, A-6	0	0	100	100	85-100	60-95	20-40	4-24
VpD: Valentine-----	0-5	Fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	5-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Pivot-----	0-10	Loamy fine sand	SP-SM, SM	A-2	0	0	100	100	50-90	10-35	15-20	NP-5
	10-28	Loamy fine sand	SP-SM, SW-SM, SM	A-2, A-3	0	0	100	95-100	50-85	5-30	10-20	NP-5
	28-60	Coarse sand	GP-GM, SP, SP-SM, GP	A-1, A-2, A-3	0	0	50-100	50-90	20-54	2-12	10-20	NP-5
VrD: Valentine-----	0-7	Loamy fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	95-100	2-35	15-20	NP-5
	7-10	Loamy fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	90-100	2-35	15-20	NP-5
	10-60	Fine sand	SP, SM, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Sandose-----	0-10	Loamy fine sand	SM, SC-SM	A-2	0	0	100	100	70-100	15-35	15-20	NP-5
	10-20	Loamy fine sand	SM, SP-SM, SC-SM	A-2, A-3	0	0	100	100	65-85	5-35	15-20	NP-5
	20-40	Sandy clay loam	CL, CL-ML, SC, SC-SM	A-4, A-6, A-7	0	0	100	100	80-100	45-80	20-45	5-20
	40-60	Very fine sandy loam	SM, SC, CL, ML	A-4	0	0	100	95-100	70-95	40-75	15-25	NP-10
VsG2: Valentine-----	0-6	Fine sand	SP-SM, SP, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	6-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Simeon-----	0-7	Loamy sand	SP-SM, SM	A-2, A-3	0	0	95-100	90-100	51-95	5-35	10-20	NP-5
	7-60	Sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	35-95	0-30	10-20	NP-5
VtE: Valentine-----	0-7	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	7-60	Fine sand	SP, SP-SM, SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Longpine-----	0-6	Fine sandy loam	SM, ML	A-4	0	0	95-100	75-100	75-100	40-65	15-25	NP-7
	6-13	Gravelly fine sandy loam	SM	A-1, A-2	0	0-5	55-100	50-75	40-60	10-35	15-25	NP-5
	13-60	Unweathered bedrock			---	---	---	---	---	---	---	---

ENGINEERING INDEX PROPERTIES--Continued  
Brown County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
VwE:	In											
Valentine-----	0-4	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
	4-60	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	70-100	2-25	15-20	NP-5
Tryon-----	0-6	Loamy fine sand	SM, SP-SM	A-2	0	0	100	100	85-100	10-30	10-20	NP-5
	6-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	51-90	5-30	5-15	NP-5
VxB:												
Vetal-----	0-16	Loam	CL, ML, CL-ML	A-4, A-6	0	0	100	95-100	90-100	50-65	20-35	NP-12
	16-31	Loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0	0	100	95-100	60-100	30-65	20-30	NP-10
	31-56	Loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	90-100	60-100	30-65	20-30	NP-10
	56-60	Loamy fine sand	SM, SC-SM	A-2	0	0	100	90-100	85-100	15-35	15-25	NP-5
zwa:												
Water-----	---	---	---	---	---	---	---	---	---	---	---	---
zwb:												
Water-----	---	---	---	---	---	---	---	---	---	---	---	---



## Physical Properties

This table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Saturated hydraulic conductivity refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity (K-sat). The estimates in the table indicate the rate of water movement, in micrometers per second (um/sec), when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In this table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in this table as the K factor (Kw and Kf) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor Kw indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor Kf indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.

PHYSICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska: Published

PAGE 2 OF 7

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					

2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.

3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.

4L. Calcareous loams, silt loams, clay loams, and silty clay loams.

4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.

5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.

6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.

7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.

8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.



PHYSICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
Aa:														
Almeria-----	0-5	87	7	3-10	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	0.5-4.0	.17	.17	5	8	0
	5-60	92	6	1-10	1.55-1.80	5.95-19.98	0.05-0.12	0.0-2.9	0.0-0.5	.15	.15			
Ae:														
Almeria-----	0-5	64	26	7-15	1.30-1.50	2.00-6.00	0.13-0.18	0.0-2.9	1.0-12	.24	.24	5	8	0
	5-60	92	6	1-10	1.55-1.80	5.95-19.98	0.05-0.12	0.0-2.9	0.0-0.5	.15	.15			
Af:														
Almeria-----	0-5	64	26	7-15	1.30-1.50	1.98-5.95	0.16-0.20	0.0-2.9	1.0-12	.24	.24	5	8	0
	5-60	92	6	1-10	1.55-1.80	5.95-19.98	0.05-0.12	0.0-2.9	0.0-0.5	.15	.15			
Histosols----	0-26			---	0.15-0.45	2.00-6.00	0.35-0.45	0.0-2.9	20-99	---	---	3	8	0
	26-60	97	1	0-5	1.60-1.80	5.95-19.98	0.06-0.07	0.0-2.9	0.0-0.5	.15	.15			
An:														
Anselmo-----	0-7	66	20	10-18	1.30-1.60	2.00-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	7-18	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	18-33	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	33-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
AnC:														
Anselmo-----	0-7	66	20	10-18	1.30-1.60	2.00-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	7-18	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	18-33	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	33-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
AnD:														
Anselmo-----	0-7	66	20	10-18	1.30-1.60	2.00-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	7-18	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	18-33	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	33-60	86	7	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.17	.17			
AtF:														
Anselmo-----	0-6	66	20	10-18	1.30-1.60	2.00-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	6-28	66	20	10-18	1.40-1.60	2.00-6.00	0.15-0.19	0.0-2.9	0.5-1.0	.20	.20			
	28-37	66	20	10-18	1.40-1.60	2.00-6.00	0.12-0.16	0.0-2.9	0.5-1.0	.24	.24			
	37-60	91	1	5-10	1.50-1.75	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Brunswick----	0-4	67	20	8-18	1.50-1.60	1.98-5.95	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	3	3	86
	4-26	67	20	8-18	1.40-1.60	1.98-5.95	0.09-0.14	0.0-2.9	0.0-0.5	.20	.20			
	26-33	66	27	4-10	1.40-1.70	1.98-19.98	0.05-0.11	0.0-2.9	0.0-0.5	.20	.20			
	33-60			---	---	0.20-0.60	---	---	---	---	---			
Ba:														
Barney-----	0-10	64	26	7-15	1.50-1.70	1.98-5.95	0.10-0.18	0.0-2.9	1.0-3.0	.20	.20	5	8	0
	10-20	92	6	2-10	1.60-1.80	1.98-19.98	0.09-0.14	0.0-2.9	0.0-0.5	.15	.15			
	20-60	91	6	0-5	1.70-1.90	5.95-19.98	0.04-0.07	0.0-2.9	0.0-0.5	.10	.10			
Bd:														
Bolent-----	0-5	66	20	7-20	1.30-1.50	2.00-6.00	0.13-0.18	0.0-2.9	1.0-2.0	.24	.24	5	3	86
	5-60	95	1	1-10	1.50-1.80	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Bo:														
Brocksburg---	0-16	44	41	10-20	1.30-1.50	0.60-2.00	0.20-0.24	0.0-2.9	1.0-3.0	.28	.28	4	5	56
	16-36	36	39	20-30	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	1.0-2.0	.28	.28			
	36-60	91	6	0-5	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
BrD:														
Brunswick----	0-4	67	20	8-18	1.50-1.60	1.98-5.95	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	3	3	86
	4-17	67	20	8-18	1.40-1.60	1.98-5.95	0.09-0.14	0.0-2.9	0.0-0.5	.20	.20			
	17-36	92	1	4-10	1.40-1.70	1.98-19.98	0.05-0.11	0.0-2.9	0.0-0.5	.15	.15			
	36-60			---	---	0.20-0.60	---	---	---	---	---			
DuB:														
Dunday-----	0-18	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	5	2	134
	18-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
DuD:														
Dunday-----	0-18	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	5	2	134
	18-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Eo:														
Els-----	0-6	84	9	3-10	1.55-1.60	5.95-19.98	0.08-0.13	0.0-2.9	0.5-3.0	.17	.17	5	2	134
	6-60	95	1	0-8	1.50-1.60	5.95-19.98	0.05-0.08	0.0-2.9	0.0-0.5	.15	.15			
EpB:														
Els-----	0-6	94	1	2-8	1.60-1.70	5.95-19.98	0.07-0.12	0.0-2.9	0.5-3.0	.15	.15	5	1	220
	6-60	95	1	0-8	1.50-1.60	5.95-19.98	0.05-0.08	0.0-2.9	0.0-0.5	.15	.15			
Ipaga-----	0-7	96	1	1-5	1.40-1.50	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	7-60	95	1	1-8	1.50-1.60	5.95-19.98	0.04-0.10	0.0-2.9	0.0-0.5	.15	.15			
Es:														
Elsmere-----	0-14	87	7	3-10	1.55-1.70	5.95-19.98	0.10-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	14-60	97	1	0-5	1.50-1.60	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Fe:														
Fluvaquents--	0-60	87	7	2-10	1.30-1.80	5.95-19.98	0.07-0.13	0.0-2.9	2.0-8.0	.17	.17	5	8	0
Gn:														
Gannett-----	0-6	62	26	7-18	1.20-1.50	2.00-6.00	0.13-0.15	0.0-2.9	4.0-12	.20	.20	4	8	0
	6-23	43	40	7-27	1.20-1.50	0.60-6.00	0.13-0.19	0.0-2.9	0.5-1.0	.20	.20			
	23-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
IdB:														
Inavale-----	0-8	87	7	2-10	1.50-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	8-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
IkB:														
Inavale-----	0-8	96	2	1-5	1.50-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	8-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.10	.10			

PHYSICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
In:	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Inavale-----	0-10	87	7	2-10	1.50-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
Barney-----	10-60	92	1	3-10	1.50-1.60	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
	0-3	64	26	7-15	1.50-1.70	1.98-5.95	0.10-0.18	0.0-2.9	1.0-3.0	.20	.20	5	8	0
	7-60	91	6	0-5	1.70-1.90	5.95-19.98	0.04-0.07	0.0-2.9	0.0-0.5	.15	.15			
	10-20	92	6	2-10	1.60-1.80	1.98-19.98	0.09-0.14	0.0-2.9	0.0-0.5	.15	.15			
IpB:														
Ipape-----	0-7	96	1	1-5	1.40-1.50	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	7-60	95	1	1-8	1.50-1.60	5.95-19.98	0.04-0.10	0.0-2.9	0.0-0.5	.15	.15			
IsB:														
Ipape-----	0-7	84	9	3-10	1.40-1.50	5.95-19.98	0.10-0.12	0.0-2.9	0.5-2.0	.17	.17	5	2	134
	7-60	95	1	1-8	1.50-1.60	5.95-19.98	0.04-0.10	0.0-2.9	0.0-0.5	.15	.15			
Jn:														
Jansen-----	0-13	67	20	8-18	1.30-1.40	2.00-6.00	0.13-0.18	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	13-30	36	39	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	30-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
JnC:														
Jansen-----	0-13	67	20	8-18	1.30-1.40	2.00-6.00	0.13-0.18	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	13-30	38	36	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	30-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
Jo:														
Jansen-----	0-13	43	40	12-22	1.20-1.30	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.28	.28	4	5	56
	13-30	38	36	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	30-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
JoC:														
Jansen-----	0-13	43	40	12-22	1.20-1.30	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.28	.28	4	5	56
	13-30	38	36	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	30-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
Jr:														
Jansen-----	0-7	67	20	8-18	1.30-1.40	2.00-6.00	0.13-0.18	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	7-22	38	36	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	22-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
Meadin-----	0-5	68	24	7-12	1.50-1.60	2.00-6.00	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	5-10	80	16	3-18	1.50-1.60	2.00-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	10-60	90	6	2-5	1.50-1.70	19.98-19.98	0.02-0.05	0.0-2.9	0.0-0.5	.05	.10			
JtB:														
Jansen-----	0-11	67	20	8-18	1.30-1.40	2.00-6.00	0.13-0.18	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	11-32	36	39	18-32	1.30-1.50	0.20-2.00	0.15-0.19	3.0-5.9	0.5-2.0	.37	.37			
	32-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
Sandose-----	0-10	79	17	0-8	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	10-38	79	17	0-8	1.55-1.80	5.95-19.98	0.06-0.11	0.0-2.9	0.5-1.0	.17	.17			
	38-60	56	18	18-35	1.30-1.65	0.20-2.00	0.15-0.19	3.0-5.9	0.5-1.0	.37	.37			
Jw:														
Johnstown----	0-21	67	20	7-20	1.50-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	21-44	27	42	27-35	1.40-1.50	0.20-2.00	0.15-0.20	3.0-5.9	0.5-2.0	.37	.37			
	44-50	7	65	15-32	1.30-1.50	0.20-2.00	0.17-0.22	3.0-5.9	0.0-0.5	.43	.43			
	50-60	91	6	0-5	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
JwB:														
Johnstown----	0-21	67	20	5-20	1.50-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	21-44	27	42	27-35	1.40-1.50	0.20-2.00	0.15-0.20	3.0-5.9	0.5-2.0	.37	.37			
	44-50	7	65	15-32	1.30-1.50	0.20-2.00	0.17-0.22	3.0-5.9	0.0-0.5	.43	.43			
	50-60	91	6	0-5	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
Jy:														
Johnstown----	0-21	41	42	12-22	1.30-1.50	0.60-2.00	0.20-0.24	3.0-5.9	1.0-3.0	.28	.28	4	5	56
	21-44	27	42	27-35	1.40-1.50	0.20-2.00	0.15-0.20	3.0-5.9	0.5-2.0	.37	.37			
	44-50	7	65	15-32	1.30-1.50	0.20-2.00	0.17-0.22	3.0-5.9	0.0-0.5	.43	.43			
	50-60	91	6	0-5	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
JyB:														
Johnstown----	0-21	41	42	12-22	1.30-1.50	0.60-2.00	0.20-0.24	3.0-5.9	1.0-3.0	.28	.28	4	5	56
	21-44	27	42	27-35	1.40-1.50	0.20-2.00	0.15-0.20	3.0-5.9	0.5-2.0	.37	.37			
	44-50	7	65	15-32	1.30-1.50	0.20-2.00	0.17-0.22	3.0-5.9	0.0-0.5	.43	.43			
	50-60	91	6	0-5	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
JyC:														
Johnstown----	0-21	41	42	12-22	1.30-1.50	0.60-2.00	0.20-0.24	3.0-5.9	1.0-3.0	.28	.28	4	5	56
	21-44	27	42	27-35	1.40-1.50	0.20-2.00	0.15-0.20	3.0-5.9	0.5-2.0	.37	.37			
	44-50	7	65	15-32	1.30-1.50	0.20-2.00	0.17-0.22	3.0-5.9	0.0-0.5	.43	.43			
	50-60	80	16	0-8	1.50-1.70	5.95-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.17	.17			
LcG:														
Labu-----	0-5	3	45	45-60	1.20-1.30	0.00-0.06	0.08-0.14	6.0-8.9	1.0-3.0	.32	.32	3	4	86
	5-23	3	45	45-60	1.20-1.30	0.00-0.06	0.08-0.14	6.0-8.9	0.5-1.0	.32	.32			
	23-60					0.06-0.20								
Sansarc-----	0-3	5	4	55-65	1.10-1.20	0.00-0.06	0.08-0.12	9.0-25.0	1.0-2.0	.37	.37	2	4	86
	3-12	12	28	55-65	1.10-1.20	0.00-0.06	0.06-0.12	9.0-25.0	0.1-1.0	.37	.37			
	12-60					0.06-0.20			0.0-0.5					
LfB:														
Libory-----	0-14	86	7	2-12	1.50-1.70	5.95-19.98	0.07-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	14-25	86	7	2-12	1.50-1.70	5.95-19.98	0.06-0.11	0.0-2.9	0.5-1.0	.17	.17			
	25-60	20	57	15-32	1.20-1.40	0.20-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
Lo:														
Loup-----	0-10	64	26	7-15	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	4.0-8.0	.20	.20	3	8	0
	10-60	95	1	2-7	1.50-1.70	5.95-19.98	0.06-0.08	0.0-2.9	0.5-1.0	.15	.15			
Lp:														
Loup-----	0-10	64	26	7-15	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	4.0-8.0	.20	.20	3	8	0
	10-60	95	1	2-7	1.50-1.70	5.95-19.98	0.06-0.08	0.0-2.9	0.5-1.0	.15	.15			

PHYSICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
LtB:														
Els-----	0-6	87	7	3-10	1.55-1.60	5.95-19.98	0.08-0.13	0.0-2.9	0.5-3.0	.17	.17	5	2	134
	6-60	95	1	0-8	1.50-1.60	5.95-19.98	0.05-0.08	0.0-2.9	0.0-0.5	.15	.15			
Loup-----	0-11	64	26	7-15	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	4.0-8.0	.20	.20	3	8	0
	11-60	95	1	2-7	1.50-1.70	5.95-19.98	0.06-0.08	0.0-2.9	0.5-1.0	.15	.15			
Ma:														
Marlake-----	0-6	66	20	10-18	1.40-1.50	1.98-5.95	0.16-0.18	0.0-2.9	4.0-8.0	.20	.20	5	8	0
	6-60	78	16	3-8	1.50-1.60	5.95-19.98	0.06-0.11	0.0-2.9	0.5-1.0	.17	.17			
McG:														
Mckelvie-----	0-6	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	6-60	95	1	1-10	1.60-1.80	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Longpine-----	0-6	65	27	7-12	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	2	3	86
	6-10	65	27	7-12	1.50-1.70	2.00-6.00	0.09-0.11	0.0-2.9	0.5-1.0	.15	.28			
	10-60			---	---	0.20-0.60	---	---	---	---	---			
MeB:														
Meadin-----	0-7	68	24	7-12	1.50-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	7-12	84	4	5-18	1.50-1.60	5.95-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	12-60	90	6	2-5	1.50-1.70	19.98-19.98	0.02-0.05	0.0-2.9	0.0-0.5	.05	.10			
MeF:														
Meadin-----	0-7	68	24	7-12	1.50-1.60	2.00-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	7-12	65	23	5-18	1.50-1.60	1.98-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.20	.20			
	12-60	90	6	2-5	1.50-1.70	19.98-19.98	0.02-0.05	0.0-2.9	0.0-0.5	.05	.10			
Oe:														
O'Neill-----	0-8	64	27	7-12	1.60-1.80	1.98-5.95	0.10-0.15	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	8-26	67	20	7-18	1.60-1.80	2.00-6.00	0.10-0.15	0.0-2.9	0.5-1.0	.20	.20			
	26-60	92	7	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
On:														
O'Neill-----	0-8	43	38	10-27	1.40-1.60	0.60-2.00	0.14-0.17	0.0-2.9	1.0-3.0	.28	.28	4	5	56
	8-26	67	20	7-18	1.60-1.80	2.00-6.00	0.10-0.15	0.0-2.9	0.5-1.0	.20	.20			
	26-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
OsC:														
O'Neill-----	0-8	67	24	7-12	1.60-1.80	1.98-5.95	0.10-0.15	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	8-30	68	20	7-18	1.60-1.80	2.00-6.00	0.10-0.15	0.0-2.9	0.5-1.0	.20	.20			
	30-60	97	2	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.15	.15			
Meadin-----	0-8	67	24	7-12	1.50-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	8-12	84	4	5-18	1.50-1.60	5.95-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	12-60	90	6	2-5	1.50-1.70	19.98-19.98	0.02-0.05	0.0-2.9	0.0-0.5	.05	.10			
OsD:														
O'Neill-----	0-6	67	24	7-12	1.60-1.80	1.98-5.95	0.10-0.15	0.0-2.9	1.0-3.0	.20	.20	4	3	86
	6-24	68	20	7-18	1.60-1.80	2.00-6.00	0.10-0.15	0.0-2.9	0.5-1.0	.20	.20			
	24-60	92	7	0-3	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.05	.10			
Meadin-----	0-9	67	24	7-12	1.50-1.60	1.98-5.95	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	9-12	84	4	5-18	1.50-1.60	5.95-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	12-60	90	6	2-5	1.50-1.70	19.98-19.98	0.02-0.05	0.0-2.9	0.0-0.5	.05	.10			
Pg:														
Pits-----	0-60	95	1	0-8	1.70-2.00	5.95-19.98	0.02-0.09	0.0-2.9	0.0-0.5	.15	.15	2	8	0
PtB:														
Pivot-----	0-16	84	9	3-12	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	5	2	134
	16-27	85	9	2-10	1.40-1.70	5.95-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	27-60	92	4	1-8	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
RtB:														
Ronson-----	0-13	66	20	10-18	1.30-1.40	2.00-6.00	0.11-0.17	0.0-2.9	2.0-4.0	.20	.20	3	3	86
	13-27	67	19	10-18	1.35-1.45	2.00-6.00	0.09-0.15	0.0-2.9	0.5-1.0	.24	.24			
	27-60			---	---	0.20-6.00	---	---	0.0-0.5	---	---			
Longpine-----	0-6	65	27	7-12	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	2	3	86
	6-17	65	27	5-12	1.50-1.70	2.00-6.00	0.09-0.11	0.0-2.9	0.5-1.0	.15	.28			
	17-60			---	---	0.20-0.60	---	---	---	---	---			
RtC:														
Ronson-----	0-13	66	20	10-18	1.30-1.40	2.00-6.00	0.11-0.17	0.0-2.9	2.0-4.0	.20	.20	3	3	86
	13-37	67	19	10-18	1.35-1.45	2.00-6.00	0.09-0.15	0.0-2.9	0.5-1.0	.24	.24			
	37-60			---	---	0.20-6.00	---	---	0.0-0.5	---	---			
Longpine-----	0-6	64	27	7-12	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	2	3	86
	6-9	65	27	5-12	1.50-1.70	2.00-6.00	0.09-0.11	0.0-2.9	0.5-1.0	.15	.28			
	9-60			---	---	0.20-0.60	---	---	---	---	---			
RtD:														
Ronson-----	0-12	66	20	10-18	1.30-1.40	2.00-6.00	0.11-0.17	0.0-2.9	2.0-4.0	.20	.20	3	3	86
	12-24	67	19	10-18	1.35-1.45	2.00-6.00	0.09-0.15	0.0-2.9	0.5-1.0	.24	.24			
	24-60			---	---	0.20-6.00	---	---	0.0-0.5	---	---			
Longpine-----	0-6	64	27	7-12	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	2	3	86
	6-11	65	27	5-12	1.50-1.70	2.00-6.00	0.09-0.11	0.0-2.9	0.5-1.0	.15	.28			
	11-60			---	---	0.20-0.60	---	---	---	---	---			
ScB:														
Sandose-----	0-15	79	16	2-8	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	15-26	79	16	2-8	1.55-1.80	5.95-19.98	0.06-0.11	0.0-2.9	0.5-1.0	.17	.17			
	26-46	56	18	18-35	1.30-1.65	0.20-2.00	0.15-0.19	3.0-5.9	0.5-1.0	.37	.37			
	46-60	67	20	5-20	1.45-1.70	0.57-5.95	0.12-0.19	0.0-2.9	0.0-0.5	.28	.28			
SkB:														
Simeon-----	0-5	82	9	5-12	1.30-1.50	5.95-19.98	0.08-0.14	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	5-60	90	4	2-10	1.50-1.70	5.95-99.90	0.05-0.10	0.0-2.9	0.0-0.5	.10	.10			
SkD:														
Simeon-----	0-5	82	9	5-12	1.30-1.50	5.95-19.98	0.08-0.14	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	5-60	90	4	2-10	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.10	.10			

PHYSICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
SvD:														
Simeon-----	0-5	92	1	3-10	1.30-1.50	5.95-19.98	0.06-0.12	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	5-60	90	4	2-10	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.10	.10			
Valentine----	0-6	95	1	2-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	6-60	96	1	2-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Tn:														
Tryon-----	0-5	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	4.0-8.0	.17	.17	5	8	0
	5-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
To:														
Tryon-----	0-6	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	4.0-8.0	.17	.17	5	8	0
	6-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
TpB:														
Tryon-----	0-5	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	4.0-8.0	.17	.17	5	8	0
	5-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Els-----	0-6	94	1	2-8	1.60-1.70	5.95-19.98	0.07-0.12	0.0-2.9	0.5-3.0	.15	.15	5	1	220
	6-60	95	1	0-8	1.50-1.60	5.95-19.98	0.05-0.08	0.0-2.9	0.0-0.5	.15	.15			
VaB:														
Valentine----	0-4	95	1	2-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	2-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VaD:														
Valentine----	0-4	95	1	2-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	2-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VaE:														
Valentine----	0-4	95	1	2-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	2-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VaF:														
Valentine----	0-4	95	1	2-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	2-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
	0-4	95	1	2-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	2-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VbB:														
Valentine----	0-7	87	7	2-10	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	7-13	87	7	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.17	.17			
	13-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VbD:														
Valentine----	0-7	87	7	2-10	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	7-13	87	7	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.17	.17			
	13-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VfD:														
Valentine----	0-4	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Els-----	0-6	94	1	2-8	1.60-1.70	5.95-19.98	0.07-0.12	0.0-2.9	0.5-3.0	.15	.15	5	1	220
	6-60	95	1	0-8	1.50-1.60	5.95-19.98	0.05-0.08	0.0-2.9	0.0-0.5	.15	.15			
VhD:														
Valentine----	0-6	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	6-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Libory-----	0-10	86	7	2-12	1.50-1.70	5.95-19.98	0.07-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	10-24	86	7	2-12	1.50-1.70	5.95-19.98	0.06-0.11	0.0-2.9	0.5-1.0	.17	.17			
	24-60	19	52	15-32	1.20-1.40	0.20-1.98	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
VpD:														
Valentine----	0-5	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	5-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Pivot-----	0-10	86	7	3-12	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	1.0-2.0	.17	.17	5	2	134
	10-28	87	7	2-10	1.40-1.70	5.95-19.98	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	28-60	92	4	1-8	1.50-1.70	19.98-19.98	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
VrD:														
Valentine----	0-7	87	7	2-10	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	7-10	87	7	2-10	1.55-1.75	5.95-19.98	0.09-0.11	0.0-2.9	0.0-0.5	.17	.17			
	10-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Sandose-----	0-10	79	17	0-8	1.35-1.55	5.95-19.98	0.10-0.12	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	10-20	79	17	0-8	1.55-1.80	5.95-19.98	0.06-0.11	0.0-2.9	0.5-1.0	.17	.17			
	20-40	38	36	18-35	1.30-1.65	0.20-2.00	0.15-0.19	3.0-5.9	0.5-1.0	.37	.37			
	40-60	63	24	5-20	1.45-1.70	0.57-5.95	0.12-0.19	0.0-2.9	0.0-0.5	.28	.28			
VsG2:														
Valentine----	0-6	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	6-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Simeon-----	0-7	82	9	5-12	1.30-1.50	5.95-19.98	0.08-0.14	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	7-60	90	4	2-10	1.50-1.70	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
VtE:														
Valentine----	0-7	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	7-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Longpine-----	0-6	65	27	5-12	1.30-1.50	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	2	3	86
	6-13	65	27	5-12	1.50-1.70	2.00-6.00	0.09-0.11	0.0-2.9	0.5-1.0	.17	.17			
	13-60			---	---	0.20-0.60	---	---	---	---	---			
VwE:														
Valentine----	0-4	96	1	0-6	1.40-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-60	96	1	0-6	1.60-1.80	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Tryon-----	0-6	87	7	3-10	1.40-1.60	5.95-19.98	0.10-0.12	0.0-2.9	4.0-8.0	.17	.17	5	8	0
	6-60	95	1	2-7	1.50-1.70	5.95-19.98	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
VxB:														
Vetal-----	0-16	43	39	10-26	1.20-1.30	2.00-6.00	0.17-0.21	0.0-2.9	1.0-3.0	.28	.28	5	5	56
	16-31	42	38	12-26	1.25-1.40	2.00-6.00	0.11-0.17	0.0-2.9	0.5-2.0	.28	.28			
	31-56	43	39	10-26	1.30-1.40	2.00-6.00	0.11-0.17	0.0-2.9	0.5-2.0	.24	.24			
	56-60	86	7	5-10	1.40-1.50	5.95-19.98	0.08-0.10	0.0-2.9	0.0-1.0	.17	.17			

PHYSICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
zwa: Water-----	---			---	---	---	---	---	---	---	---	-	---	---
zwb: Water-----	---			---	---	---	---	---	---	---	---	-	---	---



CHEMICAL PROPERTIES OF THE SOILS  
Brown County, Nebraska

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Brown County, Nebraska

PAGE 2 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Aa:							
Almeria-----	0-5	3.0-15	6.1-8.4	0-5	0	0.0-4.0	0
	5-60	1.0-8.0	5.6-7.3	0	0	0.0-4.0	0
Ae:							
Almeria-----	0-5	4.0-22	6.1-8.4	0-5	0	0.0-4.0	0
	5-60	1.0-8.0	5.6-7.3	0	0	0.0-4.0	0
Af:							
Almeria-----	0-5	4.0-22	6.1-8.4	0-5	0	0.0-4.0	0
	5-60	1.0-8.0	5.6-7.3	0-5	0	0.0-4.0	0
Histosols-----	0-26	---	5.6-7.3	---	---	0.0-2.0	---
	26-60	0.0-5.0	5.6-7.3	0	0	0.0-2.0	---
An:							
Anselmo-----	0-7	10-20	5.6-7.3	0	0	0	0
	7-18	5.0-15	6.1-7.8	0	0	0	0
	18-33	5.0-15	6.6-7.8	0-3	0	0	0
	33-60	0.0-10	6.6-7.8	0-3	0	0	0
AnC:							
Anselmo-----	0-7	10-20	5.6-7.3	0	0	0	0
	7-18	5.0-15	6.1-7.8	0	0	0	0
	18-33	5.0-15	6.6-7.8	0-3	0	0	0
	33-60	0.0-10	6.6-7.8	0-3	0	0	0
AnD:							
Anselmo-----	0-7	10-20	5.6-7.3	0	0	0	0
	7-18	5.0-15	6.1-7.8	0	0	0	0
	18-33	5.0-15	6.6-7.8	0-3	0	0	0
	33-60	0.0-10	6.6-7.8	0-3	0	0	0
AtF:							
Anselmo-----	0-6	10-20	5.6-7.3	0	0	0	0
	6-28	5.0-15	6.1-7.8	0	0	0	0
	28-37	5.0-15	6.6-7.8	0-3	0	0	0
	37-60	0.0-10	6.6-7.8	0-3	0	0	0
Brunswick-----	0-4	7.0-15	5.6-7.3	0	0	0	0
	4-26	6.0-13	5.6-7.3	0	0	0	0
	26-33	3.0-7.0	5.6-7.8	0	0	0	0
	33-60	---	---	---	---	---	---
Ba:							
Barney-----	0-10	3.0-14	6.6-8.4	0-5	0	0	0
	10-20	2.0-7.0	6.6-8.4	0-5	0	0	0
	20-60	0.0-4.0	6.6-7.8	0	0	0	0
Bd:							
Bolent-----	0-5	5.0-20	7.4-8.4	1-10	0	0	0
	5-60	1.0-10	6.6-8.4	1-10	0	0	0
Bo:							
Brocksburg-----	0-16	10-20	6.1-7.3	0	0	0	0
	16-36	15-25	6.6-7.8	0	0	0	0
	36-60	0.0-5.0	6.6-7.8	0	0	0	0
BrD:							
Brunswick-----	0-4	7.0-15	5.6-7.3	0	0	0	0
	4-17	6.0-13	5.6-7.3	0	0	0	0
	17-36	3.0-7.0	5.6-7.8	0	0	0	0
	36-60	---	---	---	---	---	---
DuB:							
Dunday-----	0-18	5.0-10	6.1-7.3	0	0	0	0
	18-60	1.0-5.0	6.1-7.8	0	0	0	0
DuD:							
Dunday-----	0-18	5.0-10	6.1-7.3	0	0	0	0
	18-60	1.0-5.0	6.1-7.8	0	0	0	0
Eo:							
Els-----	0-6	5.0-10	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	6.1-7.3	0	0	0	0
EpB:							
Els-----	0-6	0.0-5.0	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	6.1-7.3	0	0	0	0
Ipage-----	0-7	0.0-5.0	5.1-7.3	0	0	0	0
	7-60	0.0-5.0	5.1-7.3	0	0	0	0
Es:							
Elsmere-----	0-14	5.0-15	5.6-7.3	0	0	0	0
	14-60	0.0-5.0	5.6-7.8	0	0	0	0
Fe:							
Fluvaquents-----	0-60	1.0-20	6.6-8.4	0-5	0	0.0-2.0	0



CHEMICAL PROPERTIES OF THE SOILS--Continued  
Brown County, Nebraska

PAGE 3 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Gn:							
Gannett-----	0-6	8.0-25	6.6-8.4	0-10	0	0	0
	6-23	2.0-15	6.6-8.4	0-10	0	0	0
	23-60	1.0-10	6.6-8.4	0-10	0	0	0
IdB:							
Inavale-----	0-8	2.0-10	5.6-7.8	0	0	0	0
	8-60	1.0-10	5.6-7.8	0	0	0	0
IkB:							
Inavale-----	0-8	1.0-5.0	5.6-7.8	0	0	0	0
	8-60	1.0-10	5.6-7.8	0	0	0	0
In:							
Inavale-----	0-10	2.0-10	5.6-7.8	0	0	0	0
	10-60	1.0-10	5.6-7.8	0	0	0	0
Barney-----	0-3	3.0-14	6.6-8.4	0-5	0	0	0
	7-60	0.0-4.0	6.6-7.8	0	0	0	0
	10-20	2.0-7.0	6.6-8.4	0-5	0	0	0
IpB:							
Ipave-----	0-7	0.0-5.0	5.1-7.3	0	0	0	0
	7-60	0.0-5.0	5.1-7.3	0	0	0	0
IsB:							
Ipave-----	0-7	5.0-10	5.1-7.3	0	0	0	0
	7-60	0.0-5.0	5.1-7.3	0	0	0	0
Jn:							
Jansen-----	0-13	5.0-15	5.1-7.3	0	0	0	0
	13-30	12-25	5.1-7.3	0	0	0	0
	30-60	0.0-2.0	5.1-7.3	0	0	0	0
JnC:							
Jansen-----	0-13	5.0-15	5.1-7.3	0	0	0	0
	13-30	12-25	5.1-7.3	0	0	0	0
	30-60	0.0-2.0	5.1-7.3	0	0	0	0
Jo:							
Jansen-----	0-13	10-20	5.1-7.3	0	0	0	0
	13-30	12-25	5.1-7.3	0	0	0	0
	30-60	0.0-2.0	5.1-7.3	0	0	0	0
JoC:							
Jansen-----	0-13	10-20	5.1-7.3	0	0	0	0
	13-30	12-25	5.1-7.3	0	0	0	0
	30-60	0.0-2.0	5.1-7.3	0	0	0	0
Jr:							
Jansen-----	0-7	5.0-15	5.1-7.3	0	0	0	0
	7-22	12-25	5.1-7.3	0	0	0	0
	22-60	0.0-2.0	5.1-7.3	0	0	0	0
Meadin-----	0-5	10-20	5.1-7.3	0	0	0	0
	5-10	5.0-15	5.1-7.3	0	0	0	0
	10-60	0.0-5.0	6.1-7.3	0	0	0	0
JtB:							
Jansen-----	0-11	5.0-15	5.1-7.3	0	0	0	0
	11-32	12-25	5.1-7.3	0	0	0	0
	32-60	0.0-2.0	5.1-7.3	0	0	0	0
Sandose-----	0-10	1.0-9.0	5.6-7.3	0	0	0	0
	10-38	1.0-8.0	6.1-7.3	0	0	0	0
	38-60	13-27	6.6-7.8	0-5	0	0	0
Jw:							
Johnstown-----	0-21	5.0-20	5.6-7.3	0	0	0	0
	21-44	20-30	6.1-7.8	0	0	0	0
	44-50	10-25	6.6-8.4	0	0	0	0
	50-60	0.0-5.0	6.6-7.8	0-5	0	0	0
JwB:							
Johnstown-----	0-21	5.0-20	5.6-7.3	0	0	0	0
	21-44	20-30	6.1-7.8	0	0	0	0
	44-50	10-25	6.6-8.4	0	0	0	0
	50-60	0.0-5.0	6.6-7.8	0-5	0	0	0
Jy:							
Johnstown-----	0-21	15-30	5.6-7.3	0	0	0	0
	21-44	20-30	6.1-7.8	0	0	0	0
	44-50	10-25	6.6-8.4	0	0	0	0
	50-60	0.0-5.0	6.6-7.8	0-5	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Brown County, Nebraska

PAGE 4 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
JyB:							
Johnstown-----	0-21	15-30	5.6-7.3	0	0	0	0
	21-44	20-30	6.1-7.8	0	0	0	0
	44-50	10-25	6.6-8.4	0	0	0	0
	50-60	0.0-5.0	6.6-7.8	0-5	0	0	0
JyC:							
Johnstown-----	0-21	15-30	5.6-7.3	0	0	0	0
	21-44	20-30	6.1-7.8	0	0	0	0
	44-50	10-25	6.6-8.4	0	0	0	0
	50-60	0.0-5.0	6.6-7.8	0-5	0	0	0
LcG:							
Labu-----	0-5	30-45	6.6-8.4	5-15	0	0	0
	5-23	30-45	7.4-8.4	5-15	0	0	0
	23-60	---	---	---	---	---	---
Sansarc-----	0-3	40-45	6.6-8.4	0-5	0	0.0-2.0	0-1
	3-12	35-45	7.4-8.4	1-10	0-2	0.0-2.0	0-1
	12-60	35-45	7.4-8.4	1-10	1-5	0.0-2.0	0-1
LfB:							
Libory-----	0-14	2.0-12	5.6-7.3	0	0	0	0
	14-25	1.0-9.0	5.6-7.3	0	0	0	0
	25-60	10-22	5.6-7.8	0	0	0	0
Lo:							
Loup-----	0-10	5.0-20	5.6-7.8	0	0	0	0
	10-60	0.0-5.0	5.6-7.3	0	0	0	0
Lp:							
Loup-----	0-10	5.0-20	5.6-7.8	0	0	0	0
	10-60	0.0-5.0	5.6-7.3	0	0	0	0
LtB:							
Loup-----	0-11	5.0-20	5.6-7.8	0	0	0	0
	11-60	0.0-5.0	5.6-7.3	0	0	0	0
Els-----	0-6	5.0-10	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	6.1-7.3	0	0	0	0
Ma:							
Marlake-----	0-6	10-20	6.1-8.4	0-5	0	0	0
	6-60	0.0-10	6.1-7.8	0	0	0	0
McG:							
Mckelvie-----	0-6	5.0-10	6.1-7.8	0	0	0	0
	6-60	1.0-8.0	6.6-8.4	0-5	0	0	0
Longpine-----	0-6	5.0-10	6.6-8.4	0-5	0	0	0
	6-10	5.0-10	7.4-8.4	2-15	0	0	0
	10-60	---	---	---	---	---	---
MeB:							
Meadin-----	0-7	10-20	5.1-7.3	0	0	0	0
	7-12	5.0-15	5.1-7.3	0	0	0	0
	12-60	0.0-5.0	6.1-7.3	0	0	0	0
MeF:							
Meadin-----	0-7	10-20	5.1-7.3	0	0	0	0
	7-12	5.0-15	5.1-7.3	0	0	0	0
	12-60	0.0-5.0	6.1-7.3	0	0	0	0
Oe:							
O'Neill-----	0-8	10-20	5.1-6.5	0	0	0	0
	8-26	5.0-15	5.6-7.3	0	0	0	0
	26-60	0.0-5.0	5.6-7.3	0	0	0	0
On:							
O'Neill-----	0-8	10-20	5.1-6.5	0	0	0	0
	8-26	5.0-15	5.6-7.3	0	0	0	0
	26-60	0.0-5.0	5.6-7.3	0	0	0	0
OsC:							
O'Neill-----	0-8	10-20	5.1-6.5	0	0	0	0
	8-30	5.0-15	5.6-7.3	0	0	0	0
	30-60	0.0-5.0	5.6-7.3	0	0	0	0
Meadin-----	0-8	10-20	5.1-7.3	0	0	0	0
	8-12	5.0-15	5.1-7.3	0	0	0	0
	12-60	0.0-5.0	6.1-7.3	0	0	0	0
OsD:							
O'Neill-----	0-6	10-20	5.1-6.5	0	0	0	0
	6-24	5.0-15	5.6-7.3	0	0	0	0
	24-60	0.0-5.0	5.6-7.3	0	0	0	0
Meadin-----	0-9	10-20	5.1-7.3	0	0	0	0
	9-12	5.0-15	5.1-7.3	0	0	0	0
	12-60	0.0-5.0	6.1-7.3	0	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Brown County, Nebraska

PAGE 5 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Pg:							
Pits-----	0-60	0.0-5.0	6.6-8.4	0	0	0	0
PtB:							
Pivot-----	0-16	3.0-10	5.6-7.3	0	0	0	0
	16-27	1.0-8.0	5.6-7.3	0	0	0	0
	27-60	1.0-6.0	5.6-7.3	0	0	0	0
RtB:							
Ronson-----	0-13	10-20	6.1-8.4	0-5	0	0.0-2.0	0
	13-27	5.0-15	7.4-8.4	5-10	0	0.0-2.0	0
	27-60	---	---	---	---	---	---
Longpine-----	0-6	5.0-10	6.6-8.4	0-5	0	0	0
	6-17	5.0-10	7.4-8.4	2-15	0	0	0
	17-60	---	---	---	---	---	---
RtC:							
Ronson-----	0-13	10-20	6.1-8.4	0-5	0	0.0-2.0	0
	13-37	5.0-15	7.4-8.4	5-10	0	0.0-2.0	0
	37-60	---	---	---	---	---	---
Longpine-----	0-6	5.0-10	6.6-8.4	0-5	0	0	0
	6-9	5.0-10	7.4-8.4	2-15	0	0	0
	9-60	---	---	---	---	---	---
RtD:							
Ronson-----	0-12	10-20	6.1-8.4	0-5	0	0.0-2.0	0
	12-24	5.0-15	7.4-8.4	5-10	0	0.0-2.0	0
	24-60	---	---	---	---	---	---
Longpine-----	0-6	5.0-10	6.6-8.4	0-5	0	0	0
	6-11	5.0-10	7.4-8.4	2-15	0	0	0
	11-60	---	---	---	---	---	---
ScB:							
Sandose-----	0-15	1.0-9.0	5.6-7.3	0	0	0	0
	15-26	1.0-8.0	6.1-7.3	0	0	0	0
	26-46	13-27	6.6-7.8	0-5	0	0	0
	46-60	3.0-15	6.6-8.4	0-10	0	0	0
SkB:							
Simeon-----	0-5	0.0-5.0	6.1-7.8	0	0	0	0
	5-60	0.0-5.0	6.1-7.8	0	0	0	0
SkD:							
Simeon-----	0-5	0.0-5.0	6.1-7.8	0	0	0	0
	5-60	0.0-5.0	6.1-7.8	0	0	0	0
SvD:							
Simeon-----	0-5	0.0-5.0	6.1-7.8	0	0	0	0
	5-60	0.0-5.0	6.1-7.8	0	0	0	0
Valentine-----	0-6	0.0-6.0	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	5.6-7.3	0	0	0	0
Tn:							
Tryon-----	0-5	10-25	5.6-8.4	0-5	0	0	0
	5-60	0.0-5.0	5.6-7.8	0	0	0	0
To:							
Tryon-----	0-6	10-25	5.6-8.4	0-5	0	0	0
	6-60	0.0-5.0	5.6-7.8	0	0	0	0
TpB:							
Tryon-----	0-5	10-25	5.6-8.4	0-5	0	0	0
	5-60	0.0-5.0	5.6-7.8	0	0	0	0
Els-----	0-6	0.0-5.0	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	6.1-7.3	0	0	0	0
VaB:							
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0
VaD:							
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0
VaE:							
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0
VaF:							
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Brown County, Nebraska

PAGE 6 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
VbB:							
Valentine-----	0-7	2.0-8.0	5.6-7.3	0	0	0	0
	7-13	1.0-8.0	5.6-7.3	0	0	0	0
	13-60	0.0-5.0	5.6-7.3	0	0	0	0
VbD:							
Valentine-----	0-7	2.0-8.0	5.6-7.3	0	0	0	0
	7-13	1.0-8.0	5.6-7.3	0	0	0	0
	13-60	0.0-5.0	5.6-7.3	0	0	0	0
VfD:							
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0
Els-----	0-6	0.0-5.0	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	6.1-7.3	0	0	0	0
VhD:							
Valentine-----	0-6	0.0-6.0	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	5.6-7.3	0	0	0	0
Libory-----	0-10	2.0-12	5.6-7.3	0	0	0	0
	10-24	1.0-9.0	5.6-7.3	0	0	0	0
	24-60	10-22	5.6-7.8	0	0	0	0
VpD:							
Valentine-----	0-5	0.0-6.0	5.6-7.3	0	0	0	0
	5-60	0.0-5.0	5.6-7.3	0	0	0	0
Pivot-----	0-10	3.0-10	5.6-7.3	0	0	0	0
	10-28	1.0-8.0	5.6-7.3	0	0	0	0
	28-60	1.0-6.0	5.6-7.3	0	0	0	0
VrD:							
Valentine-----	0-7	2.0-8.0	5.6-7.3	0	0	0	0
	7-10	1.0-8.0	5.6-7.3	0	0	0	0
	10-60	0.0-5.0	5.6-7.3	0	0	0	0
Sandose-----	0-10	1.0-9.0	5.6-7.3	0	0	0	0
	10-20	1.0-8.0	6.1-7.3	0	0	0	0
	20-40	13-27	6.6-7.8	0-5	0	0	0
	40-60	3.0-15	6.6-8.4	0-10	0	0	0
VsG2:							
Valentine-----	0-6	0.0-6.0	5.6-7.3	0	0	0	0
	6-60	0.0-5.0	5.6-7.3	0	0	0	0
Simeon-----	0-7	0.0-5.0	6.1-7.8	0	0	0	0
	7-60	0.0-5.0	6.1-7.8	0	0	0	0
VtE:							
Valentine-----	0-7	0.0-6.0	5.6-7.3	0	0	0	0
	7-60	0.0-5.0	5.6-7.3	0	0	0	0
Longpine-----	0-6	5.0-10	6.6-8.4	0-5	0	0	0
	6-13	5.0-10	7.4-8.4	2-15	0	0	0
	13-60	---	---	---	---	---	---
VwE:							
Valentine-----	0-4	0.0-6.0	5.6-7.3	0	0	0	0
	4-60	0.0-5.0	5.6-7.3	0	0	0	0
Tryon-----	0-6	10-25	5.6-8.4	0-5	0	0	0
	6-60	0.0-5.0	5.6-7.8	0	0	0	0
VxB:							
Vetal-----	0-16	10-22	5.6-7.8	0	0	0	0
	16-31	10-22	6.1-7.8	0	0	0	0
	31-56	9.0-19	6.1-8.4	0-5	0	0	0
	56-60	7.0-17	6.1-8.4	0-5	0	0	0
zwa:							
Water-----	---	---	---	---	---	---	---
zwb:							
Water-----	---	---	---	---	---	---	---



# WATER FEATURES Brown County, Nebraska

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
Aa: Almeria-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		March	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		April	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		May	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		June	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Ae: Almeria-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		March	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		April	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		May	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		June	0.0-1.5	>6.0	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Af: Almeria-----	D	January	0.0	>6.0	0.0-0.5	Long	---	---	None
		February	0.0	>6.0	0.0-0.5	Long	---	Brief	Frequent
		March	0.0	>6.0	0.0-0.5	Long	---	Brief	Frequent
		April	0.0	>6.0	0.0-0.5	Long	---	Brief	Frequent
		May	0.0	>6.0	0.0-0.5	Long	---	Brief	Frequent
		June	0.0	>6.0	0.0-0.5	Long	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		November	0.0	>6.0	0.0-0.5	Long	---	---	None
		December	0.0	>6.0	0.0-0.5	Long	---	---	None
Histosols-----	D	January	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		February	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		March	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		April	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		May	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		June	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		November	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
		December	0.0	>6.0	0.0-2.0	Very long	---	Long	Frequent
An: Anselmo-----	B		---	---	---	---	---	---	---
AnC: Anselmo-----	B		---	---	---	---	---	---	---
AnD: Anselmo-----	B		---	---	---	---	---	---	---
AtF: Anselmo-----	B		---	---	---	---	---	---	---
Brunswick-----	B		---	---	---	---	---	---	---
Ba: Barney-----	D	January	0.0-1.0	>6.0	---	---	---	---	None
		February	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		March	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		April	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		May	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		June	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		November	0.0-1.0	>6.0	---	---	---	---	None
		December	0.0-1.0	>6.0	---	---	---	---	None
Bd: Bolent-----	A	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Bo: Brocksburg-----	B		---	---	---	---	---	---	---
BrD: Brunswick-----	B		---	---	---	---	---	---	---
DuB:			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Dunday-----	A		Ft	Ft	Ft				
DuD: Dunday-----	A		---	---	---	---	---	---	---
Es: Els-----	A		---	---	---	---	---	---	---
EpB: Els-----	A	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	---	None
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	---	None
Ipage-----	A	November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
		January	3.0-5.0	>6.0	---	---	---	---	None
		February	3.0-5.0	>6.0	---	---	---	---	None
		March	3.0-5.0	>6.0	---	---	---	---	None
		April	3.0-5.0	>6.0	---	---	---	---	None
		May	3.0-5.0	>6.0	---	---	---	---	None
		June	3.0-5.0	>6.0	---	---	---	---	None
		December	3.0-5.0	>6.0	---	---	---	---	None
Es: Elsmere-----	A	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	---	None
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Fe: Fluvaquents-----	D	January	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		February	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		March	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		April	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		May	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		June	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		July	0.0	>6.0	0.0-2.0	Very long	---	---	None
		August	0.0	>6.0	0.0-2.0	Very long	---	---	None
		September	0.0	>6.0	0.0-2.0	Very long	---	---	None
		October	0.0	>6.0	0.0-2.0	Very long	---	---	None
		November	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
		December	0.0	>6.0	0.0-2.0	Very long	---	Brief	Frequent
Gn: Gannett-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	---	None
		April	0.0-1.5	>6.0	---	---	---	---	None
		May	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
IdB: Inavale-----	A	January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
IkB: Inavale-----	A	January	---	---	---	---	---	Very brief	Occasional
		February	---	---	---	---	---	Very brief	Occasional
		March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
In:									



(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
Inavale-----	A	January	---	---	---	---	---	Very brief	Occasional
		February	---	---	---	---	---	Very brief	Occasional
		March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
Barney-----	D	January	0.0-1.0	>6.0	---	---	---	---	None
		February	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		March	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		April	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		May	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		June	0.0-1.0	>6.0	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		November	0.0-1.0	>6.0	---	---	---	---	None
		December	0.0-1.0	>6.0	---	---	---	---	None
IpB:									
Ipge-----	A	January	3.0-5.0	>6.0	---	---	---	---	None
		February	3.0-5.0	>6.0	---	---	---	---	None
		March	3.0-5.0	>6.0	---	---	---	---	None
		April	3.0-5.0	>6.0	---	---	---	---	None
		May	3.0-5.0	>6.0	---	---	---	---	None
		June	3.0-5.0	>6.0	---	---	---	---	None
		December	3.0-5.0	>6.0	---	---	---	---	None
IsB:									
Ipge-----	A	January	3.0-5.0	>6.0	---	---	---	---	None
		February	3.0-5.0	>6.0	---	---	---	---	None
		March	3.0-5.0	>6.0	---	---	---	---	None
		April	3.0-5.0	>6.0	---	---	---	---	None
		May	3.0-5.0	>6.0	---	---	---	---	None
		June	3.0-5.0	>6.0	---	---	---	---	None
		December	3.0-5.0	>6.0	---	---	---	---	None
Jn:									
Jansen-----	B		---	---	---	---	---	---	---
JnC:									
Jansen-----	B		---	---	---	---	---	---	---
Jo:									
Jansen-----	B		---	---	---	---	---	---	---
JoC:									
Jansen-----	B		---	---	---	---	---	---	---
Jr:									
Jansen-----	B		---	---	---	---	---	---	---
Meadin-----	A		---	---	---	---	---	---	---
JtB:									
Jansen-----	B		---	---	---	---	---	---	---
Sandose-----	A		---	---	---	---	---	---	---
Jw:									
Johnstown-----	B		---	---	---	---	---	---	---
JwB:									
Johnstown-----	B		---	---	---	---	---	---	---
Jy:									
Johnstown-----	B		---	---	---	---	---	---	---
JyB:									
Johnstown-----	B		---	---	---	---	---	---	---
JyC:									
Johnstown-----	B		---	---	---	---	---	---	---
LcG:									
Labu-----	D		---	---	---	---	---	---	---
Sansarc-----	D		---	---	---	---	---	---	---
LfB:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Libory-----	A		Ft	Ft	Ft				
		January	1.5-3.0	1.7-3.0	---	---	---	---	None
		March	1.5-3.0	1.5-3.0	---	---	---	---	None
		April	1.5-3.0	1.5-3.0	---	---	---	---	None
		May	1.5-3.0	1.5-3.0	---	---	---	---	None
		June	1.5-3.0	1.5-3.0	---	---	---	---	None
Lo: Loup-----	D								
		January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	---	None
		April	0.0-1.5	>6.0	---	---	---	---	None
		May	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Lp: Loup-----	D								
		January	0.0	>6.0	0.0-0.5	Long	---	---	None
		February	0.0	>6.0	0.0-0.5	Long	---	---	None
		March	0.0	>6.0	0.0-0.5	Long	---	---	None
		April	0.0	>6.0	0.0-0.5	Long	---	---	None
		May	0.0	>6.0	0.0-0.5	Long	---	---	None
		June	0.0	>6.0	0.0-0.5	Long	---	---	None
		November	0.0	>6.0	0.0-0.5	Long	---	---	None
		December	0.0	>6.0	0.0-0.5	Long	---	---	None
LtB: Els-----	A								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	---	None
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Loup-----	D								
		January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	---	None
		April	0.0-1.5	>6.0	---	---	---	---	None
		May	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Ma: Marlake-----	D								
		January	0.0	>6.0	0.0-2.0	Very long	---	---	None
		February	0.0	>6.0	0.0-2.0	Very long	---	---	None
		March	0.0	>6.0	0.0-2.0	Very long	---	---	None
		April	0.0	>6.0	0.0-2.0	Very long	---	---	None
		May	0.0	>6.0	0.0-2.0	Very long	---	---	None
		June	0.0	>6.0	0.0-2.0	Very long	---	---	None
		November	0.0	>6.0	0.0-2.0	Very long	---	---	None
		December	0.0	>6.0	0.0-2.0	Very long	---	---	None
McG: Mckelvie-----	A								
			---	---	---	---	---	---	---
Longpine-----	D								
			---	---	---	---	---	---	---
MeB: Meadin-----	A								
			---	---	---	---	---	---	---
MeF: Meadin-----	A								
			---	---	---	---	---	---	---
Oe: O'Neill-----	B								
			---	---	---	---	---	---	---
On: O'Neill-----	B								
			---	---	---	---	---	---	---
OsC: O'Neill-----	B								
			---	---	---	---	---	---	---
Meadin-----	A								
			---	---	---	---	---	---	---
OsD: O'Neill-----	B								
			---	---	---	---	---	---	---
Meadin-----	A								
			---	---	---	---	---	---	---
Pg: Pits-----	A								
			---	---	---	---	---	---	---
PtB: Pivot-----	A								
			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
RtB: Ronson-----	B		---	---	---	---	---	---	---
Longpine-----	D		---	---	---	---	---	---	---
RtC: Ronson-----	B		---	---	---	---	---	---	---
Longpine-----	D		---	---	---	---	---	---	---
RtD: Ronson-----	B		---	---	---	---	---	---	---
Longpine-----	D		---	---	---	---	---	---	---
ScB: Sandose-----	A		---	---	---	---	---	---	---
SkB: Simeon-----	A		---	---	---	---	---	---	---
SkD: Simeon-----	A		---	---	---	---	---	---	---
SvD: Simeon-----	A		---	---	---	---	---	---	---
Valentine-----	A		---	---	---	---	---	---	---
Tn: Tryon-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	---	None
		April	0.0-1.5	>6.0	---	---	---	---	None
		May	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
To: Tryon-----	D	January	0.0	>6.0	0.0-0.5	Long	---	---	None
		February	0.0	>6.0	0.0-0.5	Long	---	---	None
		March	0.0	>6.0	0.0-0.5	Long	---	---	None
		April	0.0	>6.0	0.0-0.5	Long	---	---	None
		May	0.0	>6.0	0.0-0.5	Long	---	---	None
		November	0.0	>6.0	0.0-0.5	Long	---	---	None
		December	0.0	>6.0	0.0-0.5	Long	---	---	None
TpB: Tryon-----	D	January	0.0-1.5	>6.0	---	---	---	---	None
		February	0.0-1.5	>6.0	---	---	---	---	None
		March	0.0-1.5	>6.0	---	---	---	---	None
		April	0.0-1.5	>6.0	---	---	---	---	None
		May	0.0-1.5	>6.0	---	---	---	---	None
		November	0.0-1.5	>6.0	---	---	---	---	None
		December	0.0-1.5	>6.0	---	---	---	---	None
Els-----	A	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	---	None
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
VaB: Valentine-----	A		---	---	---	---	---	---	---
VaD: Valentine-----	A		---	---	---	---	---	---	---
VaE: Valentine-----	A		---	---	---	---	---	---	---
VaF: Valentine-----	A		---	---	---	---	---	---	---
Valentine-----	A		---	---	---	---	---	---	---
VbB: Valentine-----	A		---	---	---	---	---	---	---
VbD: Valentine-----	A		---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
VfD: Valentine-----	A		---	---	---	---	---	---	---
Els-----	A		---	---	---	---	---	---	---
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	---	None
		April	1.5-3.0	>6.0	---	---	---	---	None
		May	1.5-3.0	>6.0	---	---	---	---	None
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
VhD: Valentine-----	A		---	---	---	---	---	---	---
Libory-----	A		---	---	---	---	---	---	---
		January	1.5-3.0	1.7-3.0	---	---	---	---	None
		March	1.5-3.0	1.5-3.0	---	---	---	---	None
		April	1.5-3.0	1.5-3.0	---	---	---	---	None
		May	1.5-3.0	1.5-3.0	---	---	---	---	None
		June	1.5-3.0	1.5-3.0	---	---	---	---	None
VpD: Valentine-----	A		---	---	---	---	---	---	---
Pivot-----	A		---	---	---	---	---	---	---
VrD: Valentine-----	A		---	---	---	---	---	---	---
Sandose-----	A		---	---	---	---	---	---	---
VsG2: Valentine-----	A		---	---	---	---	---	---	---
Simeon-----	A		---	---	---	---	---	---	---
VtE: Valentine-----	A		---	---	---	---	---	---	---
Longpine-----	D		---	---	---	---	---	---	---
VwE: Valentine-----	A		---	---	---	---	---	---	---
Tryon-----	D		---	---	---	---	---	---	---
		January	0.0	>6.0	0.0-0.5	Long	---	---	None
		February	0.0	>6.0	0.0-0.5	Long	---	---	None
		March	0.0	>6.0	0.0-0.5	Long	---	---	None
		April	0.0	>6.0	0.0-0.5	Long	---	---	None
		May	0.0	>6.0	0.0-0.5	Long	---	---	None
		November	0.0	>6.0	0.0-0.5	Long	---	---	None
		December	0.0	>6.0	0.0-0.5	Long	---	---	None
VxB: Vetal-----	B		---	---	---	---	---	---	---
zwa: Water-----	---		---	---	---	---	---	---	---
zwb: Water-----	---		---	---	---	---	---	---	---
			---	---	---	---	---	---	---



The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top In	Thickness In	Hardness		Uncoated Steel	Concrete
Aa:							
Almeria-----	---	---	---	---	Moderate	High	Low
Ae:							
Almeria-----	---	---	---	---	Moderate	High	Low
Af:							
Almeria-----	---	---	---	---	Moderate	High	Low
Histosols-----	---	---	---	---	---	High	Moderate
An:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
AnC:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
AnD:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
AtF:							
Anselmo-----	---	---	---	---	Moderate	Moderate	Low
Brunswick-----	20-40	Bedrock (paralithic)	---	---	Moderate	High	Low
Ba:							
Barney-----	---	---	---	---	Moderate	High	Low
Bd:							
Bolent-----	---	---	---	---	Moderate	Low	Low
Bo:							
Brocksburg-----	---	---	---	---	Moderate	Low	Low
BrD:							
Brunswick-----	20-40	Bedrock (paralithic)	---	---	Moderate	High	Low
DuB:							
Dunday-----	---	---	---	---	Low	Low	Low
DuD:							
Dunday-----	---	---	---	---	Low	Low	Low
Eo:							
Els-----	---	---	---	---	Moderate	Moderate	Low
EpB:							
Els-----	---	---	---	---	Moderate	Moderate	Low
Ipige-----	---	---	---	---	Moderate	Low	Moderate
Es:							
Elsmere-----	---	---	---	---	Moderate	Moderate	Low
Fe:							
Fluvaquents-----	---	---	---	---	Moderate	High	Low
Gn:							
Gannett-----	---	---	---	---	High	High	Low
IdB:							
Inavale-----	---	---	---	---	Low	Moderate	Low
IkB:							
Inavale-----	---	---	---	---	Low	Moderate	Low
In:							
Inavale-----	---	---	---	---	Low	Moderate	Low
Barney-----	---	---	---	---	Moderate	High	Low
IpB:							
Ipige-----	---	---	---	---	Moderate	Low	Moderate
IsB:							
Ipige-----	---	---	---	---	Moderate	Low	Moderate
Jn:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
JnC:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
Jo:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
JoC:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
Jr:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
Meadin-----	---	---	---	---	Low	Low	Moderate
JtB:							
Jansen-----	---	---	---	---	Moderate	Moderate	Low
Sandose-----	---	---	---	---	Low	Moderate	Low
Jw:							
Johnstown-----	---	---	---	---	Moderate	Moderate	Low
JwB:							
Johnstown-----	---	---	---	---	Moderate	Moderate	Low
Jy:							
Johnstown-----	---	---	---	---	Moderate	Moderate	Low
JyB:							
Johnstown-----	---	---	---	---	Moderate	Moderate	Low
JyC:							
Johnstown-----	---	---	---	---	Moderate	Moderate	Low
LcG:							
Labu-----	20-40	Bedrock (paralithic)	---	---	Low	High	Moderate
Sansarc-----	4-20	Bedrock (paralithic)	---	---	Low	High	Moderate
LfB:							
Libory-----	---	---	---	---	Low	Moderate	Low
Lo:							
Loup-----	---	---	---	---	Moderate	High	Low
Lp:							
Loup-----	---	---	---	---	Moderate	High	Low

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top In	Thickness In	Hardness		Uncoated Steel	Concrete
LtB:							
Els-----	---	---	---	---	Moderate	Moderate	Low
Loup-----	---	---	---	---	Moderate	High	Low
Ma:							
Marlake-----	---	---	---	---	Moderate	High	Low
McG:							
Mckelvie-----	---	---	---	---	Low	Low	Low
Longpine-----	10-20	Bedrock (paralithic)	---	---	Low	High	Low
MeB:							
Meadin-----	---	---	---	---	Low	Low	Moderate
MeF:							
Meadin-----	---	---	---	---	Low	Low	Moderate
Oe:							
O'neill-----	---	---	---	---	Moderate	Moderate	Low
On:							
O'neill-----	---	---	---	---	Moderate	Moderate	Low
OsC:							
O'neill-----	---	---	---	---	Moderate	Moderate	Low
Meadin-----	---	---	---	---	Low	Low	Moderate
OsD:							
O'neill-----	---	---	---	---	Moderate	Moderate	Low
Meadin-----	---	---	---	---	Low	Low	Moderate
Pg:							
Pits-----	---	---	---	---	Low	Low	Low
PtB:							
Pivot-----	---	---	---	---	Low	Low	Low
RtB:							
Ronson-----	20-40	Bedrock (paralithic)	---	---	Moderate	Moderate	Low
Longpine-----	10-20	Bedrock (paralithic)	---	---	Low	High	Low
RtC:							
Ronson-----	20-40	Bedrock (paralithic)	---	---	Moderate	Moderate	Low
Longpine-----	10-20	Bedrock (paralithic)	---	---	Low	High	Low
RtD:							
Ronson-----	20-40	Bedrock (paralithic)	---	---	Moderate	Moderate	Low
Longpine-----	10-20	Bedrock (paralithic)	---	---	Low	High	Low
ScB:							
Sandose-----	---	---	---	---	Low	Moderate	Low
SkB:							
Simeon-----	---	---	---	---	Low	Low	Low
SkD:							
Simeon-----	---	---	---	---	Low	Low	Low
SvD:							
Simeon-----	---	---	---	---	Low	Low	Low
Valentine-----	---	---	---	---	Low	Low	Low
Tn:							
Tryon-----	---	---	---	---	Moderate	High	Low
To:							
Tryon-----	---	---	---	---	Moderate	High	Low
TpB:							
Tryon-----	---	---	---	---	Moderate	High	Low
Els-----	---	---	---	---	Moderate	Moderate	Low
VaB:							
Valentine-----	---	---	---	---	Low	Low	Low
VaD:							
Valentine-----	---	---	---	---	Low	Low	Low
VaE:							
Valentine-----	---	---	---	---	Low	Low	Low
VaF:							
Valentine-----	---	---	---	---	Low	Low	Low
Valentine-----	---	---	---	---	Low	Low	Low
VbB:							
Valentine-----	---	---	---	---	Low	Low	Low
VbD:							
Valentine-----	---	---	---	---	Low	Low	Low
VfD:							
Valentine-----	---	---	---	---	Low	Low	Low
Els-----	---	---	---	---	Moderate	Moderate	Low
VhD:							
Valentine-----	---	---	---	---	Low	Low	Low
Libory-----	---	---	---	---	Low	Moderate	Low
VpD:							
Valentine-----	---	---	---	---	Low	Low	Low
Pivot-----	---	---	---	---	Low	Low	Low
VrD:							
Valentine-----	---	---	---	---	Low	Low	Low
Sandose-----	---	---	---	---	Low	Moderate	Low
VsG2:							
Valentine-----	---	---	---	---	Low	Low	Low
Simeon-----	---	---	---	---	Low	Low	Low



Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
		In	In				
VtE:							
Valentine-----	---	---	---	---	Low	Low	Low
Longpine-----	10-20	Bedrock (paralithic)	---	---	Low	High	Low
VwE:							
Valentine-----	---	---	---	---	Low	Low	Low
Tryon-----	---	---	---	---	Moderate	High	Low
VxB:							
Vetal-----	---	---	---	---	Moderate	Moderate	Low
zwa:							
Water-----	---	---	---	---	---	---	---
zwb:							
Water-----	---	---	---	---	---	---	---



WATER MANAGEMENT  
Brown County, Nebraska

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Aa: Almeria-----	Limitation: flooding cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
Ae: Almeria-----	Limitation: flooding cutbanks cave	Limitation: wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
Af: Almeria-----	Limitation: flooding ponding cutbanks cave	Limitation: ponding droughty	Limitation: too sandy ponding	Limitation: rooting depth wetness droughty
Histosols-----	Limitation: flooding subsides ponding	Limitation: flooding ponding	Limitation: ponding	Limitation: wetness
An: Anselmo-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
AnC: Anselmo-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: soil blowing	Favorable
AnD: Anselmo-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope soil blowing	Limitation: slope
AtF: Anselmo-----	Limitation: deep to water	Limitation: slope	Limitation: slope soil blowing	Limitation: slope
Brunswick-----	Limitation: deep to water	Limitation: slope droughty	Limitation: slope soil blowing depth to rock	Limitation: slope depth to rock droughty
Ba: Barney-----	Limitation: flooding cutbanks cave	Limitation: rooting depth wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
Bd: Bolent-----	Limitation: flooding cutbanks cave	Limitation: wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: rooting depth droughty
Bo: Brocksburg-----	Limitation: deep to water	Favorable	Limitation: too sandy	Favorable
BrD: Brunswick-----	Limitation: deep to water	Limitation: slope droughty	Limitation: soil blowing depth to rock	Limitation: depth to rock droughty
DuB: Dunday-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
DuD: Dunday-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Eo: Els-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty
EpB: Els-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty
Ipage-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
Es: Elsmere-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Fe: Fluvaquents-----	Limitation: flooding ponding	Limitation: rooting depth ponding droughty	Limitation: too sandy ponding	Limitation: rooting depth wetness droughty
Gn: Gannett-----	Limitation: frost action cutbanks cave	Limitation: wetness droughty	Limitation: too sandy wetness	Limitation: wetness droughty
IdB: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
IkB: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
In: Inavale-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
Barney-----	Limitation: flooding cutbanks cave	Limitation: rooting depth wetness droughty	Limitation: too sandy wetness	Limitation: rooting depth wetness droughty
IpB: Ipage-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
IsB: Ipage-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
Jn: Jansen-----	Limitation: deep to water	Limitation: droughty	Limitation: erodes easily too sandy	Limitation: erodes easily droughty
JnC: Jansen-----	Limitation: deep to water	Limitation: slope droughty	Limitation: erodes easily too sandy	Limitation: erodes easily droughty
Jo: Jansen-----	Limitation: deep to water	Favorable	Limitation: erodes easily too sandy	Limitation: erodes easily
JoC: Jansen-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily too sandy	Limitation: erodes easily
Jr: Jansen-----	Limitation: deep to water	Limitation: droughty	Limitation: erodes easily too sandy	Limitation: erodes easily droughty
Meadin-----	Limitation: deep to water	Limitation: droughty	Limitation: too sandy soil blowing	Limitation: droughty
JtB: Jansen-----	Limitation: deep to water	Limitation: droughty	Limitation: erodes easily too sandy	Limitation: erodes easily droughty
Sandose-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: erodes easily soil blowing	Limitation: erodes easily droughty
Jw: Johnstown-----	Limitation: deep to water	Limitation: soil blowing	Limitation: erodes easily soil blowing	Limitation: erodes easily
JwB: Johnstown-----	Limitation: deep to water	Limitation: soil blowing	Limitation: erodes easily soil blowing	Limitation: erodes easily
Jy: Johnstown-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
JyB: Johnstown-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
JyC: Johnstown-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
LcG: Labu-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: percs slowly slope depth to rock	Limitation: slope depth to rock droughty

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Sansarc-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope droughty
LfB: Libory-----	Favorable	Limitation: wetness droughty	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily droughty
Lo: Loup-----	Limitation: cutbanks cave	Limitation: wetness droughty	Limitation: too sandy wetness	Limitation: wetness droughty
Lp: Loup-----	Limitation: ponding cutbanks cave	Limitation: ponding droughty	Limitation: too sandy ponding	Limitation: wetness droughty
LtB: Els-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty
Loup-----	Limitation: cutbanks cave	Limitation: wetness droughty	Limitation: too sandy wetness	Limitation: wetness droughty
Ma: Marlake-----	Limitation: ponding cutbanks cave	Limitation: ponding droughty	Limitation: too sandy ponding	Limitation: wetness droughty
McG: Mckelvie-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
Longpine-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope depth to rock	Limitation: slope depth to rock
MeB: Meadin-----	Limitation: deep to water	Limitation: droughty	Limitation: too sandy soil blowing	Limitation: droughty
MeF: Meadin-----	Limitation: deep to water	Limitation: slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Oe: O'Neill-----	Limitation: deep to water	Limitation: droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
On: O'Neill-----	Limitation: deep to water	Limitation: droughty	Limitation: too sandy	Limitation: rooting depth droughty
OsC: O'Neill-----	Limitation: deep to water	Limitation: slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Meadin-----	Limitation: deep to water	Limitation: slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
OsD: O'Neill-----	Limitation: deep to water	Limitation: slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
Meadin-----	Limitation: deep to water	Limitation: slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Pg: Pits-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy	Limitation: rooting depth slope droughty
PtB: Pivot-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
RtB: Ronson-----	Limitation: deep to water	Limitation: soil blowing droughty	Limitation: area reclaim soil blowing	Limitation: area reclaim droughty
Longpine-----	Limitation: deep to water	Limitation: soil blowing	Limitation: depth to rock	Limitation: depth to rock

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
RtC: Ronson-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: area reclaim soil blowing	Limitation: area reclaim droughty
Longpine-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: depth to rock	Limitation: depth to rock
RtD: Ronson-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: area reclaim slope soil blowing	Limitation: area reclaim slope droughty
Longpine-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope depth to rock	Limitation: slope depth to rock
ScB: Sandose-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: erodes easily soil blowing	Limitation: erodes easily droughty
SkB: Simeon-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
SkD: Simeon-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
SvD: Simeon-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Tn: Tryon-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness	Limitation: wetness droughty
To: Tryon-----	Limitation: ponding cutbanks cave	Limitation: fast intake ponding droughty	Limitation: too sandy ponding	Limitation: wetness droughty
TpB: Tryon-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness	Limitation: wetness droughty
Els-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty
VaB: Valentine-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
VaD: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
VaE: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
VaF: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
VbB: Valentine-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
VbD: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
VfD: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Els-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty
VhD: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Libory-----	Favorable	Limitation: wetness droughty	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily droughty
VpD: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Pivot-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: too sandy soil blowing	Limitation: droughty
VrD: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: rooting depth droughty
Sandose-----	Limitation: deep to water	Limitation: fast intake droughty	Limitation: erodes easily soil blowing	Limitation: erodes easily droughty
VsG2: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
Simeon-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
VtE: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
Longpine-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope depth to rock	Limitation: slope depth to rock
VvE: Valentine-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: rooting depth slope droughty
Tryon-----	Limitation: ponding cutbanks cave	Limitation: fast intake ponding droughty	Limitation: too sandy ponding	Limitation: wetness droughty
VxB: Vetal-----	Limitation: deep to water	Favorable	Favorable	Favorable
zwa: Water-----	---	---	---	---
zwb: Water-----	---	---	---	---



WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.97	Very limited Cutbanks cave	1.00
Ae: Almeria-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.97	Very limited Cutbanks cave	1.00
Af: Almeria-----	55	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.97	Very limited Cutbanks cave	1.00
Histosols-----	45	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
An: Anselmo-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
AnC: Anselmo-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
AnD: Anselmo-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
AtF: Anselmo-----	60	Very limited Seepage Slope	1.00 0.24	Somewhat limited Seepage	0.96	Very limited Deep to water	1.00
Brunswick-----	40	Very limited Seepage Slope Depth to bedrock	1.00 0.15 0.11	Somewhat limited Thin layer Seepage	0.85 0.09	Very limited Deep to water	1.00
Ba: Barney-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.97	Very limited Cutbanks cave	1.00
Bd: Bolent-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
Bo: Brocksburg-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.95	Very limited Deep to water	1.00
BrD: Brunswick-----	100	Very limited Seepage Depth to bedrock	1.00 0.11	Somewhat limited Seepage Thin layer	0.97 0.85	Very limited Deep to water	1.00
DuB: Dunday-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
DuD: Dunday-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Eo: Els-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
EpB: Els-----	65	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
IpB:-----	35	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Cutbanks cave Deep to water	1.00 0.81
Es: Elsmere-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
Fe: Fluvaquents-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.70	Very limited Cutbanks cave	1.00
Gn: Gannett-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
IdB: Inavale-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
IkB: Inavale-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
In: Inavale-----	60	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
Barney-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.97	Very limited Cutbanks cave	1.00
IpB: IpB:-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Cutbanks cave Deep to water	1.00 0.81
IsB: IpB:-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Cutbanks cave Deep to water	1.00 0.81
Jn: Jansen-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
JnC: Jansen-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Jo: Jansen-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
JoC: Jansen-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Jr: Jansen-----	50	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Meadin-----	50	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
JtB: Jansen-----	50	Very limited		Very limited		Very limited	

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sandose-----	50	Seepage	1.00	Seepage	1.00	Deep to water	1.00
		Very limited Seepage	1.00	Somewhat limited Seepage	0.20	Very limited Deep to water	1.00
Jw: Johnstown-----	100	Very limited Seepage	1.00	Somewhat limited Seepage Piping	0.95 0.82	Very limited Deep to water	1.00
JwB: Johnstown-----	100	Very limited Seepage	1.00	Somewhat limited Seepage Piping	0.95 0.82	Very limited Deep to water	1.00
Jy: Johnstown-----	100	Very limited Seepage	1.00	Somewhat limited Seepage Piping	0.95 0.55	Very limited Deep to water	1.00
JyB: Johnstown-----	100	Very limited Seepage	1.00	Somewhat limited Seepage Piping	0.95 0.55	Very limited Deep to water	1.00
JyC: Johnstown-----	100	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.55 0.22	Very limited Deep to water	1.00
LcG: Labu-----	55	Somewhat limited Slope Depth to bedrock	0.15 0.11	Somewhat limited Thin layer	0.85	Very limited Deep to water	1.00
Sansarc-----	45	Somewhat limited Depth to bedrock Slope	0.76 0.32	Very limited Thin layer Hard to pack	1.00 1.00	Very limited Deep to water	1.00
LfB: Libory-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping Seepage	1.00 1.00 0.62	Very limited Cutbanks cave Slow refill Deep to water	1.00 0.30 0.02
Lo: Loup-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
Lp: Loup-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
LtB: Els-----	50	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
Loup-----	50	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
Ma: Marlake-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.17	Very limited Cutbanks cave	1.00
McG: Mckelvie-----	50	Very limited Seepage Slope	1.00 0.94	Very limited Seepage	1.00	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Longpine-----	30	Somewhat limited Slope Depth to bedrock Seepage	0.94 0.66 0.05	Very limited Thin layer Seepage	1.00 0.08	Very limited Deep to water	1.00
MeB: Meadin-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
MeF: Meadin-----	100	Very limited Seepage Slope	1.00 0.06	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
Oe: O'Neill-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.97	Very limited Deep to water	1.00
On: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
OsC: O'Neill-----	60	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Meadin-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
OsD: O'Neill-----	55	Very limited Seepage	1.00	Somewhat limited Seepage	0.97	Very limited Deep to water	1.00
Meadin-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
Pg: Pits-----	100	Not rated		Not rated		Not rated	
PtB: Pivot-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.96	Very limited Deep to water	1.00
RtB: Ronson-----	55	Very limited Seepage Depth to bedrock	1.00 0.11	Somewhat limited Thin layer Seepage	0.85 0.09	Very limited Deep to water	1.00
Longpine-----	45	Somewhat limited Depth to bedrock Seepage	0.66 0.05	Very limited Thin layer Seepage	1.00 0.08	Very limited Deep to water	1.00
RtC: Ronson-----	55	Very limited Seepage Depth to bedrock	1.00 0.11	Somewhat limited Thin layer Seepage	0.85 0.09	Very limited Deep to water	1.00
Longpine-----	45	Somewhat limited Depth to bedrock Seepage	0.66 0.05	Very limited Thin layer Seepage	1.00 0.08	Very limited Deep to water	1.00
RtD: Ronson-----	55	Very limited Seepage Depth to bedrock	1.00 0.11	Somewhat limited Thin layer Seepage	0.85 0.09	Very limited Deep to water	1.00
Longpine-----	45	Somewhat limited Depth to bedrock Seepage	0.66 0.05	Very limited Thin layer Seepage	1.00 0.08	Very limited Deep to water	1.00
ScB: Sandose-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
SkB: Simeon-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
SkD: Simeon-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
SvD: Simeon-----	60	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
Valentine-----	40	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Tn: Tryon-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
To: Tryon-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
TpB: Tryon-----	55	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
Els-----	45	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
VaB: Valentine-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
VaD: Valentine-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
VaE: Valentine-----	100	Very limited Seepage Slope	1.00 0.06	Very limited Seepage	1.00	Very limited Deep to water	1.00
VaF: Valentine-----	70	Very limited Seepage Slope	1.00 0.06	Very limited Seepage	1.00	Very limited Deep to water	1.00
Valentine-----	30	Very limited Seepage Slope	1.00 0.92	Very limited Seepage	1.00	Very limited Deep to water	1.00
VbB: Valentine-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
VbD: Valentine-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
VfD: Valentine-----	55	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Els-----	45	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
VhD: Valentine-----	60	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Libory-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Cutbanks cave	1.00

WATER MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
				Piping Seepage	0.99 0.62	Slow refill Deep to water	0.96 0.02
VpD: Valentine-----	60	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Pivot-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.96	Very limited Deep to water	1.00
VrD: Valentine-----	65	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Sandose-----	35	Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.07	Very limited Deep to water	1.00
VsG2: Valentine-----	65	Very limited Seepage Slope	1.00 0.28	Very limited Seepage	1.00	Very limited Deep to water	1.00
Simeon-----	35	Very limited Seepage Slope	1.00 0.12	Somewhat limited Seepage	0.91	Very limited Deep to water	1.00
VtE: Valentine-----	60	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Longpine-----	40	Somewhat limited Depth to bedrock Seepage	0.66 0.05	Very limited Thin layer Seepage	1.00 0.08	Very limited Deep to water	1.00
VwE: Valentine-----	65	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Tryon-----	35	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Cutbanks cave	1.00
VxB: Vetal-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	



SANITARY FACILITIES  
Brown County, Nebraska

Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.



SANITARY FACILITIES  
Brown County, Nebraska

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Ae: Almeria-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Af: Almeria-----	55	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Histosols-----	45	Very limited Flooding Depth to saturated zone Filtering capacity Subsidence	1.00 1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone Content of organic matter	1.00 1.00 1.00 1.00
An: Anselmo-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
AnC: Anselmo-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.33
AnD: Anselmo-----	100	Very limited Filtering capacity Slope	1.00 0.04	Very limited Seepage Slope	1.00 1.00
AtF: Anselmo-----	60	Very limited Slope Filtering capacity	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Brunswick-----	40	Very limited Depth to bedrock Slope Filtering capacity	1.00 1.00 1.00	Very limited Depth to soft bedrock Slope Seepage	1.00 1.00 1.00
Ba: Barney-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Bd: Bolent-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
Bo: Brocksburg-----	100	Very limited Filtering capacity Restricted permeability	1.00 0.71	Very limited Seepage	1.00
BrD: Brunswick-----	100	Very limited Depth to bedrock Filtering capacity	1.00 1.00	Very limited Depth to soft bedrock Seepage	1.00 1.00

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
DuB: Dunday-----	100	Very limited Filtering capacity	1.00	Slope	0.91
DuD: Dunday-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
Es: Els-----	100	Very limited Depth to saturated zone Filtering capacity	1.00	Slope	0.00
EpB: Els-----	65	Very limited Depth to saturated zone Filtering capacity	1.00	Very limited Seepage	1.00
IpB: IpB-----	100	Very limited Filtering capacity	1.00	Depth to saturated zone	1.00
Es: Elsmere-----	100	Very limited Depth to saturated zone Filtering capacity	1.00	Slope	0.91
Fe: Fluvaquents-----	100	Very limited Flooding Depth to saturated zone	1.00	Very limited Seepage	1.00
Gn: Gannett-----	100	Very limited Depth to saturated zone Filtering capacity	1.00	Depth to saturated zone	1.00
IdB: Inavale-----	100	Very limited Filtering capacity Flooding	1.00	Very limited Seepage	1.00
IkB: Inavale-----	100	Very limited Flooding Filtering capacity	1.00	Depth to saturated zone	1.00
In: Inavale-----	60	Very limited Flooding Filtering capacity	1.00	Flooding Slope	0.40
Barney-----	40	Very limited Flooding Depth to saturated zone Filtering capacity	1.00	Slope	0.00
IpB: IpB-----	100	Very limited Filtering capacity Depth to saturated zone	1.00	Very limited Flooding Seepage	1.00
IsB: IpB-----	100	Very limited	1.00	Depth to saturated zone	1.00
				Slope	0.71
				Slope	0.00

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Jn: Jansen-----	100	Filtering capacity	1.00	Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	0.71
				Slope	0.00
JnC: Jansen-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Restricted permeability	0.71		
Jo: Jansen-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Restricted permeability	0.71	Slope	0.33
JoC: Jansen-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Restricted permeability	1.00	Slope	0.33
Jr: Jansen-----	50	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Meadin-----	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00		
JtB: Jansen-----	50	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Restricted permeability	1.00		
Sandose-----	50	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Restricted permeability	0.71	Slope	0.00
Jw: Johnstown-----	100	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00		
JwB: Johnstown-----	100	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00	Slope	0.00
Jy: Johnstown-----	100	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00		
JyB: Johnstown-----	100	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00	Slope	0.00
JyC: Johnstown-----	100	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00	Slope	0.67
LcG: Labu-----	55	Very limited		Very limited	

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Sansarc-----	45	Depth to bedrock	1.00	Depth to soft bedrock	1.00
		Slope	1.00	Slope	1.00
		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to soft bedrock	1.00
LfB: Libory-----	100	Slope	1.00	Slope	1.00
		Very limited		Very limited	
Lo: Loup-----	100	Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	0.00
		Restricted permeability	0.50	Slope	0.00
		Very limited		Very limited	
Lp: Loup-----	100	Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
LtB: Els-----	50	Very limited		Very limited	
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
		Very limited		Slope	0.00
Loup-----	50	Depth to saturated zone	1.00	Very limited	1.00
		Filtering capacity	1.00	Seepage	1.00
Ma: Marlake-----	100	Very limited		Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Seepage	1.00
McG: Mckelvie-----	50	Filtering capacity	1.00	Depth to saturated zone	1.00
		Very limited		Very limited	
Longpine-----	30	Slope	1.00	Slope	1.00
		Filtering capacity	1.00	Seepage	1.00
MeB: Meadin-----	100	Very limited		Very limited	
		Depth to bedrock	1.00	Depth to soft bedrock	1.00
MeF: Meadin-----	100	Slope	1.00	Slope	1.00
		Very limited		Very limited	
Oe: O'Neill-----	100	Filtering capacity	1.00	Seepage	1.00
		Very limited		Very limited	
On: O'Neill-----	100	Filtering capacity	1.00	Seepage	1.00
		Very limited		Very limited	
OsC: O'Neill-----	60	Filtering capacity	1.00	Seepage	1.00
		Very limited		Very limited	
Meadin-----	40	Very limited		Slope	0.33
		Filtering capacity	1.00	Very limited	1.00

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
OsD: O'Neill-----	55	Very limited Filtering capacity Slope	1.00 0.04	Slope Very limited Seepage	0.33 1.00
Meadin-----	45	Very limited Filtering capacity Slope	1.00 0.04	Slope Very limited Seepage	1.00 1.00
Pg: Pits-----	100	Not rated		Slope Not rated	1.00
PtB: Pivot-----	100	Very limited Filtering capacity	1.00	Not rated	
RtB: Ronson-----	55	Very limited Depth to bedrock	1.00	Very limited Seepage Slope	1.00 0.00
Longpine-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Seepage Slope	1.00 1.00 0.00
RtC: Ronson-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Seepage Slope	1.00 1.00 0.67
Longpine-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Slope	1.00 0.67
RtD: Ronson-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Seepage Slope	1.00 1.00 1.00
Longpine-----	45	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to soft bedrock Slope	1.00 1.00
ScB: Sandose-----	100	Very limited Filtering capacity Restricted permeability	1.00 0.50	Very limited Seepage Slope	1.00 0.00
SkB: Simeon-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
SkD: Simeon-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
SvD: Simeon-----	60	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
Valentine-----	40	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
Tn: Tryon-----	100	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
To: Tryon-----	100	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
TpB: Tryon-----	55	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
Els-----	45	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 1.00 0.00
VaB: Valentine-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
VaD: Valentine-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
VaE: Valentine-----	100	Very limited Slope Filtering capacity	1.00 1.00	Very limited Slope Seepage	1.00 1.00
VaF: Valentine-----	70	Very limited Slope Filtering capacity	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Valentine-----	30	Very limited Slope Filtering capacity	1.00 1.00	Very limited Slope Seepage	1.00 1.00
VbB: Valentine-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.00
VbD: Valentine-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
VfD: Valentine-----	55	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
Els-----	45	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
VhD: Valentine-----	60	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
Libory-----	40	Very limited Depth to saturated zone Restricted permeability Filtering capacity	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 0.00 0.00
VpD: Valentine-----	60	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
Pivot-----	40	Very limited		Very limited	

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
VrD: Valentine-----	65	Filtering capacity	1.00	Seepage	1.00
				Slope	0.00
Sandose-----	35	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.91
VsG2: Valentine-----	65	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
				Slope	0.00
Simeon-----	35	Very limited Slope	1.00	Very limited Slope	1.00
		Filtering capacity	1.00	Seepage	1.00
VtE: Valentine-----	60	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.16	Slope	1.00
Longpine-----	40	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	0.16	Seepage	1.00
VwE: Valentine-----	65	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.16	Slope	1.00
Tryon-----	35	Very limited Depth to saturated zone	1.00	Very limited Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
VxB: Vetal-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.00
zwa: Water-----	100	Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated	



SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage Depth to saturated zone	1.00 1.00 1.00
Ae: Almeria-----	100	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage Depth to saturated zone	1.00 1.00 1.00
Af: Almeria-----	55	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Too Sandy Seepage	1.00 1.00 1.00
Histosols-----	45	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Too Sandy Seepage	1.00 1.00 1.00
An: Anselmo-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
AnC: Anselmo-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
AnD: Anselmo-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 0.04	Very limited Seepage Too Sandy Slope	1.00 0.50 0.04
AtF: Anselmo-----	60	Very limited Slope Seepage Too Sandy	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00	Very limited Slope Too Sandy Seepage	1.00 1.00 1.00
Brunswick-----	40	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to bedrock Slope Seepage	1.00 1.00 0.50
Ba: Barney-----	100	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Too Sandy Seepage	1.00 1.00 1.00
Bd: Bolent-----	100	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 0.68
Bo: Brocksburg-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
BrD: Brunswick-----	100	Very limited Depth to bedrock Too Sandy	1.00 1.00	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock Too Sandy Seepage	1.00 1.00 1.00
DuB: Dunday-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
DuD: Dunday-----	100	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Eo: Els-----	100	Seepage Too Sandy	1.00 1.00	Seepage	1.00	Too Sandy Seepage	1.00 1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage Depth to saturated zone	1.00 0.68
EpB: Els-----	65	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage Depth to saturated zone	1.00 0.68
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
Ipage-----	35	Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
Es: Elsmere-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage Depth to saturated zone	1.00 0.68
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
Fe: Fluvaquents-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Too Sandy	0.50
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
Gn: Gannett-----	100	Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage Depth to saturated zone	1.00 1.00
		Very limited Seepage Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage	1.00
		Very limited Flooding Seepage	1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Very limited Seepage	1.00
IdB: Inavale-----	100	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage	1.00
		Very limited Flooding Seepage	1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Very limited Seepage	1.00
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
In: Inavale-----	60	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Too Sandy Seepage	1.00 1.00	Too Sandy Seepage	1.00 1.00	Seepage	1.00
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
Barney-----	40	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Too Sandy Seepage	1.00 1.00	Too Sandy Seepage	1.00 1.00	Seepage	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
IpB: Ipage-----	100	Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
IsB: Ipage-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
Jn: Jansen-----	100	Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
JnC: Jansen-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00
		Seepage Too Sandy	1.00 1.00	Seepage	1.00	Seepage	1.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Too Sandy	1.00

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Jo: Jansen-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
JoC: Jansen-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
Jr: Jansen-----	50	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
Meadin-----	50	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage Gravel content	1.00 1.00 0.93
JtB: Jansen-----	50	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
Sandose-----	50	Very limited Too Sandy	1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
Jw: Johnstown-----	100	Very limited Seepage Too clayey	1.00 0.50	Very limited Seepage	1.00	Somewhat limited Too clayey	0.50
JwB: Johnstown-----	100	Very limited Seepage Too clayey	1.00 0.50	Very limited Seepage	1.00	Somewhat limited Too clayey	0.50
Jy: Johnstown-----	100	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
JyB: Johnstown-----	100	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
JyC: Johnstown-----	100	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
LcG: Labu-----	55	Very limited Depth to bedrock Too clayey Slope	1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Too clayey Hard to compact Slope	1.00 1.00 1.00 1.00
Sansarc-----	45	Very limited Depth to bedrock Too clayey Slope	1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Too clayey Hard to compact Slope	1.00 1.00 1.00 1.00
LfB: Libory-----	100	Very limited Depth to saturated zone	1.00	Very limited Seepage  Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
Lo: Loup-----	100	Very limited Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Too Sandy  Seepage Depth to saturated zone	1.00 1.00 1.00
Lp: Loup-----	100	Very limited Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Too Sandy Seepage	1.00 1.00 1.00
LtB: Els-----	50	Very limited Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Too Sandy  Seepage Depth to saturated zone	1.00 1.00 1.00 0.68
Loup-----	50	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ma: Marlake-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Depth to saturated zone	1.00
McG: Mckelvie-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Too Sandy	0.50
Longpine-----	30	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00	Slope	1.00	Seepage	1.00
		Slope	1.00			Slope	1.00
MeB: Meadin-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Slope	1.00
MeF: Meadin-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
		Slope	1.00			Gravel content	0.89
Oe: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
On: O'Neill-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
OsC: O'Neill-----	60	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Meadin-----	40	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
						Gravel content	0.88
OsD: O'Neill-----	55	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00	Slope	0.04	Seepage	1.00
		Slope	0.04			Slope	0.04
Meadin-----	45	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
		Too Sandy	1.00	Slope	0.04	Seepage	1.00
		Slope	0.04			Gravel content	0.87
Pg: Pits-----	100	Not rated		Not rated		Not rated	0.04
PtB: Pivot-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Too Sandy	1.00
RtB: Ronson-----	55	Very limited Depth to bedrock	1.00	Very limited Seepage	1.00	Very limited Depth to bedrock	1.00
Longpine-----	45	Seepage	1.00	Depth to bedrock	1.00	Seepage	0.50
		Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
						Seepage	0.50
RtC: Ronson-----	55	Very limited Depth to bedrock	1.00	Very limited Seepage	1.00	Gravel content	0.01
		Seepage	1.00	Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
						Seepage	0.50
Longpine-----	45	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
RtD: Ronson-----	55	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Longpine-----	45	Depth to bedrock	1.00	Seepage	1.00	Depth to bedrock	1.00
		Seepage	1.00	Depth to bedrock	1.00	Seepage	0.50
		Slope	0.04	Slope	0.04	Slope	0.04
		Very limited		Very limited		Very limited	
ScB: Sandose-----	100	Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Slope	0.04	Slope	0.04	Slope	0.04
		Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Not limited	
SkB: Simeon-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
SkD: Simeon-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
SvD: Simeon-----	60	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Valentine-----	40	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Tn: Tryon-----	100	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Depth to saturated zone	1.00
To: Tryon-----	100	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
TpB: Tryon-----	55	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Depth to saturated zone	1.00
Els-----	45	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Depth to saturated zone	0.68
VaB: Valentine-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
VaD: Valentine-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
VaE: Valentine-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00	Slope	1.00	Seepage	1.00
		Slope	1.00			Slope	1.00
VaF: Valentine-----	70	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00	Slope	1.00	Seepage	1.00
		Slope	1.00			Slope	1.00
Valentine-----	30	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
VbB: Valentine-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
VbD: Valentine-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
VfD: Valentine-----	55	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Els-----	45	Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
		Very limited		Very limited		Very limited	
		Depth to	1.00	Depth to	1.00	Too Sandy	1.00
		saturated zone		saturated zone			
VhD: Valentine-----	60	Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Depth to	0.68
		Very limited		Very limited		saturated zone	
		Depth to	1.00	Seepage	1.00	Very limited	
		saturated zone				Too Sandy	1.00
Libory-----	40	Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Too Sandy	1.00
		Very limited		Very limited		Seepage	
		Depth to	1.00	Seepage	1.00	Depth to	1.00
		saturated zone				saturated zone	
VpD: Valentine-----	60	Too clayey	0.50	Depth to	1.00	Too clayey	0.50
				saturated zone			
		Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
Pivot-----	40	Very limited		Very limited		Seepage	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
						Too Sandy	
						Seepage	1.00
VrD: Valentine-----	65	Seepage	1.00	Seepage	1.00	Very limited	
		Too Sandy	1.00			Too Sandy	1.00
		Very limited		Very limited		Seepage	1.00
		Seepage	1.00	Seepage	1.00	Not limited	
		Too Sandy	1.00				
Sandose-----	35	Seepage	1.00	Seepage	1.00		
VsG2: Valentine-----	65	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00	Slope	1.00	Seepage	1.00
		Slope	1.00			Slope	1.00
Simeon-----	35	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00	Slope	1.00	Seepage	1.00
		Slope	1.00			Slope	1.00
VtE: Valentine-----	60	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00	Slope	0.16	Seepage	1.00
		Slope	0.16			Slope	0.16
Longpine-----	40	Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Slope	0.16	Slope	0.16	Seepage	0.50
						Slope	0.16
VwE: Valentine-----	65	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00	Slope	0.16	Seepage	1.00
		Slope	0.16			Slope	0.16
Tryon-----	35	Very limited		Very limited		Very limited	
		Depth to	1.00	Depth to	1.00	Depth to	1.00
		saturated zone		saturated zone		saturated zone	
		Seepage	1.00	Seepage	1.00	Too Sandy	1.00
		Too Sandy	1.00			Seepage	1.00
VxB: Vetal-----	100	Very limited		Very limited		Somewhat limited	
		Seepage	1.00	Seepage	1.00	Seepage	0.50
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	



# AGRICULTURAL WASTE MANAGEMENT Brown County, Nebraska

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.



AGRICULTURAL WASTE MANAGEMENT  
Brown County, Nebraska

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Aa: Almeria-----	100	Very limited Depth to saturated zone Flooding Filtering capacity Runoff limitation Droughty	1.00 1.00 1.00 0.40 0.04	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.04	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.04
Ae: Almeria-----	100	Very limited Depth to saturated zone Flooding Filtering capacity Runoff limitation Droughty	1.00 1.00 1.00 0.40 0.01	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.01	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.01
Af: Almeria-----	55	Very limited Depth to saturated zone Flooding Filtering capacity Runoff limitation Droughty	1.00 1.00 1.00 0.40 0.00	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.00	Very limited Depth to saturated zone Flooding Filtering capacity Droughty	1.00 1.00 1.00 0.00
Histosols-----	45	Very limited Depth to saturated zone Flooding Filtering capacity Runoff limitation	1.00 1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Flooding Low adsorption Filtering capacity	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Flooding Filtering capacity	1.00 1.00 1.00
An: Anselmo-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
AnC: Anselmo-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity Too steep for surface application	1.00 0.08
AnD: Anselmo-----	100	Very limited Filtering capacity  Slope	1.00  0.04	Very limited Filtering capacity  Slope	1.00  0.04	Very limited Too steep for surface application Filtering capacity Too steep for sprinkler application	1.00 1.00 0.22
AtF: Anselmo-----	60	Very limited Slope  Filtering capacity	1.00  1.00	Very limited Slope  Filtering capacity	1.00  1.00	Very limited Too steep for surface application Too steep for sprinkler application Filtering capacity	1.00 1.00 1.00
Brunswick-----	40	Very limited Slope  Droughty  Depth to bedrock	1.00  0.89  0.42	Very limited Slope  Droughty  Depth to bedrock	1.00  0.89  0.42	Very limited Too steep for surface application Too steep for sprinkler application Droughty Depth to bedrock	1.00 1.00 0.89 0.42
Ba: Barney-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Bd: Bolent-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	1.00	Flooding	1.00
		Depth to dense layer	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Filtering capacity	1.00	Droughty	0.19	Droughty	0.19
		Runoff limitation	0.40				
Bo: Brocksburg-----	100	Very limited Filtering capacity	1.00	Very limited Flooding	1.00	Very limited Filtering capacity	1.00
		Depth to saturated zone	0.95	Filtering capacity	1.00	Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
		Leaching limitation	0.45	Droughty	0.11	Droughty	0.11
		Droughty	0.11				
BrD: Brunswick-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Somewhat limited Droughty	0.97	Somewhat limited Droughty	0.97	Somewhat limited Droughty	0.97
		Depth to bedrock	0.42	Depth to bedrock	0.42	Too steep for surface application	0.66
						Depth to bedrock	0.42
						Too steep for sprinkler application	0.00
DuB: Dunday-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.08	Droughty	0.08
		Droughty	0.08				
DuD: Dunday-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.08	Too steep for surface application	0.66
						Droughty	0.08
		Droughty	0.08			Too steep for sprinkler application	0.00
Eo: Els-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Depth to saturated zone	0.95	Depth to saturated zone	0.95	Depth to saturated zone	0.95
		Droughty	0.50	Droughty	0.50	Droughty	0.50
		Leaching limitation	0.45				
EpB: Els-----	65	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Depth to saturated zone	0.95	Depth to saturated zone	0.95	Depth to saturated zone	0.95
		Droughty	0.54	Droughty	0.54	Droughty	0.54
		Leaching limitation	0.45				
Ipage-----	35	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Droughty	0.61	Droughty	0.61	Droughty	0.61
		Leaching limitation	0.45	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
Es: Elsmere-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Fe: Fluvaquents-----	100	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to saturated zone	0.95	Depth to saturated zone	0.95	Depth to saturated zone	0.95
		Droughty	0.58	Droughty	0.58	Droughty	0.58
		Leaching limitation	0.45				
Gn: Gannett-----	100	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	1.00	Flooding	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
IdB: Inavale-----	100	Runoff limitation	0.40				
		Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
IkB: Inavale-----	100	Runoff limitation	0.40	Droughty	0.00	Droughty	0.00
		Droughty	0.00				
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
In: Inavale-----	100	Leaching limitation	0.45	Flooding	0.40	Droughty	0.18
		Droughty	0.18	Droughty	0.18		
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Flooding	1.00	Filtering capacity	1.00
IpB: Ipage-----	100	Flooding	0.60	Filtering capacity	1.00	Flooding	0.60
		Leaching limitation	0.45	Droughty	0.29	Droughty	0.29
		Droughty	0.29				
		Very limited		Very limited		Very limited	
IsB: Ipage-----	100	Filtering capacity	1.00	Flooding	1.00	Filtering capacity	1.00
		Depth to saturated zone	1.00	Filtering capacity	1.00	Flooding	0.60
		Flooding	1.00	Droughty	0.15	Droughty	0.15
		Leaching limitation	0.45				
Jn: Jansen-----	100	Droughty	0.15	Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	1.00	Flooding	1.00
		Depth to dense layer	1.00	Filtering capacity	1.00	Filtering capacity	1.00
Jn: Jansen-----	100	Filtering capacity	1.00	Droughty	0.26	Droughty	0.26
		Runoff limitation	0.40				
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
Jn: Jansen-----	100	Droughty	0.61	Droughty	0.61	Droughty	0.61
		Leaching limitation	0.45	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
		Very limited		Very limited		Very limited	
Jn: Jansen-----	100	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty	0.47	Droughty	0.47	Droughty	0.47
		Leaching limitation	0.45	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
Jn: Jansen-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Too acid	0.02	Too acid	0.07	Too acid	0.07

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
JnC: Jansen-----	100	Droughty	0.00	Droughty	0.00	Droughty	0.00
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Too acid	0.02	Too acid	0.07	Too steep for surface application	0.08
Jo: Jansen-----	100	Droughty	0.00	Droughty	0.00	Too acid	0.07
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Droughty	0.00
		Too acid	0.02	Too acid	0.07	Very limited Filtering capacity	1.00
JoC: Jansen-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Too acid	0.07
		Restricted permeability	0.43	Restricted permeability	0.32	Very limited Filtering capacity	1.00
		Too acid	0.02	Too acid	0.07	Restricted permeability	0.32
Jr: Jansen-----	50	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Too steep for surface application	0.08
		Restricted permeability	0.43	Restricted permeability	0.32	Too acid	0.07
		Droughty	0.27	Droughty	0.27	Very limited Filtering capacity	1.00
Meadin-----	50	Too acid	0.02	Too acid	0.07	Restricted permeability	0.32
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Too steep for surface application	0.08
		Droughty	0.97	Droughty	0.97	Too acid	0.07
JtB: Jansen-----	50	Leaching limitation	0.45	Too acid	0.07	Very limited Filtering capacity	1.00
		Too acid	0.02	Very limited Filtering capacity	1.00	Restricted permeability	0.32
		Very limited Filtering capacity	1.00	Restricted permeability	0.32	Too acid	0.07
Sandose-----	50	Too acid	0.02	Too acid	0.07	Very limited Filtering capacity	1.00
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Leaching limitation	0.45	Filtering capacity			
Jw: Johnstown-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00		
		Restricted permeability	0.43	Restricted permeability	0.32	Very limited Filtering capacity	1.00
						Restricted permeability	0.32
JwB: Johnstown-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Restricted permeability	0.43	Restricted permeability	0.32	Restricted permeability	0.32
Jy: Johnstown-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Restricted permeability	0.43	Restricted permeability	0.32	Restricted permeability	0.32
JyB: Johnstown-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Restricted permeability	0.43	Restricted permeability	0.32	Restricted permeability	0.32
JyC: Johnstown-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Restricted permeability	0.43	Restricted permeability	0.32	Restricted permeability	0.32
JyC: Johnstown-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
LcG: Labu-----	55	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Restricted permeability	0.43	Restricted permeability	0.32	Restricted permeability	0.32
						Too steep for surface application	0.31
		Very limited		Very limited		Very limited	
		Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Droughty	1.00	Droughty	1.00	Too steep for surface application	1.00
Sansarc-----	45	Slope	1.00	Slope	1.00	Droughty	1.00
		Depth to bedrock	0.42	Depth to bedrock	0.42	Too steep for sprinkler application	1.00
						Depth to bedrock	0.42
		Runoff limitation	0.40	Very limited		Very limited	
		Restricted permeability	1.00	Droughty	1.00	Droughty	1.00
		Depth to bedrock	1.00	Restricted permeability	1.00	Restricted permeability	1.00
LfB: Libory-----	100	Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
						Too steep for sprinkler application	
		Runoff limitation	0.40			Too steep for sprinkler application	1.00
		Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
Lo: Loup-----	100	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45				
		Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Runoff limitation	0.40	Droughty	0.12	Droughty	0.12
Lp: Loup-----	100	Droughty	0.12				
		Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Runoff limitation	0.40	Droughty	0.12	Droughty	0.12
		Droughty	0.12				
LtB: Els-----	50	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to saturated zone	0.95	Depth to saturated zone	0.95	Depth to saturated zone	0.95
		Droughty	0.50	Droughty	0.50	Droughty	0.50
		Leaching limitation	0.45				
		Very limited		Very limited		Very limited	
Loup-----	50	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Runoff limitation	0.40	Droughty	0.09	Droughty	0.09
		Droughty	0.09				
		Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
Ma: Marlake-----	100	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Runoff limitation	0.40	Droughty	0.00	Droughty	0.00
		Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Runoff limitation	0.40	Droughty	0.00	Droughty	0.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
McG: Mckelvie-----	50	Droughty	0.00				
		Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Too steep for sprinkler application	1.00
		Leaching limitation	0.45	Droughty	0.20	Filtering capacity	1.00
Longpine-----	30	Droughty	0.20			Droughty	0.20
		Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Droughty	1.00	Droughty	1.00
		Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Runoff limitation	0.40	Filtering capacity	0.00	Too steep for sprinkler application	1.00
MeB: Meadin-----	100	Low adsorption	0.08			Low adsorption	0.08
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty	0.92	Droughty	0.92	Droughty	0.92
MeF: Meadin-----	100	Leaching limitation	0.45	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
Oe: O'Neill-----	100	Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Droughty	0.92	Droughty	0.92	Too steep for sprinkler application	1.00
		Leaching limitation	0.45	Too acid	0.07	Droughty	0.92
		Too acid	0.02			Too acid	0.07
On: O'Neill-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to dense layer	1.00	Droughty	0.52	Droughty	0.52
		Droughty	0.52	Too acid	0.42	Too acid	0.42
OsC: O'Neill-----	60	Too acid	0.11				
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to dense layer	1.00	Too acid	0.42	Too acid	0.42
Meadin-----	40	Droughty	0.29	Droughty	0.29	Droughty	0.29
		Too acid	0.11			Too steep for surface application	0.08
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty	0.91	Droughty	0.91	Droughty	0.91
		Leaching limitation	0.45	Too acid	0.07	Too steep for surface application	0.08
		Too acid	0.02			Too acid	0.07

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
OsD: O'Neill-----	55	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to dense layer	1.00	Droughty	0.65	Too steep for surface application	1.00
		Droughty	0.65	Too acid	0.42	Droughty	0.65
Meadin-----	45	Too acid	0.11	Slope	0.04	Too acid	0.42
		Slope	0.04			Too steep for sprinkler application	0.22
						Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
Pg: Pits-----	100	Droughty	0.89	Droughty	0.89	Too steep for surface application	1.00
		Leaching limitation	0.45	Too acid	0.07	Droughty	0.89
		Slope	0.04	Slope	0.04	Too steep for sprinkler application	0.22
		Too acid	0.02			Too acid	0.07
PtB: Pivot-----	100	Not rated		Not rated		Not rated	
RtB: Ronson-----	55	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty	0.81	Droughty	0.81	Droughty	0.81
		Leaching limitation	0.45				
Longpine-----	45	Somewhat limited		Somewhat limited		Somewhat limited	
		Droughty	0.93	Droughty	0.93	Droughty	0.93
		Depth to bedrock	0.42	Depth to bedrock	0.42	Depth to bedrock	0.42
		Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00
RtC: Ronson-----	55	Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Droughty	1.00	Droughty	1.00
		Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Runoff limitation	0.40	Filtering capacity	0.00	Filtering capacity	0.00
Longpine-----	45	Filtering capacity	0.00				
		Somewhat limited		Somewhat limited		Somewhat limited	
		Droughty	0.80	Droughty	0.80	Droughty	0.80
		Depth to bedrock	0.42	Depth to bedrock	0.42	Depth to bedrock	0.42
RtD: Ronson-----	55	Filtering capacity	0.00	Filtering capacity	0.00	Too steep for surface application	0.31
						Filtering capacity	0.00
		Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Droughty	1.00	Droughty	1.00
Longpine-----	45	Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Runoff limitation	0.40	Filtering capacity	0.00	Too steep for surface application	0.31
						Low adsorption	0.13
		Low adsorption	0.13			Filtering capacity	0.00
RtD: Ronson-----	55	Filtering capacity	0.00				
		Very limited		Very limited		Very limited	
		Droughty	0.99	Droughty	0.99	Too steep for surface application	1.00
						Droughty	0.99
Longpine-----	45	Depth to bedrock	0.42	Depth to bedrock	0.42	Depth to bedrock	0.42
		Slope	0.04	Slope	0.04	Too steep for sprinkler application	0.22
		Filtering capacity	0.00	Filtering capacity	0.00		



AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Longpine-----	45	Very limited		Very limited		Filtering capacity	0.00
		Depth to bedrock	1.00	Droughty	1.00	Very limited	
		Droughty	1.00	Depth to bedrock	1.00	Droughty	1.00
		Runoff limitation	0.40	Slope	0.04	Depth to bedrock	1.00
ScB: Sandose-----	100	Slope	0.04	Filtering capacity	0.00	Too steep for surface application	1.00
		Filtering capacity	0.00			Too steep for sprinkler application	0.22
						Filtering capacity	0.00
SkB: Simeon-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45				
SkD: Simeon-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.21	Droughty	0.21
SvD: Simeon-----	100	Droughty	0.21				
		Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45	Droughty	0.21	Too steep for surface application	0.66
Valentine-----	40	Droughty	0.21			Droughty	0.21
						Too steep for sprinkler application	0.00
Tn: Tryon-----	100	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
To: Tryon-----	100	Droughty	0.82	Droughty	0.82	Droughty	0.82
		Runoff limitation	0.40				
TpB: Tryon-----	55	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty	0.82	Droughty	0.82	Droughty	0.82
		Runoff limitation	0.40				

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Els-----	45	Very limited Filtering capacity Depth to saturated zone Droughty Leaching limitation	1.00 0.95 0.54 0.45	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 0.95 0.54	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 0.95 0.54
VaB: Valentine-----	100	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.88 0.45	Very limited Filtering capacity Droughty	1.00 0.88	Very limited Filtering capacity Droughty	1.00 0.88
VaD: Valentine-----	100	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.88 0.45	Very limited Filtering capacity Droughty	1.00 0.88	Very limited Filtering capacity Droughty Too steep for surface application Too steep for sprinkler application	1.00 0.88 0.66 0.00
VaE: Valentine-----	100	Very limited Slope  Filtering capacity Droughty Leaching limitation	1.00  1.00 0.88 0.45	Very limited Slope  Filtering capacity Droughty	1.00  1.00 0.88	Very limited Too steep for surface application Too steep for sprinkler application Filtering capacity Droughty	1.00  1.00 1.00 0.88
VaF: Valentine-----	70	Very limited Slope  Filtering capacity Droughty Leaching limitation	1.00  1.00 0.88 0.45	Very limited Slope  Filtering capacity Droughty	1.00  1.00 0.88	Very limited Too steep for surface application Too steep for sprinkler application Filtering capacity Droughty	1.00  1.00 1.00 0.88
Valentine-----	30	Very limited Slope  Filtering capacity Droughty Leaching limitation	1.00  1.00 0.88 0.45	Very limited Slope  Filtering capacity Droughty	1.00  1.00 0.88	Very limited Too steep for surface application Too steep for sprinkler application Filtering capacity Droughty	1.00  1.00 1.00 0.88
VbB: Valentine-----	100	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.65 0.45	Very limited Filtering capacity Droughty	1.00 0.65	Very limited Filtering capacity Droughty	1.00 0.65
VbD: Valentine-----	100	Very limited Filtering capacity Droughty  Leaching limitation	1.00 0.65 0.45	Very limited Filtering capacity Droughty	1.00 0.65	Very limited Filtering capacity Too steep for surface application Droughty	1.00 0.66 0.65

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VfD: Valentine-----	55	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.88 0.45	Very limited Filtering capacity Droughty	1.00 0.88	Too steep for sprinkler application	0.00
						Very limited Filtering capacity	1.00
						Droughty	0.88
						Too steep for surface application	0.66
Els-----	45	Very limited Filtering capacity Depth to saturated zone Droughty Leaching limitation	1.00 0.95 0.54 0.45	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 0.95 0.54	Too steep for sprinkler application	0.00
						Very limited Filtering capacity	1.00
						Droughty	0.95
						Depth to saturated zone Droughty	0.54
VhD: Valentine-----	60	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.87 0.45	Very limited Filtering capacity Droughty	1.00 0.87	Too steep for surface application	0.00
						Too steep for sprinkler application	0.00
						Very limited Filtering capacity	1.00
						Droughty	0.87
Libory-----	40	Very limited Depth to saturated zone Filtering capacity Leaching limitation Restricted permeability	1.00 1.00 0.45 0.43	Very limited Depth to saturated zone Filtering capacity Restricted permeability	1.00 1.00 0.32	Too steep for surface application	0.66
						Too steep for sprinkler application	0.00
						Very limited Filtering capacity	1.00
						Droughty	0.82
VpD: Valentine-----	60	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.88 0.45	Very limited Filtering capacity Droughty	1.00 0.88	Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
Pivot-----	40	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.82 0.45	Very limited Filtering capacity Droughty	1.00 0.82	Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
VrD: Valentine-----	65	Very limited Filtering capacity Droughty Leaching limitation	1.00 0.72 0.45	Very limited Filtering capacity Droughty	1.00 0.72	Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
Sandose-----	35	Very limited Filtering capacity Leaching limitation Restricted permeability	1.00 0.45 0.43	Very limited Filtering capacity Restricted permeability	1.00 0.32	Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66
						Too steep for surface application	0.66

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
VsG2: Valentine-----	65	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Too steep for sprinkler application	1.00
		Droughty	0.87	Droughty	0.87	Filtering capacity Droughty	1.00
		Leaching limitation	0.45				0.87
Simeon-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Too steep for sprinkler application	1.00
		Leaching limitation Droughty	0.45 0.19	Droughty	0.19	Filtering capacity Droughty	1.00 0.19
VtE: Valentine-----	60	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Droughty	0.86	Droughty	0.86	Filtering capacity	1.00
		Leaching limitation Slope	0.45 0.16	Slope	0.16	Droughty	0.86
						Too steep for sprinkler application	0.39
Longpine-----	40	Very limited Depth to bedrock Droughty Runoff limitation	1.00 1.00 0.40	Very limited Droughty Depth to bedrock Slope	1.00 1.00 0.16	Very limited Droughty Depth to bedrock Too steep for surface application	1.00 1.00 1.00
		Slope	0.16	Filtering capacity	0.00	Too steep for sprinkler application	0.39
		Filtering capacity	0.00			Filtering capacity	0.00
VwE: Valentine-----	65	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Droughty	0.88	Droughty	0.88	Filtering capacity	1.00
		Leaching limitation Slope	0.45 0.16	Slope	0.16	Droughty	0.88
						Too steep for sprinkler application	0.39
Tryon-----	35	Very limited Depth to saturated zone Filtering capacity Droughty Runoff limitation	1.00 1.00 0.80 0.40	Very limited Depth to saturated zone Filtering capacity Droughty	1.00 1.00 0.80	Very limited Depth to saturated zone Filtering capacity Droughty	1.00 1.00 0.80
VxB: Vetal-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
zwa: Water-----	100	Not rated		Not rated		Not rated	
zwb: Water-----	100	Not rated		Not rated		Not rated	

AGRICULTURAL WASTE MANAGEMENT--Continued  
Brown County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	



In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Brown County, Nebraska

PAGE 2 of 7

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Aa: ALMERIA LOAMY FINE SAND, 0 TO 2 PERCENT SLOPES	ALMERIA	Yes	flood plain	2B3	YES	NO	NO
Ae: ALMERIA FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	ALMERIA	Yes	flood plain	2B3	YES	NO	NO
Af: ALMERIA-HISTOSOLS COMPLEX, CHANNELED	ALMERIA	Yes	flood plain	2B3,3	YES	NO	YES
	HISTOSOLS	Yes	fen	1,3,4	NO	YES	YES
An: ANSELMO FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	ANSELMO	No	flat	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
AnC: ANSELMO FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	ANSELMO	No	hill, valley side	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
AnD: ANSELMO FINE SANDY LOAM, 6 TO 11 PERCENT SLOPES	ANSELMO	No	hill, valley side	---	---	---	---
AtF: ANSELMO-BRUNSWICK FINE SANDY LOAMS, 11 TO 30 PERCENT SLOPES	ANSELMO	No	valley side	---	---	---	---
	BRUNSWICK	No	valley side	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
Ba: BARNEY FINE SANDY LOAM, CHANNELED	BARNEY	Yes	flood plain	2B3	YES	NO	NO
Bd: BOLENT FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	BOLENT	No	flood plain	---	---	---	---
	ALMERIA	Yes	flood plain	2B3	YES	NO	NO
Bo: BROCKSBURG LOAM, 0 TO 1 PERCENT SLOPES	BROCKSBURG	No	flat	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
BrD: BRUNSWICK FINE SANDY LOAM, 3 TO 9 PERCENT SLOPES	BRUNSWICK	No	hill, valley side	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO
DuB: DUNDAY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	DUNDAY	No	hummock, interdune	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
DuD: DUNDAY LOAMY FINE SAND, 3 TO 9 PERCENT SLOPES	DUNDAY	No	dune, interdune	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
Eo: ELS LOAMY SAND, 0 TO 2 PERCENT SLOPES	ELS	No	interdune, swale	---	---	---	---
	MARLAKE	Yes	depression	2B1,3	YES	NO	YES
	TRYON	Yes	swale	2B2,3	YES	NO	YES
EpB: ELS-IPAGE FINE SANDS, 0 TO 3 PERCENT SLOPES	ELS	No	interdune, swale	---	---	---	---
	IPAGE	No	hummock, interdune	---	---	---	---
	MARLAKE	Yes	depression	2B1,3	YES	NO	YES
	TRYON	Yes	swale	2B2,3	YES	NO	YES
Es: ELSMERE LOAMY FINE SAND, 0 TO 2 PERCENT SLOPES	ELSMERE	No	interdune, swale	---	---	---	---
	LOUP	Yes	swale	2B3	YES	NO	NO
Fe: FLUVAQUENTS, SANDY	FLUVAQUENTS	Yes	flood plain	2B2,3	YES	NO	YES



HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Brown County, Nebraska

PAGE 3 of 7

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Gn: GANNETT FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	GANNETT	Yes	interdune, swale	2B3	YES	NO	NO
IdB: INAVALE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	INAVALE	No	flood plain	---	---	---	---
IkB: INAVALE SAND, CHanneLED	ALMERIA FLUVAQUENTS	Yes Yes	flood plain flood plain	2B3 2B1,3	YES YES	NO NO	NO YES
	INAVALE	No	flood plain	---	---	---	---
In: INAVALE-BARNEY COMPLEX, CHanneLED	ALMERIA FLUVAQUENTS	Yes Yes	flood plain flood plain	2B3 2B1,3	YES YES	NO NO	NO YES
	INAVALE	No	flood plain	---	---	---	---
IpB: IPAGE FINE SAND, 0 TO 3 PERCENT SLOPES	BARNEY FLUVAQUENTS	Yes Yes	flood plain flood plain	2B3 3,2B1	YES YES	NO NO	NO YES
	IPAGE	No	hummock, interdune, stream terrace	---	---	---	---
IsB: IPAGE LOAMY SAND, 0 TO 3 PERCENT SLOPES	TRYON	Yes	swale	2B2,3	YES	NO	YES
	IPAGE	No	hummock, interdune, stream terrace	---	---	---	---
Jn: JANSEN FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	TRYON	Yes	swale	2B2,3	YES	NO	YES
	JANSEN	No	flat	---	---	---	---
JnC: JANSEN FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	PERCHED WT	Yes	depression	2A	YES	NO	NO
	JANSEN	No	hill, valley side	---	---	---	---
Jo: JANSEN LOAM, 0 TO 2 PERCENT SLOPES	PERCHED WT	Yes	depression	2A	YES	NO	NO
	JANSEN	No	flat	---	---	---	---
JoC: JANSEN LOAM, 2 TO 6 PERCENT SLOPES	PERCHED WT	Yes	depression	2A	YES	NO	NO
	JANSEN	No	hill, valley side	---	---	---	---
Jr: JANSEN-MEADIN COMPLEX, 0 TO 2 PERCENT SLOPES	PERCHED WT	Yes	depression	2A	YES	NO	NO
	JANSEN	No	knoll	---	---	---	---
JtB: JANSEN-SANDOSE COMPLEX, 0 TO 3 PERCENT SLOPES	MEADIN	No	flat	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
Jw: JOHNSTOWN FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	JANSEN	No	flat	---	---	---	---
	SANDOSE	No	knoll	---	---	---	---
JwB: JOHNSTOWN FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	JOHNSTOWN	No	flat	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
Jy: JOHNSTOWN LOAM, 0 TO 1 PERCENT SLOPES	JOHNSTOWN	No	hill	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
JyB: JOHNSTOWN LOAM, 1 TO 3 PERCENT SLOPES	JOHNSTOWN	No	hill	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
JyC: JOHNSTOWN LOAM, 3 TO 6 PERCENT SLOPES	JOHNSTOWN	No	hill, valley side	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Brown County, Nebraska

PAGE 4 of 7

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
LcG: LABU-SANSARC SILTY CLAYS, 11 TO 40 PERCENT SLOPES	LABU	No	valley side	---	---	---	---
	SANSARC	No	valley side	---	---	---	---
LfB: LIBORY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	LIBORY	No	interdune, swale	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
Lo: LOUP FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	LOUP	Yes	interdune, swale	2B3	YES	NO	NO
Lp: LOUP FINE SANDY LOAM, WET, 0 TO 2 PERCENT SLOPES	LOUP	Yes	interdune, swale	2B3,3	YES	NO	YES
LtB: LOUP-ELS COMPLEX, 0 TO 3 PERCENT SLOPES	ELS	No	hummock, interdune	---	---	---	---
	LOUP	Yes	interdune, swale	2B3	YES	NO	NO
	LOUP MARLAKE	Yes	swale	2B3,3	YES	NO	YES
	MARLAKE	Yes	depression	2B1,3	YES	NO	YES
Ma: MARLAKE FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	MARLAKE	Yes	depression, interdune	2B2,3	YES	NO	YES
McG: MCKELVIE-TASSEL-RONSON COMPLEX, 15 TO 70 PERCENT SLOPES	MCKELVIE	No	valley side	---	---	---	---
	LONGPINE	No	valley side	---	---	---	---
	RONSON	No	valley side	---	---	---	---
	BARNEY	Yes	flood plain	2B3,3	YES	NO	YES
	FLUVAQUENTS	Yes	flood plain	2B1,3	YES	NO	YES
MeB: MEADIN SANDY LOAM, 0 TO 3 PERCENT SLOPES	MEADIN	No	flat, knoll	---	---	---	---
MeF: MEADIN SANDY LOAM, 3 TO 30 PERCENT SLOPES	MEADIN	No	hill, valley side	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
Oe: O'NEILL FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	O'NEILL	No	flat	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
On: O'NEILL LOAM, 0 TO 2 PERCENT SLOPES	O'NEILL	No	flat	---	---	---	---
OsC: O'NEILL-MEADIN SANDY LOAMS, 2 TO 6 PERCENT SLOPES	O'NEILL	No	hill, valley side	---	---	---	---
	MEADIN	No	hill, valley side	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
OsD: O'NEILL-MEADIN SANDY LOAMS, 6 TO 11 PERCENT SLOPES	O'NEILL	No	hill, valley side	---	---	---	---
	MEADIN	No	hill, valley side	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B3	YES	NO	NO
Pg: PITS, SAND AND GRAVEL	PITS	Unranked	---	---	---	---	---
	MARLAKE	Yes	depression	3,2B1	YES	NO	YES
PtB: PIVOT LOAMY SAND, 0 TO 3 PERCENT SLOPES	PIVOT	No	hummock, interdune	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
RtB: RONSON-TASSEL FINE SANDY LOAMS, 0 TO 3 PERCENT SLOPES	RONSON	No	swale	---	---	---	---
	LONGPINE	No	knoll	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Brown County, Nebraska

PAGE 5 of 7

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
RtC: RONSON-TASSEL FINE SANDY LOAMS, 3 TO 6 PERCENT SLOPES	RONSON	No	hill	---	---	---	---
	LONGPINE PERCHED WT	No Yes	hill depression	---	---	---	---
RtD: RONSON-TASSEL FINE SANDY LOAMS, 6 TO 11 PERCENT SLOPES	RONSON	No	hill	---	---	---	---
	LONGPINE PERCHED WT	No Yes	hill depression	---	---	---	---
ScB: SANDOSE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	SANDOSE	No	interdune, swale	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
SkB: SIMEON LOAMY SAND, 0 TO 3 PERCENT SLOPES	SIMEON	No	hummock, interdune	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
SkD: SIMEON LOAMY SAND, 3 TO 9 PERCENT SLOPES	SIMEON	No	hill, valley side	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
SvD: SIMEON-VALENTINE FINE SANDS, 0 TO 9 PERCENT SLOPES	SIMEON	No	interdune, swale	---	---	---	---
	VALENTINE	No	dune, interdune	---	---	---	---
Tn: TRYON LOAMY FINE SAND, 0 TO 2 PERCENT SLOPES	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
	TRYON	Yes	interdune, swale	2B2	YES	NO	NO
To: TRYON LOAMY FINE SAND, WET, 0 TO 2 PERCENT SLOPES	TRYON	Yes	interdune, swale	2B2,3	YES	NO	YES
	TRYON	Yes	interdune, swale	2B2	YES	NO	NO
TpB: TRYON-ELS COMPLEX, 0 TO 3 PERCENT SLOPES	ELS	No	hummock, interdune	---	---	---	---
	MARLAKE	Yes	depression	2B1,3	YES	NO	YES
VaB: VALENTINE FINE SAND, 0 TO 3 PERCENT SLOPES	VALENTINE	No	hummock, interdune	---	---	---	---
	TRYON	Yes	swale	2B2,3	YES	NO	YES
VaD: VALENTINE FINE SAND, 3 TO 9 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	MARLAKE TRYON	Yes Yes	depression swale	2B1,3 2B2,3	YES YES	NO NO	YES YES
VaE: VALENTINE FINE SAND, ROLLING	VALENTINE	No	dune	---	---	---	---
	TRYON	Yes	swale	2B2,3	YES	NO	YES
VaF: VALENTINE FINE SAND, ROLLING AND HILLY	VALENTINE	No	dune	---	---	---	---
	VALENTINE TRYON	No Yes	dune swale	---	---	---	---
VbB: VALENTINE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	VALENTINE	No	hummock, interdune	---	---	---	---
	TRYON	Yes	swale	2B2,3	YES	NO	YES
VbD: VALENTINE LOAMY FINE SAND, 3 TO 9 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	TRYON	Yes	swale	3,2B2	YES	NO	YES
VfD: VALENTINE-ELS FINE SANDS, 0 TO 9 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	ELS	No	interdune, swale	---	---	---	---
	MARLAKE TRYON	Yes Yes	depression swale	2B1,3 2B2,3	YES YES	NO NO	YES YES

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Brown County, Nebraska

PAGE 6 of 7

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
VhD: VALENTINE-LIBORY COMPLEX, 0 TO 9 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	LIBORY	No	interdune, swale	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
VpD: VALENTINE-PIVOT COMPLEX, 0 TO 9 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	PIVOT	No	interdune, swale	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
VrD: VALENTINE-SANDOSE LOAMY FINE SANDS, 0 TO 9 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	SANDOSE	No	interdune, swale	---	---	---	---
	PERCHED WT	Yes	depression	2A	YES	NO	NO
VsG2: VALENTINE-SIMEON COMPLEX, 9 TO 40 PERCENT SLOPES, ERODED	VALENTINE	No	valley side	---	---	---	---
	SIMEON MARLAKE	No Yes	valley side depression	---	---	---	---
				2B1,3	YES	NO	YES
VtE: VALENTINE-TASSEL COMPLEX, 3 TO 17 PERCENT SLOPES	VALENTINE	No	dune	---	---	---	---
	LONGPINE	No	hill	---	---	---	---
VwE: VALENTINE-TRYON COMPLEX, 0 TO 17 PERCENT SLOPES	VALENTINE	No	dune, interdune	---	---	---	---
	TRYON	Yes	interdune, swale	2B2,3	YES	NO	YES
	MARLAKE	Yes	depression	2B1,3	YES	NO	YES
VxB: VETAL LOAM, 1 TO 3 PERCENT SLOPES	VETAL	No	swale, valley side	---	---	---	---
	WT AT 0-1 FOOT	Yes	swale	2B2	YES	NO	NO
zwa: WATER > 40 ACRES	WATER	Unranked	---	---	---	---	---
zwb: WATER < 40 ACRES	WATER	Unranked	---	---	---	---	---

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Brown County, Nebraska

PAGE 7 of 7

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II.

Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
  - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
  - b. poorly drained or very poorly drained and have either:
    - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in), or for other soils
    - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
    - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.



HIGHLY ERODIBLE LANDS REPORT

Survey Area- BROWN COUNTY, NEBRASKA

Map Symbol	Soil Mapunit Name	HEL Classifications		
		C=30	R=100	wnd wat mu
Aa	ALMERIA LOAMY FINE SAND, 0 TO 2 PERCENT SLOPES	3	3	3
Ae	ALMERIA FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Af	ALMERIA-HISTOSOLS COMPLEX, CHANNELED	3	3	3
An	ANSELMO FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
AnC	ANSELMO FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
AnD	ANSELMO FINE SANDY LOAM, 6 TO 11 PERCENT SLOPES	3	2	2
AtF	ANSELMO-BRUNSWICK FINE SANDY LOAMS, 11 TO 30 PERCENT SLOPES	3	2	2
Ba	BARNEY FINE SANDY LOAM, CHANNELED	3	3	3
Bd	BOLENT FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Bo	BROCKSBURG LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
BrD	BRUNSWICK FINE SANDY LOAM, 3 TO 9 PERCENT SLOPES	3	2	2
DuB	DUNDAY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
DuD	DUNDAY LOAMY FINE SAND, 3 TO 9 PERCENT SLOPES	1	2	1
Eo	ELS LOAMY SAND, 0 TO 2 PERCENT SLOPES	1	3	1
EpB	ELS-IPAGE FINE SANDS, 0 TO 3 PERCENT SLOPES	1	3	1
Es	ELSMERE LOAMY FINE SAND, 0 TO 2 PERCENT SLOPES	1	3	1
Fe	FLUVAQUENTS, SANDY	1	3	1
Gn	GANNETT FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
IdB	INAVALE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
IkB	INAVALE SAND, CHANNELED	1	3	1
In	INAVALE-BARNEY COMPLEX, CHANNELED	1	3	1
IpB	IPAGE FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
IsB	IPAGE LOAMY SAND, 0 TO 3 PERCENT SLOPES	1	3	1
Jn	JANSEN FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
JnC	JANSEN FINE SANDY LOAM, 2 TO 6 PERCENT SLOPES	3	3	3
Jo	JANSEN LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
JoC	JANSEN LOAM, 2 TO 6 PERCENT SLOPES	3	2	2
Jr	JANSEN-MEADIN COMPLEX, 0 TO 2 PERCENT SLOPES	2	3	2
JtB	JANSEN-SANDOSE COMPLEX, 0 TO 3 PERCENT SLOPES	2	3	2
Jw	JOHNSTOWN FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
JwB	JOHNSTOWN FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	3	3	3
Jy	JOHNSTOWN LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
JyB	JOHNSTOWN LOAM, 1 TO 3 PERCENT SLOPES	3	3	3
JyC	JOHNSTOWN LOAM, 3 TO 6 PERCENT SLOPES	3	3	3
LcG	LABU-SANSARC SILTY CLAYS, 11 TO 40 PERCENT SLOPES	2	1	1
LfB	LIBORY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
Lo	LOUP FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
Lp	LOUP FINE SANDY LOAM, WET, 0 TO 2 PERCENT SLOPES	3	3	3
LtB	LOUP-ELS COMPLEX, 0 TO 3 PERCENT SLOPES	2	3	2
Ma	MARLAKE FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	3	3	3
McG	MCKELVIE-TASSEL-RONSON COMPLEX, 15 TO 70 PERCENT SLOPES	1	1	1
MeB	MEADIN SANDY LOAM, 0 TO 3 PERCENT SLOPES	1	3	1
MeF	MEADIN SANDY LOAM, 3 TO 30 PERCENT SLOPES	1	2	1
Oe	O'NEILL FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
On	O'NEILL LOAM, 0 TO 2 PERCENT SLOPES	3	3	3
OsC	O'NEILL-MEADIN SANDY LOAMS, 2 TO 6 PERCENT SLOPES	2	3	2
OsD	O'NEILL-MEADIN SANDY LOAMS, 6 TO 11 PERCENT SLOPES	2	2	2
Pg	PITS, SAND AND GRAVEL	3	3	3
PtB	PIVOT LOAMY SAND, 0 TO 3 PERCENT SLOPES	1	3	1
RtB	RONSON-TASSEL FINE SANDY LOAMS, 0 TO 3 PERCENT SLOPES	2	3	2
RtC	RONSON-TASSEL FINE SANDY LOAMS, 3 TO 6 PERCENT SLOPES	2	2	2
RtD	RONSON-TASSEL FINE SANDY LOAMS, 6 TO 11 PERCENT SLOPES	2	2	2
ScB	SANDOSE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
SkB	SIMEON LOAMY SAND, 0 TO 3 PERCENT SLOPES	1	3	1
SkD	SIMEON LOAMY SAND, 3 TO 9 PERCENT SLOPES	1	2	1

SvD	SIMEON-VALENTINE FINE SANDS, 0 TO 9 PERCENT SLOPES	1	3	1
Tn	TRYON LOAMY FINE SAND, 0 TO 2 PERCENT SLOPES	3	3	3
To	TRYON LOAMY FINE SAND, WET, 0 TO 2 PERCENT SLOPES	3	3	3
TpB	TRYON-ELS COMPLEX, 0 TO 3 PERCENT SLOPES	2	3	2
VaB	VALENTINE FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
VaD	VALENTINE FINE SAND, 3 TO 9 PERCENT SLOPES	1	3	1
VaE	VALENTINE FINE SAND, ROLLING	1	2	1
VaF	VALENTINE FINE SAND, ROLLING AND HILLY	1	2	1
VbB	VALENTINE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES	1	3	1
VbD	VALENTINE LOAMY FINE SAND, 3 TO 9 PERCENT SLOPES	1	3	1
VfD	VALENTINE-ELS FINE SANDS, 0 TO 9 PERCENT SLOPES	1	3	1
VhD	VALENTINE-LIBORY COMPLEX, 0 TO 9 PERCENT SLOPES	1	3	1
VpD	VALENTINE-PIVOT COMPLEX, 0 TO 9 PERCENT SLOPES	1	3	1
VrD	VALENTINE-SANDOSE LOAMY FINE SANDS, 0 TO 9 PERCENT SLOPES	1	3	1
VsG2	VALENTINE-SIMEON COMPLEX, 9 TO 40 PERCENT SLOPES, ERODED	1	2	1
VtE	VALENTINE-TASSEL COMPLEX, 3 TO 17 PERCENT SLOPES	1	2	1
VwE	VALENTINE-TRYON COMPLEX, 0 TO 17 PERCENT SLOPES	2	2	2
VxB	VETAL LOAM, 1 TO 3 PERCENT SLOPES	3	3	3